Test yourself: a 49-year-old lady with a history of chronic left hip pain following a fall twenty months earlier

Muhammad Dharas¹ · Petra Balogh² · Michele Calleja¹ · Asif Saifuddin¹

Published online: 14 April 2020
© ISS 2020

Left sided hip pain and reduced range of motion

History: A 49-year-old lady with a history of chronic left hip pain following a fall twenty months earlier Figs. 1, 2 and 3.

Fig. 1 a Coronal STIR and b axial PDW FSE MR images

Skeletal Radiology: Test Yourself Manuscript.
Left sided hip pain and reduced range of motion.

The diagnosis can be found at <https://doi.org/10.1007/s00256-020-03428-6>

✉ Muhammad Dharas
muhammad.dharas@nhs.net

¹ Imaging Department, Royal National Orthopaedic Hospital, Brockley Hill, Stanmore, England HA7 4LP, UK

² Department of Histopathology, Royal National Orthopaedic Hospital, Brockley Hill, Stanmore, England HA7 4LP, UK

Fig. 2 **a** Axial and **b** sagittal MPR CT images

Fig. 3 **a** Microphotograph (H&E, 10x magnification). **b** Polarised light image

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.