



Painless wrist lump

Rupert Berkeley¹ · Daniel Lindsay² · Rob Pollock³ · Asif Saifuddin¹

Received: 27 October 2020 / Revised: 7 December 2020 / Accepted: 13 December 2020 / Published online: 18 January 2021
© ISS 2021

Question

A 17-year-old female patient presented with a 4-week history of a painless, hard lump on the volar aspect of the right wrist. Range of movement, distal neurovascular examination and the overlying skin were normal. There was no history of trauma.



Fig. 1 Lateral radiograph of the right wrist

The diagnosis can be found at <https://doi.org/10.1007/s00256-020-03699-z>.

✉ Rupert Berkeley
rupert.berkeley@nhs.net

¹ Department of Radiology, Royal National Orthopaedic Hospital, Brockley Hill, Stanmore, Middlesex HA7 4LP, UK

² Department of Pathology, Royal National Orthopaedic Hospital, Brockley Hill, Stanmore, Middlesex HA7 4LP, UK

³ Department of Surgery, Royal National Orthopaedic Hospital, Brockley Hill, Stanmore, Middlesex HA7 4LP, UK




Fig. 2 Axial CT study of the right wrist




Fig. 4 Microphotographs, H&E. **a** $\times 4$. **b** $\times 10$




Fig. 3 MRI of the right wrist. **a** Sagittal T1W TSE. **b** Axial fat-suppressed T2W FSE. **c** Sagittal fat-suppressed post-contrast T1W FSE images

Compliance with ethical standards

Conflict of interest The authors declare that they have no conflict of interest.

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.