Enlarging knee masses

Hatem Mawi¹ · Denis Gravel² · Joel Werier³ · Zaid Jibri¹

Received: 16 April 2021 / Revised: 25 June 2021 / Accepted: 30 June 2021 / Published online: 11 July 2021
© ISS 2021

Question: A 67-year-old man presented with an 8-year history of progressive right knee swelling, increasing pain, and limited range of motion. MRI was performed (Fig. 1). Ultrasound-guided biopsy was obtained. Follow-up MRI was performed (Fig. 2), which was followed by surgical resection with the histopathologic examination (Fig. 3).

The diagnosis can be found at doi: 10.1007/s00256-021-03,860-2

✉ Zaid Jibri
zaid.jibri@icloud.com

¹ Department of Medical Imaging, Department of Radiology, The Ottawa Hospital, Ottawa, ON K1H 8L6, Canada

² Division of Anatomical Pathology, The Ottawa Hospital, Ottawa, ON, Canada

³ Department of Surgery, The Ottawa Hospital, Ottawa, ON, Canada

Fig. 1 MRI of the right knee including **a** and **b** sagittal T2 fat suppressed (FS), **c** sagittal proton density (PD), and **d** axial PDFS sequences

Fig. 2 Follow-up (20 months) MRI images of the right knee including **a** sagittal PD, **b** axial T1, and **c** sagittal fat-suppressed T1 post-IV gadolinium

Fig. 3 Photomicrographs including hematoxylin and eosin (**a–c**), as well as immunohistochemical nuclear staining for STAT-6 (**d**)

Declarations

Conflict of interest The authors declare no competing interests.

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.