TEST YOURSELF: QUESTION



Newborn with abnormality of the clavicle

Roberto Franceschi^{1,2} · Giovanna Stringari³ · Fiorenza Soli⁴ · Annalisa Pedrolli¹ · Evelina Maines¹

Received: 29 March 2022 / Revised: 27 May 2022 / Accepted: 13 June 2022 / Published online: 21 June 2022 © The Author(s), under exclusive licence to International Skeletal Society (ISS) 2022

A 32-week gestational age female neonate, born by caesarean section, presented with respiratory distress, and a chest x-ray was performed (Fig. 1). Physical examination showed large fontanelles and open sutures, micrognathia and glossoptosis, facial dysmorphisms, pectus excavatum, and bulging eyes.

At 11 months of corrected age, due to palpable clavicular discontinuity on physical examination, the proband performed a second X-ray (Fig. 2).

Declarations

Ethics approval Informed consent was obtained from all individual participants included in the study. This study was conducted in accordance with the ethical standards of the institutional and/or national research committee and with the 1964 Helsinki declaration and its later amendments or comparable ethical standards.

Conflict of interest The authors declare no competing interests.

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

The diagnosis can be found at https://doi.org/10.1007/s00256-022-04095-5

Roberto Franceschi roberto.franceschi@apss.tn.it

- ¹ Pediatric Department, S.Chiara Hospital of Trento, Trento, Italy
- ² Division of Paediatrics, S. Chiara General Hospital, Largo Medaglie d'Oro 9, 38122 Trento, Italy
- ³ Neonatology Unit, S.Chiara Hospital of Trento, Trento, Italy
- ⁴ Genetic Unit, S.Chiara Hospital of Trento, Trento, Italy



Fig. 1 Chest x-ray of the propositus at birth

Fig. 2 Chest x-ray of the propositus at 11 months of corrected age

