

Effects of ROI definition and reconstruction method on quantitative outcome and applicability in a response monitoring trial

Nanda C. Krak¹, R. Boellaard¹, Otto S. Hoekstra¹, Jos W. R. Twisk², Corneline J. Hoekstra¹, Adriaan A. Lammertsma¹

¹ Clinical PET Centre, VU University Medical Centre, Amsterdam, The Netherlands

² Department of Clinical Epidemiology and Biostatistics, VU University Medical Centre, Amsterdam, The Netherlands

Published online: 26 August 2005

© Springer-Verlag 2005

Eur J Nucl Med Mol Imaging (2005) 32:1245

DOI 10.1007/s00259-005-1926-5

Eur J Nucl Med (2005) 32:294–301

The following acknowledgement should have been part of the original article:

Acknowledgement. The authors would like to thank Dr. Johan Nuyts from the Department of Nuclear Medicine, University Hospital Gasthuisberg, Leuven, Belgium, for providing the semi-automatic program that was used to draw the three-dimensional regions of interest.

The online version of the original article can be found at
<http://dx.doi.org/10.1007/s00259-004-1566-1>

Adriaan A. Lammertsma (✉)
Clinical PET Centre,
VU University Medical Centre,
Amsterdam, The Netherlands