

Direct comparison of [18F]FDG PET/CT with PET alone and with side-by-side PET and CT in patients with malignant melanoma

Felix M. Mottaghy · Cord Sunderkötter ·
Roland Schubert · Petra Wohlfart ·
Norbert M. Blumstein · Bernd Neumaier ·
Gerhard Glatting · Cueneyt Özdemir ·
Andreas K. Buck · Karin Scharffetter-Kochanek ·
Sven N. Reske

Published online: 1 June 2007
© Springer-Verlag 2007

Eur J Nucl Med Mol Imaging (2007)34:XX

Unfortunately the surname of one of the co-authors, Karin Scharffetter-Kochanek, was rendered incorrectly in the original publication.

The online version of the original article can be found at <http://dx.doi.org/10.1007/s00259-006-0358-1>.

F. M. Mottaghy (✉) · P. Wohlfart · N. M. Blumstein ·
B. Neumaier · G. Glatting · A. K. Buck · S. N. Reske
Department of Nuclear Medicine, University Hospital Ulm,
Robert-Koch-Strasse 8,
89073 Ulm, Germany
e-mail: felix.mottaghy@uz.kuleuven.ac.be

C. Sunderkötter · R. Schubert · C. Özdemir ·
K. Scharffetter-Kochanek
Department of Dermatology, University Hospital Ulm,
Ulm, Germany

C. Sunderkötter
Department of Dermatology, University Hospital Münster,
Munster, Germany