

Mean and maximum standardized uptake values in [^{18}F]FDG-PET for assessment of histopathological response in oesophageal squamous cell carcinoma or adenocarcinoma after radiochemotherapy

Matthias Schmidt · Elfriede Bollschweiler · Markus Dietlein · Stefan P. Mönig · Carsten Kobe · Daniel Vallböhmer · Wolfgang Eschner · Arnulf Hölscher · Harald Schicha

Published online: 10 February 2009
© Springer-Verlag 2009

Erratum to: Eur J Nucl Med Mol Imaging
DOI 10.1007/s00259-008-1011-y

The correct name of the sixth author is Daniel Vallböhmer, not Vallboehmer as originally published.”

The online version of the original article can be found at <http://dx.doi.org/10.1007/s00259-008-1011-y>.

M. Schmidt (✉) · M. Dietlein · C. Kobe · W. Eschner · H. Schicha
Department of Nuclear Medicine, University of Cologne,
Kerpener Str. 62,
50937 Cologne, Germany
e-mail: Matthias.Schmidt@uni-koeln.de

E. Bollschweiler · S. P. Mönig · D. Vallböhmer · A. Hölscher
Department of General-, Visceral and Cancer Surgery,
University of Cologne,
Cologne, Germany