Correction to: Tumor fragmentation estimated by volume surface ratio of tumors measured on ^{18}F -FDG PET/CT is an independent prognostic factor of diffuse large B-cell lymphoma

Pierre Decazes^{1,2} • Stéphanie Becker^{1,2} • Mathieu Nessim Toledano^{1,2} • Pierre Vera^{1,2} • Paul Desbordes^{1,2} • Fabrice Jardin^{3,4} • Hervé Tilly^{3,4} • Isabelle Gardin^{1,2}

Published online: 26 May 2018

© Springer-Verlag GmbH Germany, part of Springer Nature 2018

Correction to: European Journal of Nuclear Medicine and Molecular Imaging

<https://doi.org/10.1007/s00259-018-4041-0>

A unit error concerning the tumor volume surface ratio (TVSR) is present throughout the article. The unit reported is "cm" but is actually "mm".

This error is present in the abstract, the figures 1 and 2, the legend of figure 1, the table 2 (line of TVSR), in the results part (paragraphs 'PET/CT metrics and correlations' and 'ROC curve analysis') and in the discussion.

Correct figures are presented here.

Fig. 1

The online version of the original article can be found at <https://doi.org/10.1007/s00259-018-4041-0>

✉ Pierre Decazes
pierre.decazes@chb.unicancer.fr

¹ Department of Nuclear Medicine, Henri Becquerel Cancer Centre, Rue d'Amiens – CS 11516, 76038 Rouen Cedex 1, France

² LITIS Quantif-EA4108, University of Rouen, Rouen, France

³ Department of Haematology, Henri Becquerel Cancer Centre, Rouen, France

⁴ INSERM U918, Centre Henri Becquerel, Rouen, France

Fig. 2