Correction to: Staging and quantification of florbetaben PET images using machine learning: impact of predicted regional cortical tracer uptake and amyloid stage on clinical outcomes

Jun Pyo Kim^{1,2,3} · Jeonghun Kim⁴ · Yeshin Kim⁵ · Seung Hwan Moon⁶ · Yu Hyun Park^{1,2} · Sole Yoo⁷ · Hyemin Jang^{1,2,3} · Hee Jin Kim^{1,2,3} · Duk L. Na^{1,2,3,8} · Sang Won Seo^{1,2,3,9,10} · Joon-Kyung Seong^{4,11}

Published online: 10 February 2020

© Springer-Verlag GmbH Germany, part of Springer Nature 2020

Correction to: European Journal of Nuclear Medicine and Molecular Imaging

<https://doi.org/10.1007/s00259-019-04663-3>

The Table 2 in the original version of this article contained a mistake in the alignment. Correct Table 2 presentation is presented here.

The original article has been corrected.

This article is part of the Topical Collection on Erratum 2.

The online version of the original article can be found at <https://doi.org/10.1007/s00259-019-04663-3>

✉ Sang Won Seo
sangwonseo@empal.com

✉ Joon-Kyung Seong
jkseong@korea.ac.kr

¹ Department of Neurology, Samsung Medical Center, Seoul, South Korea

² Samsung Alzheimer Research Center, Samsung Medical Center, Seoul, South Korea

³ Neuroscience Center, Samsung Medical Center, Seoul, South Korea

⁴ Department of Bio-convergence Engineering, Korea University, Seoul, South Korea

⁵ Department of Neurology, Kangwon National University Hospital, Chuncheon, South Korea

⁶ Department of Nuclear Medicine, Samsung Medical Center, Seoul, South Korea

⁷ Department of Cognitive Science, Yonsei University, Seoul, South Korea

⁸ Department of Health Sciences and Technology, SAIHST, Sungkyunkwan University, Seoul, South Korea

⁹ Department of Clinical Research Design & Evaluation, SAIHST, Sungkyunkwan University, Seoul, South Korea

¹⁰ Center for Clinical Epidemiology, Samsung Medical Center, Seoul, South Korea

¹¹ School of Biomedical Engineering, Korea University, Seoul, South Korea

Table 2 Results from linear regression analysis between mean pRCTU and neuropsychological/structural parameters

A. Structural parameters																		
	Frontal cortex		Parietal cortex		Temporal cortex		Occipital cortex		Global cortex		Hippocampal volume							
	<i>B</i> (SE)	<i>p</i>	Adj. <i>R</i> ²	<i>B</i> (SE)	<i>p</i>	Adj. <i>R</i> ²	<i>B</i> (SE)	<i>p</i>	Adj. <i>R</i> ²	<i>B</i> (SE)	<i>p</i>	Adj. <i>R</i> ²						
All subjects	-0.042 (0.010)	<0.001*	0.196	-0.082 (0.012)	<0.001*	0.238	-0.092 (0.012)	<0.001*	0.279	-0.068 (0.012)	<0.001*	0.215	-0.066 (0.010)	<0.001*	0.261	-598.9 (75.7)	<0.001*	0.298
Subjects in stage 1 and stage 2	-0.035 (0.029)	0.232	0.127	-0.087 (0.035)	0.014*	0.142	-0.0115 (0.036)	0.002*	0.137	-0.062 (0.033)	0.06	0.089	-0.067 (0.029)	0.023*	0.134	-264.5 (214.5)	0.219	0.104
B. Neuropsychological parameters																		
	K-MMSE		Attention		Language		Visuospatial		Memory		Frontal/executive							
	<i>B</i> (SE)	<i>p</i>	Adj. <i>R</i> ²	<i>B</i> (SE)	<i>p</i>	Adj. <i>R</i> ²	<i>B</i> (SE)	<i>p</i>	Adj. <i>R</i> ²	<i>B</i> (SE)	<i>p</i>	Adj. <i>R</i> ²	<i>B</i> (SE)	<i>p</i>	Adj. <i>R</i> ²			
All subjects	-4.266 (0.363)	<0.001*	0.345	-0.858 (0.171)	<0.001*	0.223	-0.382 (0.056)	<0.001*	0.142	-0.718 (0.085)	<0.001*	0.252	-0.739 (0.056)	<0.001*	<0.001*	0.362 (0.071)	-0.721	<0.001*
Subjects in stage 1 and stage 2	-4.994 (1.081)	<0.001*	0.205	-1.692 (0.436)	<0.001*	0.246	-0.707 (0.169)	<0.001*	0.08	-1.143 (0.255)	<0.001*	0.197	-0.656 (0.134)	<0.001*	0.164	-0.900 (0.194)	<0.001*	0.212

Models with neuropsychological outcomes included age, sex, and years of education as covariates. Models with structural outcome variables included age, sex, and total intracranial volume as covariates
pRCTU/predicted regional cortical tracer uptake, *K-MMSE* Korean version of mini-mental status examination, *Adj. R²* adjusted *R* squared