



Correction to: Superior performance of 18F-fluorocholine digital PET/CT in the detection of parathyroid adenomas

Diego Alfonso López-Mora¹ · Marina Sizova¹ · Montserrat Estorch¹ · Albert Flotats¹ · Valle Camacho¹ · Alejandro Fernández¹ · Safae Abouzian¹ · Francisco Fuentes-Ocampo¹ · José Ignacio Pérez García² · Ana Isabel Chico Ballesteros^{3,4} · Joan Duch¹ · Anna Domènech¹ · Antonio Moral Duarte² · Ignasi Carrió¹

Published online: 16 March 2020
© Springer-Verlag GmbH Germany, part of Springer Nature 2020

Correction to: European Journal of Nuclear Medicine and Molecular Imaging (2020) 47:572–578

<https://doi.org/10.1007/s00259-020-04680-7>

The materials and methods section of the original version of this article contained a mistake regarding the intravenous administered activity of 18F-fluorocholine (18F-FCH). The correct administered activity of 18F-FCH is 0.1–0.14 mCi/Kg, which is equivalent to 3.7–5.2 MBq/kg.

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

This article is part of the Topical Collection on Erratum

The online version of the original article can be found at <https://doi.org/10.1007/s00259-020-04680-7>

✉ Diego Alfonso López-Mora
dlopezmo@santpau.cat

¹ Nuclear Medicine Department, Hospital de la Santa Creu i Sant Pau, Universitat Autònoma de Barcelona, Pare Claret 167, 08025 Barcelona, Spain

² General Surgery Department, Hospital de la Santa Creu i Sant Pau, Universitat Autònoma de Barcelona, Barcelona, Spain

³ Endocrinology Department, Hospital de la Santa Creu i Sant Pau, Universitat Autònoma de Barcelona, Barcelona, Spain

⁴ Centro de Investigación Biomédica en Red de Bioingeniería, Biomateriales y Nanomedicina (CIBER-BBN), Madrid, Spain