



## Correction to: $^{68}\text{Ga}$ -FAPI outperforms $^{18}\text{F}$ -FDG PET/CT in identifying solitary fibrous tumor

Rongxi Wang<sup>1,2</sup> · Qingxing Liu<sup>1,2</sup> · Huimin Sui<sup>1,2</sup> · Mu Zhang<sup>3</sup> · Zhaohui Zhu<sup>1,2</sup> · Ruixue Cui<sup>1,2</sup>

Published online: 3 March 2021

© Springer-Verlag GmbH Germany, part of Springer Nature 2021

### Correction to: Eur J Nucl Med Mol Imaging

<https://doi.org/10.1007/s00259-020-05181-3>

There was a mismatch of text and image in the original article. The figure shown in the original article was the revised version, but the text part was the original version, so the figure legends in the text do not correspond to the figure. The correct figure is shown in the next page.

The original article has been corrected.

---

This article is part of the Topical Collection on Erratum.

The online version of the original article can be found at <https://doi.org/10.1007/s00259-020-05181-3>


✉ Zhaohui Zhu  
13611093752@163.com

✉ Ruixue Cui  
mmdhmm@126.com

<sup>1</sup> Department of Nuclear Medicine, Peking Union Medical College Hospital, Chinese Academy of Medical Sciences & Peking Union Medical College, Beijing 100730, China

<sup>2</sup> Beijing Key Laboratory of Molecular Targeted Diagnosis and Therapy in Nuclear Medicine, Beijing 100730, China

<sup>3</sup> Department of Pathology, Peking Union Medical College Hospital, Chinese Academy of Medical Sciences & Peking Union Medical College, Beijing 100730, China



**Publisher's note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.