Correction to: Diagnostic performance of preoperative MR imaging findings for differentiation of uterine leiomyoma with intraligamentous growth from subserosal leiomyoma

Ryo Yajima¹ · Aki Kido¹ · Ryo Kuwahara² · Yusaku Moribata¹ · Yoshitsugu Chigusa³ · Yuki Himoto¹ · Yasuhisa Kurata¹ · Yuka Matsumoto¹ · Satoshi Otani¹ · Naoko Nishio¹ · Sachiko Minamiguchi⁴ · Masaki Mandai³ · Yuji Nakamoto¹

Published online: 16 July 2021

© Springer Science+Business Media, LLC, part of Springer Nature 2021

Correction to: Abdominal Radiology

<https://doi.org/10.1007/s00261-021-03042-7>

The original version of this article unfortunately contained a mistake in Fig. 7.

The correct Fig. 7 is given below.

Fig. 7 Receiver operating characteristic (ROC) curves of the number of positive imaging findings for intraligamentous leiomyoma among four imaging findings: tumor shape deformity, lower attachment to the uterus, ovary elevation, and separation of round ligament and uterine artery

The original article can be found online at <https://doi.org/10.1007/s00261-021-03042-7>.

✉ Aki Kido
akikido@kuhp.kyoto-u.ac.jp

¹ Department of Diagnostic Imaging and Nuclear Medicine, Kyoto University Graduate School of Medicine, 54 Kawaharacho, Shogoin, Sakyo-ku, Kyoto 606-8507, Japan

² Department of Radiology, National Hospital Organization Kyoto Medical Center, 1-1 Mukaihatacho, Fushimi-ku, Kyoto 612-8555, Japan

³ Department of Gynecology and Obstetrics, Kyoto University Graduate School of Medicine, 54 Kawaharacho, Shogoin, Sakyo-ku, Kyoto 606-8507, Japan

⁴ Department of Diagnostic Pathology, Kyoto University Graduate School of Medicine, 54 Kawaharacho, Shogoin, Sakyo-ku, Kyoto 606-8507, Japan

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.