




## Correction to: The gallbladder: what's new in 2022?

Rachel Runde<sup>1</sup> · Edward D. Auyang<sup>2</sup> · Raye Ng<sup>2</sup> · Kaysey Llorente<sup>2</sup> · Hina Arif Tiwari<sup>3</sup> · Shana Elman<sup>1</sup> · William M. Thompson<sup>1</sup> 

Published online: 30 April 2022  
© Springer Science+Business Media, LLC, part of Springer Nature 2022

**Correction to: Abdominal Radiology**  
<https://doi.org/10.1007/s00261-022-03429-0>

The original version of this article unfortunately contained a mistake in figure 8. The correct Fig. 8 is given below.

---

The original article can be found online at <https://doi.org/10.1007/s00261-022-03429-0>.


---

✉ William M. Thompson  
wthompson@salud.UNM.edu

<sup>1</sup> Department of Radiology, University of New Mexico, MSC10 5530, 1 University of New Mexico, Albuquerque, NM 87131, USA

<sup>2</sup> Department of Surgery, University of New Mexico, Albuquerque, NM, USA

<sup>3</sup> Department of Radiology, University of Arizona, Tucson, AZ, USA



**Fig. 8** Chronic cholecystitis in a 40-year-old man with intermittent biliary colic for 7 years. **a** and **b**. CT (**a**) and single image from one of seven ultrasounds (**b**) the patient had prior and subsequent to the HIDA scan, all of which showed a normal gallbladder, with no gallstones, sludge, gallbladder wall thickening, or pericholecystic fluid. **c**. Anterior planar images following injection of <sup>99m</sup>Tc-Mebrofenin

show normal visualization of gallbladder within the first 60 minutes of imaging, excluding acute cholecystitis. Gallbladder ejection fraction obtained after a fatty meal challenge (Boost Plus) was low at 22%, suggestive of biliary dyskinesia/chronic cholecystitis. Subsequent cholecystectomy confirmed chronic cholecystitis in the absence of cholelithiasis

**Publisher's Note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.