



Correction to: LILRB2-containing small extracellular vesicles from glioblastoma promote tumor progression by promoting the formation and expansion of myeloid-derived suppressor cells

Peitao Wu¹ · Yuhang Guo² · Li Xiao¹ · Jiaqi Yuan¹ · Chao Tang³ · Jun Dong¹ · Zhiyuan Qian¹

Published online: 17 April 2023
© The Author(s) 2023

Correction to: Cancer Immunology, Immunotherapy
<https://doi.org/10.1007/s00262-023-03395-6>

The original version of this article unfortunately contained a mistake. The corrected details are given below for your reading.

Equal contributor statement was missing. Equal contributor statement should be "Yuhang Guo is the co-first author".

The statistical result in Fig. 2B has some mistakes.

The statistical analysis was not marked in Fig. 3A, C and F, and the statistical analysis result of Fig. 3B was misplaced.

The corrected Figs. 2 and 3 are given in the following page.

The original article has been corrected.

The original article can be found online at <https://doi.org/10.1007/s00262-023-03395-6>.

✉ Chao Tang
Chaotang@fudan.edu.cn

✉ Jun Dong
Dongjun@suda.edu.cn

✉ Zhiyuan Qian
zhiyuanqian-sz@outlook.com

Peitao Wu
peitaowuu@163.com

Yuhang Guo
guoyh0713@163.com

Li Xiao
xiaoli0107@suda.edu.cn; a1055335273@163.com

¹ Department of Neurosurgery, The Second Affiliated Hospital of Soochow University, Soochow 215000, People's Republic of China

² Department of Neurosurgery, The First Affiliated Hospital of Wenzhou Medical University, Nanbaixiang, Wenzhou, China

³ Department of Neurosurgery, Huashan Hospital, Shanghai, China

Fig. 2 Pirb promotes GBM progression through an immunosuppressive TME. **A** The proliferation of GL261-LV-nc, GL261-pirb⁺, GL261-nc and GL261-pirb⁻ cells detected by CCK8 assays at 24, 48 and 72 h. Statistical analysis of cell apoptosis **B**, migration **C** and invasion **D** in GL261-LV-nc, GL261-pirb⁺, GL261-nc and GL261-pirb⁻ cells. The presence of MDSCs **E**, Tregs **F**, CD4⁺ and CD8⁺ T cells **G** in human GBM tissue detected by IF. * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$, or **** $P < 0.0001$, ns indicates no statistical significance





Fig. 3 High pirb expression leads to immunosuppression. The percentages of MDSCs **A**, Tregs **B**, CD8⁺ T cells **C** and CD4⁺ cells **D** in the tumor sites detected by flow cytometry. The percentages of CD8⁺ T cells **E**, MDSCs **F**, CD4⁺ T cells **G** and Treg cells **H** in the spleens of tumor-bearing mice detected by flow cytometry. * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$, or **** $P < 0.0001$, ns indicates no statistical significance



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in

the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.