

## **Erratum to: Relationship of serum IL-6, C-reactive protein, erythrocyte sedimentation rate, and knee skin temperature after total knee arthroplasty: a prospective study**

**Sittisak Honsawek · Benjamad Deepaisarnsakul · Aree Tanavalee ·  
Manoon Sakdinakiattikoon · Srihatach Ngarmukos · Kanok Preativatanyou ·  
Piyanuch Bumrunghanichthaworn**

Received: 9 July 2011 / Accepted: 9 July 2011 / Published online: 22 July 2011  
© Springer-Verlag 2011

**Erratum to: Int Orthop 2011 Jan; 35 (1): 31-35**  
**DOI 10.1007/s00264-010-0973-0**

We regret that the value of serum CRP was reported incorrectly. Serum CRP levels were reported incorrectly to be 10 times higher than their true value. The value of serum CRP in osteoarthritis patients following total knee arthroplasty should have been reported in milligrams per litre (mg/l), instead of milligrams per decilitre (mg/dl) for the whole article. The authors apologise for this error.

---

The online version of the original article can be found at <http://dx.doi.org/10.1007/s00264-010-0973-0>.

---

S. Honsawek (✉) · B. Deepaisarnsakul · A. Tanavalee ·  
M. Sakdinakiattikoon · S. Ngarmukos · K. Preativatanyou ·  
P. Bumrunghanichthaworn  
Faculty of Medicine, Chulalongkorn University, King  
Chulalongkorn Memorial Hospital, Thai Red Cross Society,  
1873 Rama IV road,  
Patumwan, Bangkok 10330, Thailand  
e-mail: [sittisak.h@chula.ac.th](mailto:sittisak.h@chula.ac.th)