ERRATUM

Erratum to: Causes for early and late surgical re-intervention after radial head arthroplasty

Pierre Laumonerie¹ • David Ancelin¹ • Nicolas Reina¹ • Meagan E. Tibbo² •
Panagiotis Kerezoudis² • Stephanie Delclaux¹ • Nicolas Bonneville¹ • Pierre Mansat¹

Published online: 27 May 2017
© SICOT aisbl 2017

Erratum to: International Orthopaedics (SICOT)
DOI: 10.1007/s00264-017-3496-0

Figure 2 in the original publication contains a few mistakes. The radial head prosthesis designs in Fig. 2a, b are Evolutive® (Aston Medical, Saint-Etienne, France) and Guepar® (Aston Medical, Saint Etienne France) respectively. Below is the appropriate correction:

Fig. 2 Three bipolar radial head prostheses: Evolutive® (a) (Aston Medical, Saint-Etienne, France), and Guepar® (b) (Aston Medical, Saint-Etienne, France) are implants with a long, cemented stems anchored beyond the neck; the rHead® Recon implant (c) (Stryker-Small Bone

Innovation, Morrisville, Pennsylvania, USA) has a short, press-fit stem for high neck fixation and possible cementation if initial stability is not obtained

The online version of the original article can be found at <http://dx.doi.org/10.1007/s00264-017-3496-0>

✉ Pierre Laumonerie
laumonerie.pierre@hotmail.fr

¹ Hôpital Pierre-Paul Riquet, Place du Docteur Baylac,
31059 Toulouse, France

² Mayo Clinic, Rochester, MN 55905, USA