

## Erratum to: Individual voice recognition and an auditory map of neighbours in free-ranging black howler monkeys (*Alouatta pigra*)

Margarita Briseño-Jaramillo<sup>1</sup> · Alejandro Estrada<sup>2</sup> · Alban Lemasson<sup>3,4</sup>

Published online: 7 April 2015  
© Springer-Verlag Berlin Heidelberg 2015

**Erratum to: Behavioral Ecology and Sociobiology 69(1): 13–25**  
**DOI 10.1007/s00265-014-1813-9**

We apologize for a mistake in Table 7. Speed listed in Table 7 (values for “Approach to speaker - Speed” in coherent situation) was erroneously calculated. The correct values are for coherent situation:  $0.07 \pm 0.05$  and for incoherent situation:  $0.23 \pm 0.19$ . Results of GLMM are now  $Z=6.1$ ,  $P=0.004$ .

---

The online version of the original article can be found at <http://dx.doi.org/10.1007/s00265-014-1813-9>.

✉ Margarita Briseño-Jaramillo  
mar.briseno.jaramillo@gmail.com

<sup>1</sup> Instituto de Biología, Universidad Nacional Autónoma de México, Coyoacán, 04510 México City, Mexico

<sup>2</sup> Estación de Biología Tropical Los Tuxtlas, Instituto de Biología, Universidad Nacional Autónoma de México, Veracruz, Mexico

<sup>3</sup> Ethologie Animale et Humaine, U.M.R. 6552-C.N.R.S., Université de Rennes 1, Station Biologique, 35380 Paimpont, France

<sup>4</sup> Institut Universitaire de France, Maison des Universités, 103, Boulevard Saint-Michel, 75005 Paris, France