Dipyron Increases the Blood Flow of Arterial Dorsal Skin Flaps

S. E. Gülmez · S. Serel · A. Uluç · Z. Can ·
H. Ergün

Published online: 1 May 2008
© Springer Science+Business Media, LLC 2008

Erratum to: Aesth Plast Surg DOI 10.1007/s00266-008-9126-x

In the online and printed version of this article, the following errors occurred: the sentence “Dipyron significantly decreases skin blood flow at the acute phase of flap elevation although the necrotic area does not reach the significance level.” located in the Conclusion section of the Abstract is incorrect. The correct sentence appears below.

Dipyron significantly increases skin blood flow at the acute phase of flap elevation although the necrotic area does not reach the significance level.

In addition, the sentence “According to the described results, we conclude that dipyron significantly decreases skin blood flow, observed at the acute phase of flap elevation.” located in the Conclusion section of the article is inaccurate. The accurate sentence is shown below.

According to the described results, we conclude that dipyron significantly increases skin blood flow, observed at the acute phase of flap elevation.

The online version of the original article can be found under doi:
[10.1007/s00266-008-9126-x](https://doi.org/10.1007/s00266-008-9126-x).

S. E. Gülmez · S. Serel (✉) · A. Uluç · Z. Can · H. Ergün
Departments of Pharmacology and Clinical Pharmacology, and
Plastic and Reconstructive Surgery, Ankara University Faculty
of Medicine, Ankara 06590, Turkey
e-mail: savassarel@hotmail.com; savassarel@yahoo.com