

## Erratum to: Glasgow Prognostic Score and Prognosis After Hepatectomy for Hepatocellular Carcinoma

Tomoyuki Abe<sup>1</sup> · Hirotaka Tashiro<sup>2</sup> · Tsuyoshi Kobayashi<sup>1</sup> ·  
Minoru Hattori<sup>3</sup> · Shintaro Kuroda<sup>1</sup> · Hideki Ohdan<sup>1</sup>

Published online: 8 March 2017  
© Société Internationale de Chirurgie 2017

**Erratum to: World J Surg**  
**DOI 10.1007/s00268-017-3909-7**

In the original article, there are several errors in Table 2.  
Following is the corrected Table 2:

---

The online version of the original article can be found under doi:[10.1007/s00268-017-3909-7](https://doi.org/10.1007/s00268-017-3909-7).

---

✉ Hirotaka Tashiro  
tashiroh@kure-nh.go.jp; htashiro@hiroshima-u.ac.jp

Tomoyuki Abe  
t.abe.hiroshima@gmail.com

Tsuyoshi Kobayashi  
tsukoba@hiroshima-u.ac.jp

Minoru Hattori  
m-hattori@hiroshima-u.ac.jp

Shintaro Kuroda  
df26@smn.enjoy.ne.jp

Hideki Ohdan  
hohdan@hiroshima-u.ac.jp

<sup>1</sup> Department of Gastroenterological and Transplant Surgery, Applied Life Sciences, Institute of Biomedical and Health Sciences, Hiroshima University, Hiroshima, Japan

<sup>2</sup> Department of Surgery, Kure Medical Center, National Hospital Organization, 3-1 Aoyama, Kure, Hiroshima 737-0023, Japan

<sup>3</sup> Advanced Medical Skills Training Center, Institute of Biomedical and Health Sciences, Hiroshima University, Hiroshima, Japan

**Table 2** Univariate and multivariate analyses of prognostic factors for recurrence-free survival (RFS)

Characteristic	Patients, <i>n</i> (%)	Univariate analysis			Multivariate analysis		
		3-year RFS (%)	5-year RFS (%)	<i>P</i> value	HR	95% CI	<i>P</i> value
Sex				0.715			
M	336 (74.2)	50.0	36.2				
F	117 (25.8)	48.3	33.2				
Median age (years)				0.943			
<69	221 (48.8)	46.9	35.0				
≥69	232 (51.2)	52.2	35.6				
Median BMI (kg/m <sup>2</sup> )				0.738			
<25.0	340 (75.1)	50.1	36.2				
≥25.0	113 (24.9)	47.3	32.1				
Hepatitis C				0.002*			0.007*
Y	266 (58.7)	44.1	26.3		1.48	1.11–1.96	
N	187 (41.3)	57.6	49.6		1		
Hepatitis B				0.359			
Y	79 (17.4)	55.3	42.2				
N	374 (82.6)	48.2	33.7				
Child-Pugh				0.178			
Grade A	427 (94.3)	50.6	35.7				
Grade B	26 (5.7)	27.8	27.8				
GPS				0.040*			0.208
0	339 (74.8)	54.4	35.9		1		
1–2	114 (25.2)	35.8	32.8		1.20	0.90–1.59	
Median NLR				0.654			
<1.96	224 (49.8)	53.4	37.7				
≥1.96	226 (50.2)	51.9	39.6				
AFP levels (normal range) (ng/mL)				0.003*			0.109
<20.0	271 (59.8)	55.3	40.0		1		
≥20.0	182 (40.2)	41.1	28.6		1.24	0.95–1.61	
DCP levels (normal range) (mAU/nL)				0.008*			0.004*
<40.0	224 (49.4)	56.3	38.6		1		
≥40.0	229 (50.6)	42.5	32.2		1.48	1.14–1.92	
T-Bil levels (normal range) (mg/dL)				0.035*			0.038*
<1.2	402 (88.7)	51.1	37.5		1		
≥1.2	51 (11.3)	35.7	17.3		1.48	1.02–2.14	
AST levels (normal range) (IU/I)				<0.001*			0.075
<32	202 (44.6)	58.7	44.2		1		
≥32	251 (55.4)	42.2	28.6		1.35	0.97–1.89	
ALT levels (normal range) (IU/I)				0.001*			0.670
<37	275 (60.7)	54.8	40.5		1		
≥37	178 (39.3)	41.9	28.4		1.07	0.78–1.47	
PT (normal range) (%)				0.464			
<70	22 (4.90)	43.1	16.2				
≥70	431 (95.1)	49.7	35.9				
ICGR15 (normal range) (%)				0.056			
<10	124 (27.4)	57.4	43.3				
≥10	329 (72.6)	46.3	32.2				

**Table 2** continued

Characteristic	Patients, <i>n</i> (%)	Univariate analysis			Multivariate analysis		
		3-year RFS (%)	5-year RFS (%)	<i>P</i> value	HR	95% CI	<i>P</i> value
Alb levels (normal range) (g/dL)				0.285			
<3.5	66 (14.6)	57.0	38.0				
≥3.5	387 (85.4)	59.4	33.3				
Total ischemic time (mins)				0.366			
<57	226 (50.2)	54.2	42.4				
≥57	224 (49.8)	51.8	35.6				
Median blood loss (mL)				0.094			
<250	219 (48.3)	54.8	39.2				
≥250	234 (51.7)	44.1	31.4				
Median operation time (mins)				0.650			
<275	223 (49.2)	50.9	37.2				
≥275	230 (50.8)	48.0	33.1				
Tumor number				<0.001*			<0.001*
1	318 (70.2)	56.5	41.6		1		
>1	135 (29.8)	33.7	21.6		1.81	1.39–2.35	
Maximum tumor diameter (mm)				0.973			
<26	225 (49.7)	50.9	31.4				
≥26	228 (50.3)	48.1	39.3				
Hepatic resection				0.347			
Hr0 (limited resection)	183 (40.4)	45.2	32.3				
HrS (segmentectomy)	139 (30.7)	58.5	40.1				
Hr1 (sectionectomy)	82 (18.1)	45.3	33.8				
Hr2 (hemihepatectomy)	49 (10.8)	45.1	37.2				
Histological grade				0.124			
Well/moderately differentiated	374 (82.6)	47.4	33.2				
Poorly differentiated/other	79 (17.4)	58.7	44.8				
MVI				0.212			
Y	71 (15.7)	43.2	37.8				
N	382 (84.3)	50.7	35.2				

*Alb* albumin, *AFP* alpha-fetoprotein, *ALT* alanine aminotransferase, *AST* aspartate aminotransferase, *BMI* body mass index, *CI* confidence interval, *DCP* des-gamma carboxyprothrombin, *F* female, *GPS* Glasgow Prognostic Score, *HR* hazard ratio, *Hr* hepatic resection, *ICGR15* indocyanine green retention rate at 15 minutes, *M* male, *MVI* microvascular invasion, *N* no, *NLR* neutrophil-to-lymphocyte ratio, *PT* prothrombin time, *T-Bil* total bilirubin, *Y* yes

\*  $P < 0.05$