



CORRECTION

Correction to: Hypoparathyroidism After Total Thyroidectomy: Importance of the Intraoperative Management of the Parathyroid Glands

Guillermo Ponce de León-Ballesteros¹ · David Velázquez-Fernández¹ ·
F. Javier Hernández-Calderón¹ · Carlos Bonilla-Ramírez¹ · Rafael H. Pérez-Soto¹ ·
Juan Pablo Pantoja¹ · Mauricio Sierra¹ · Miguel F. Herrera¹

Published online: 15 April 2019
© Société Internationale de Chirurgie 2019

Correction to: World J Surg
<https://doi.org/10.1007/s00268-019-04987-z>

In the original version of the article, the last three column headings in Table 3 were mislabeled. The original article has been corrected. Following is the corrected table:

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Table 3 Rate of postoperative hypocalcemia, protracted and permanent hypoparathyroidism according PGRIS score

Group	<i>n</i> (%)	Postoperative hypocalcemia <i>n</i> (%)	Protracted hypoparathyroidism <i>n</i> (%)	Permanent hypoparathyroidism <i>n</i> (%)
1 PGRIS score 4	649 (67.9)	193 (29.7)	99 (15.3)	16 (2.5)
2 PGRIS score 3	209 (21.9)	68 (32.5)	41 (19.6)	8 (3.8)
3 PGRIS score 1–2	98 (10.3)	53 (54.1)	36 (36.7)	13 (13.3)
		$p \leq 0.0001^*$	$p \leq 0.0001^*$	$p \leq 0.0001^*$

**P* value Pearson's Chi-square test. Comparison between group rates

The original article can be found online at <https://doi.org/10.1007/s00268-019-04987-z>.

✉ Miguel F. Herrera
miguelherrera@gmail.com

¹ Service of Endocrine and Advanced Laparoscopic Surgery/
Department of Surgery, Instituto Nacional de Ciencias
Médicas y Nutrición Salvador Zubiran, Vasco de Quiroga 15,
Seccion XVI, Tlalpan, 14000 Mexico City, Mexico