

Erratum to: Subclavian Vein Versus Arm Vein for Totally Implantable Central Venous Port for Patients with Head and Neck Cancer: A Retrospective Comparative Analysis

Akio Akahane · Miyuki Sone · Shigeru Ehara ·
Kenichi Kato · Ryoichi Tanaka · Tatsuhiko Nakasato

Published online: 26 May 2011

© Springer Science+Business Media, LLC and the Cardiovascular and Interventional Radiological Society of Europe (CIRSE) 2011

Erratum to: Cardiovasc Intervent Radiol
DOI 10.1007/s00270-010-0051-4

This erratum is being published to correct the error of the article entitled “Subclavian Vein Versus Arm Vein for Totally Implantable Central Venous Port for Patients with Head and Neck Cancer: A Retrospective Comparative Analysis” DOI:[10.1007/s00270-010-0051-4](https://doi.org/10.1007/s00270-010-0051-4), Editorial manuscript number: CVR-2010-0478.R2, as follows. Table 3 of the summary of adverse event should read as follows.

The online version of the original article can be found under
doi:[10.1007/s00270-010-0051-4](https://doi.org/10.1007/s00270-010-0051-4).

A. Akahane (✉) · M. Sone · S. Ehara · K. Kato · R. Tanaka ·
T. Nakasato
Department of Radiology, Iwate Medical University School
of Medicine, 19-1 Uchimaru, Morioka 020-8505, Japan
e-mail: a.akahane@gmail.com

Table 3 Adverse events

Adverse event	Subclavian ports		Arm ports		<i>P</i> Value
	No. (%)	Median duration (days)	No. (%)	Median duration (days)	
Periprocedural					
Pneumothorax	1 (2.1)		0 (0)		0.290
Arterial puncture	1 (2.1)		0 (0)		0.290
Primary malposition	0 (0)		0 (0)		
Total	2 (4.3)		0 (0)		0.830
Postprocedural					
Infection**	4 (8.5)	49	9 (7.8)	44.5	0.758
Phlebitis	0 (0)		11 (9.5)	33	0.035*
Fibrin sheath	0 (0)		0 (0)		
System occlusion	0 (0)		5 (4.3)	53	0.323
Subcutaneous extravasation	0 (0)		1 (0.9)	34	1.000
Venous thrombosis	0 (0)		0 (0)		
Catheter detachment	0 (0)		0 (0)		
Total	4 (8.5)	49	26 (22.6)	66	0.044*

CVP central venous port

* $P < 0.05$

** Details of infectious AEs are listed in Table 4