## CORRECTION



## Correction to: The Current Use of Drug-Eluting Balloons and Stents in Peripheral Arterial Disease: An Online Survey by the Cardiovascular and Interventional Radiological Society of Europe (CIRSE)

Robert A. Morgan $^1$  · Stefan Müller-Hülsbeck $^2$  · Fabrizio Fanelli $^3$  · Patrick Haage $^4$  · Mohamad Hamady $^5$  · Romaric Loffroy $^6$  · Gerard O'Sullivan $^7$  · Florian Wolf $^8$  · Birgit Slijepčević $^9$ 

Published online: 1 November 2023

© Springer Science+Business Media, LLC, part of Springer Nature and the Cardiovascular and Interventional Radiological Society of Europe (CIRSE) 2023

## **Correction to: Cardiovasc Intervent Radiol**

https://doi.org/10.1007/s00270-023-03562-3

The original article was published without the following visual abstract and has been corrected:

The original article can be found online at https://doi.org/10.1007/s00270-023-03562-3.

⊠ Robert A. Morgan rmorgan@sgul.ac.uk

Stefan Müller-Hülsbeck muehue@diako.de

Fabrizio Fanelli fabrizio.fanelli@unifi.it

Patrick Haage patrick.haage@helios-gesundheit.de

Mohamad Hamady mhamady@imperial.ac.uk

Romaric Loffroy romaric.loffroy@chu-dijon.fr

Gerard O'Sullivan lahinchman1923@gmail.com

Florian Wolf florian.wolf@meduniwien.ac.at

Birgit Slijepčević slijepcevic@cirse.org


- St George's University of London, London, UK
- <sup>2</sup> Academic Hospital Christian-Albrechts-University Kiel, Kiel, Germany

- <sup>3</sup> Vascular and Interventional Radiology Department, "Careggi" University Hospital – University of Florence, Florence, Italy
- <sup>4</sup> Helios University Hospital, University Witten/Herdecke, Wuppertal, Germany
- 5 Imperial College, St Mary's Campus, London, UK
- Department of Diagnostic and Interventional Radiology, University of Burgundy, Dijon, France
- University Hospital Galway, Galway, Ireland
- Division of Cardiovascular and Interventional Radiology, Medical University of Vienna, Vienna, Austria
- 9 CIRSE Central Office, Vienna, Austria



R. A. Morgan, S. Müller-Hülsbeck, F. Fanelli, P. Haage, M. Hamady, R. Loffroy, G. O'Sullivan, F. Wolf, B. Slijepčević

The Current Use of Drug-Eluting Balloons and Stents in Peripheral Arterial Disease: An Online Survey by the Cardiovascular and Interventional Radiological Society of Europe (CIRSE)



The 2018 meta-analysis had an impact on the daily IR practice with drug-eluting devices in PAD of a majority of respondents. Meanwhile, with more available evidence and recent statements, the use of drug-eluting devices is considered as safe and is increasing again for a majority of respondents.

**Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not

included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit http://creativecommons.org/licenses/by/4.0/.

**Publisher's Note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

