Correction to: Wnt/ β -catenin signaling mediates the suppressive effects of diallyl trisulfide on colorectal cancer stem cells

Qi Zhang¹ · Xiao-Ting Li^{1,2} · Yue Chen¹ · Jia-Qi Chen¹ · Jian-Yun Zhu¹ · Yu Meng¹ · Xiao-Qian Wang¹ · Yuan Li¹ · Shan-Shan Geng^{1,2} · Chun-Feng Xie^{1,2} · Jie-Shu Wu^{1,2} · Cai-Yun Zhong^{1,2} · Hong-Yu Han³

Published online: 30 April 2018

© Springer-Verlag GmbH Germany, part of Springer Nature 2018

Correction to: Cancer Chemotherapy and Pharmacology
<https://doi.org/10.1007/s00280-018-3565-0>

Unfortunately, the online published article has error in Fig. 4. The correct Fig. 4 is given here.

The original article can be found online at <https://doi.org/10.1007/s00280-018-3565-0>.

✉ Cai-Yun Zhong
cyzhong@njmu.edu.cn

✉ Hong-Yu Han
hanhy@sysucc.org.cn

¹ Department of Nutrition and Food Safety, School of Public Health, Nanjing Medical University, 818 East Tianyuan Rd, Jiangning, Nanjing 211166, China

² The Key Laboratory of Modern Toxicology, Ministry of Education, School of Public Health, Nanjing Medical University, Nanjing 211166, China

³ Department of Clinical Nutrition, Sun Yat-Sen University Cancer Center, State Key Laboratory of Oncology in South China, Collaborative Innovation Center for Cancer Medicine, Guangzhou 510060, China

Fig. 4 DATS inhibits Wnt/ β -catenin pathway in colorectal CSCs. The tumorspheres enriched from SW480 and DLD-1 were treated with different concentrations of DATS for 5 days. **a**, **b** SW480 and DLD-1 colonospheres were treated with DATS for 5 days. The expression levels of the critical molecules in Wnt/ β -catenin pathway were determined by qRT-PCR (**a**) and Western blotting (**b**), respectively. **c–f** SW480 and DLD-1 sphere-forming cells were cultured with 40 μ M DATS in the presence or absence of 20 mM LiCl for 5 days, and the protein levels for Wnt/ β -catenin pathway were detected by

Western blotting (**c**). **d** The images of tumorspheres were obtained. Bar 100 μ m. **e** Western blot analysis of colorectal CSCs markers. **f** SW480 cells were transfected with control vector or β -catenin plasmids, and then treated with or without DATS (40 μ M) for 3 days. The levels for β -CATENIN, CD133 and ALDH1 were detected by Western blotting. Data are expressed as mean $\pm$ SD of three independent experiments. * P < 0.05, ** P < 0.01 compared with control group. # P < 0.05, ## P < 0.01 compared with 40 μ M DATS group