Correction to: Cyclooxygenase activity mediates colorectal cancer cell resistance to the omega-3 polyunsaturated fatty acid eicosapentaenoic acid

Milene Volpato¹ · Nicola Ingram¹ · Sarah L Perry¹ · Jade Spencer² · Amanda D Race² · Catriona Marshall¹ · John M Hutchinson¹ · Anna Nicolaou^{3,4} · Paul M Loadman² · P Louise Coletta¹ · Mark A Hull¹

Published online: 17 November 2020
© Springer-Verlag GmbH Germany, part of Springer Nature 2020

Correction to: Cancer Chemotherapy and Pharmacology
<https://doi.org/10.1007/s00280-020-04157-2>

In the original publication of the article, curves, more data point and lines were missing in the Fig. 2A. The corrected Fig. 2 is given below. The original article has been corrected.

The original article can be found online at <https://doi.org/10.1007/s00280-020-04157-2>.

✉ Milene Volpato
m.volpato@leeds.ac.uk

¹ Leeds Institute of Medical Research at St James's, University of Leeds, St James's University Hospital, Leeds LS9 7TF, UK

² Institute of Cancer Therapeutics, University of Bradford, Bradford BD7 1DP, UK

³ Laboratory for Lipidomics and Lipid Biology, Division of Pharmacy and Optometry, School of Health Sciences, University of Manchester, Manchester M13 9PT, UK

⁴ Lydia Becker Institute of Immunology and Inflammation, University of Manchester, Manchester M13 9PT, UK

Fig. 2 COX expression mediates EPA sensitivity of CRC cells in vitro. **a** MTT cell viability assay curves for the effect of EPA on MC38 (closed circles) and MC38r mouse CRC cells (open circles). Data represent the mean $\pm$ SEM % growth compared to untreated control cells for three independent replicates for each mouse CRC cell line. **b** Western blot analysis of COX isoform protein expression in MC38 and MC38r mouse CRC cells. **c** PGE_2 levels in medium conditioned by MC38 and MC38r cells for 24 h. Data compared with Student's t test. **d** Western blot analysis of COX isoform protein

expression in CT26 and COX^{low} -CT26 mouse CRC cells. **e** PGE_2 levels in medium conditioned by CT26 and COX^{low} -CT26 mouse CRC cells for 24 h. Data compared with Student's t test. **f** MTT cell viability assay curves for the effect of EPA on CT26 (closed circles) and COX^{low} -CT26 mouse CRC cells (open circles). Data represent the mean $\pm$ SEM % growth compared to untreated control cells for three independent replicates for each mouse CRC cell line. Two-way ANOVA confirmed a significant increase in EPA sensitivity of COX^{low} -CT26 cells compared with CT26 mouse CRC cells ($p = 0.03$)