Two Definite Integrals That Are Definitely (and Surprisingly!) Equal

SHALOSH B. EKHAD, DORON ZEILBERGER,
AND WADIM ZUDILIN

This column is a place for those bits of contagious mathematics that travel from person to person in the community, because they are so elegant, surprising, or appealing that one has an urge to pass them on.

Contributions are most welcome.

To our good friend Gert Almkvist (1934–2018)
In Memoriam.

PROPOSITION 1. *For real $a > b > 0$ and nonnegative integer n , the following beautiful and surprising identity holds:*

$$\int_0^1 \frac{x^n(1-x)^n}{((x+a)(x+b))^{n+1}} dx = \int_0^1 \frac{x^n(1-x)^n}{((a-b)x + (a+1)b)^{n+1}} dx.$$

PROOF. Fix a and b , let $L(n)$ and $R(n)$ be the integrals on the left and right sides respectively, and let $F_1(n, x)$ and $F_2(n, x)$ be the corresponding integrands, so that $L(n) = \int_0^1 F_1(n, x) dx$ and $R(n) = \int_0^1 F_2(n, x) dx$. We cleverly construct the rational functions

$$R_1(x) = \frac{x(x-1)((a+b+1)x^2 + 2abx - ab)}{(x+b)(x+a)}$$

and

$$R_2(x) = \frac{x(x-1)((a-b)x^2 + 2b(a+1)x - (a+1)b)}{(a-b)x - (a+1)b}$$

motivated by the fact that (check!)

$$\begin{aligned} & (n+1)F_1(n, x) - (2n+3)(2ba+a+b)F_1(n+1, x) \\ & + (a-b)^2(n+2)F_1(n+2, x) \\ & = \frac{d}{dx}(R_1(x)F_1(n, x)) \end{aligned}$$

and

$$\begin{aligned} & (n+1)F_2(n, x) - (2n+3)(2ba+a+b)F_2(n+1, x) \\ & + (a-b)^2(n+2)F_2(n+2, x) \\ & = \frac{d}{dx}(R_2(x)F_2(n, x)). \end{aligned}$$

Integrating both identities from $x = 0$ to $x = 1$ and noting that the right-hand sides vanish, we have

$$\begin{aligned} & (n+1)L(n) - (2n+3)(2ba+a+b)L(n+1) \\ & + (a-b)^2(n+2)L(n+2) = 0 \end{aligned}$$

and

$$\begin{aligned} & (n+1)R(n) - (2n+3)(2ba+a+b)R(n+1) \\ & + (a-b)^2(n+2)R(n+2) = 0. \end{aligned}$$

Since $L(0) = R(0)$ and $L(1) = R(1)$ (check!), the proposition follows by mathematical induction. $\square$

➤ Submissions should be uploaded to <http://tmin.edmgr.com> or sent directly to **Sergei Tabachnikov**, e-mail: tabachni@math.psu.edu

REMARK 1. This beautiful identity is equivalent to an identity buried in Bailey's classic book [3, Section 9.5, formula (2)], but you need an expert (like the third-named author) to realize that!

REMARK 2. Our proof was obtained by the first-named author by running a Maple program¹ written by the second-named author that implements the Almkvist–Zeilberger algorithm [2] designed by Zeilberger and our good mutual friend Gert Almkvist, to whose memory this note is dedicated.

REMARK 3. Our integrals are not taken from a pool of no-one-cares-about-them analytic creatures: the right-hand side covers a famous sequence of rational approximations to $\log(1 + (a - b)/((a + 1)b))$ [1], and hence the left-hand side does, too.

REMARK 4. We thank Greg Egan for spotting a sign error in an earlier version.

REMARK 5. Watch Greg Egan's beautiful animation.²

REMARK 6. To our surprise, the identity turned out to be not as surprising as we had believed. Mikael Sundquist noticed that the change of variable $x = b(1 - u)/(b + u)$ gives a “Calculus 1 proof.” Indeed,

$$dx = -\frac{b(1+b)}{(b+u)^2} du,$$

and we have

$$\begin{aligned} & \int_0^1 \frac{x^n(1-x)^n}{((x+a)(x+b))^{n+1}} dx \\ &= \int_0^1 \frac{(b(1-u)/(b+u))^n(1-(b(1-u)/(b+u)))^n}{(b(1-u)/(b+u)+a)(b(1-u)/(b+u)+b)^{n+1}} \\ & \quad \times \frac{b(1+b)}{(b+u)^2} du \\ &= \int_0^1 \frac{(1-u)^n u^n b^{n+1} (1+b)^{n+1}}{(b(1-u)+a(b+u))(b(1-u)+b(b+u))^{n+1}} du \\ &= \int_0^1 \frac{u^n(1-u)^n}{((a-b)u+(a+1)b)^{n+1}} du. \end{aligned}$$

Note that n does not have to be an integer.

REMARK 7. Alin Bostan has two further insightful proofs.³

OPEN ACCESS

This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

Shalosh B. Ekhad
Department of Mathematics
Rutgers University (New Brunswick)
Hill Center-Busch Campus, 110 Frelinghuysen Rd.
Piscataway, NJ 08854-8019
USA
e-mail: ShaloshBEkhad@gmail.com

Doron Zeilberger
Department of Mathematics
Rutgers University (New Brunswick)
Hill Center-Busch Campus, 110 Frelinghuysen Rd.
Piscataway, NJ 08854-8019
USA
website: <https://sites.math.rutgers.edu/~zeilberg/>

Wadim Zudilin
Institute for Mathematics, Astrophysics and Particle Physics
Radboud University
PO Box 9010, 6500 GL Nijmegen
The Netherlands
website: <https://www.math.ru.nl/~zudilin/>

REFERENCES

- [1] K. Alladi and M. L. Robinson. Legendre polynomials and irrationality. *J. Reine Angew. Math.* 318 (1980), 137–155.
- [2] Gert Almkvist and Doron Zeilberger. The method of differentiating under the integral sign. *J. Symbolic Computation* 10 (1990), 571–591.
- [3] W. N. Bailey. *Generalized Hypergeometric Series*. Cambridge University Press, 1935.

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

¹Available online at <http://sites.math.rutgers.edu/~zeilberg/tokhniot/EKHAD.txt>.

²Available online at <https://twitter.com/gregeganSF/status/1192309179119104000>.

³See <https://specfun.inria.fr/bostan/publications/EZZ2.pdf> and <https://specfun.inria.fr/bostan/publications/EZZ2.pdf>.