

Erratum

Extranuclear Expression of the Bacterial Xylose Isomerase (*xylA*) and the UDP-Glucose Dehydrogenase (*hasB*) Genes in Yeast with *Kluyveromyces lactis* Linear Killer Plasmids as Vectors

Jürgen Schründer,¹ Norio Gunge,² Friedhelm Meinhardt¹

¹Institut für Mikrobiologie, Westfälische Wilhelms-Universität, Corrensstr. 3, 48149 Münster, Germany

²Applied Microbial Technology, Kumamoto Institute of Technology, 4-22-1 Ikeda, Kumamoto 860, Japan

Re: Curr Microbiol 33(5) 323–330 (1996). Please note that in the published version of this article, Figure 4 was misprinted. The correct version of this figure, along with its caption, appears below.




Fig. 4. Genetic organization of the *Saccharomyces cerevisiae* linear hybrid plasmid pRSC128 encoding the UDP glucose dehydrogenase gene (*hasB*) from *Streptococcus pyogenes*. As for the xylose isomerase (see Fig. 2), the presence of the *hasB* encoding hybrid plasmid in the recombinant yeast is demonstrated by a Southern blot with *hasB* as a probe and the native *Kluyveromyces lactis* killer plasmids as a negative control (A). A schematic diagram of the recombinant plasmid with the integrated *LEU2** selectable marker and the heterologous *hasB* gene is presented in B. The aminoterminal modification, resulting in addition of two amino acids to the polypeptide, is shown in part C.