CORRECTION



## Correction to: Matrix group structure and Markov invariants in the strand symmetric phylogenetic substitution model

Peter D. Jarvis<sup>1</sup> · Jeremy G. Sumner<sup>1</sup>

Accepted: 28 May 2021 / Published online: 8 June 2021 © Springer-Verlag GmbH Germany, part of Springer Nature 2021

## Correction to: J. Math. Biol. (2016) 73:259–282 https://doi.org/10.1007/s00285-015-0951-7

The original version of this article was published on 11 December 2015 in Volume 73, issue 2, page 259–282, contained a mistake in the Abstract. The abstract gave a formula for the numbers of such invariants for this model, for L taxa, of quadratic and cubic degree, which contained a pre factor  $\frac{1}{3}$  which was a typographical error.

The sentence in Abstract which previously read as

"In particular, in the quadratic and cubic cases we prove there are precisely  $\frac{1}{3}(3^{L} + (-1)^{L})$  and  $6^{L-1}$  linearly independent Markov invariants, respectively."

Should read as

"In particular, in the quadratic and cubic cases we prove that there are precisely  $\frac{1}{2}(3^{L} + (-1)^{L})$  and  $6^{L-1}$  linearly independent Markov invariants, respectively."

**Publisher's Note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

The original article can be found online at https://doi.org/10.1007/s00285-015-0951-7.

Jeremy G. Sumner Jeremy.Sumner@utas.edu.au

<sup>&</sup>lt;sup>1</sup> School of Physical Sciences, University of Tasmania, Private Bag 37, GPO, Hobart, TAS 7001, Australia