




Correction to: A prospective, longitudinal monocentric study on laser Doppler imaging of microcirculation: comparison with macrovascular pathophysiology and effect of adalimumab treatment in early rheumatoid arthritis

László Dávida¹ · Vanda Pongrácz¹ · Emir Awad Mohamed¹ · Szilvia Szamosi² · Gabriella Szücs² · Andrea Váncsa² · Orsolya Tímár^{1,2} · Zoltán Csiki³ · Edit Végh² · Pál Soltész^{1,4} · Zoltán Szekanez²  · György Kerekes¹

Published online: 6 January 2020
© Springer-Verlag GmbH Germany, part of Springer Nature 2020

Correction to: Rheumatology International
<https://doi.org/10.1007/s00296-019-04503-5>

In the Original article, the legend of Figs. 3 and 4 are interchanged and line number 387 is incomplete in the discussion section. The correct Figs. 3 and 4 legends and the complete sentence are given below:

The complete line in the discussion section is “Our study may have important limitations. The limitations include the small number of patients in the second, pilot part of our study.”

Publisher’s Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations

The original article can be found online at <https://doi.org/10.1007/s00296-019-04503-5>.

✉ Zoltán Szekanez
szekanez.zoltan@med.unideb.hu

¹ Intensive Care Unit, Faculty of Medicine, University of Debrecen, Debrecen, Hungary

² Department of Rheumatology, Faculty of Medicine, University of Debrecen, Nagyerdei str 98, 4032 Debrecen, Hungary

³ Division of Clinical Immunology, Department of Medicine, Faculty of Medicine, University of Debrecen, Debrecen, Hungary

⁴ Division of Angiology, Department of Medicine, Faculty of Medicine, University of Debrecen, Debrecen, Hungary




Fig. 3 Changes in macrovascular endothelial dysfunction (flow-mediated vasodilation) and von Willebrand factor antigen levels upon 12 weeks of adalimumab treatment in patients with early rheumatoid arthritis




Fig. 4 Changes in time to half after hyperemia (TH2) of the postocclusive reactive hyperemia curve representing microcirculation upon 12 weeks of adalimumab treatment in patients with early rheumatoid arthritis