

# Erratum to: Intravenous injection of gadobutrol in an epidemiological study group did not lead to a difference in relative signal intensities of certain brain structures after 5 years

Marie-Luise Kromrey<sup>1</sup> · Kim Rouven Liedtke<sup>1</sup> · Till Ittermann<sup>2</sup> · Sönke Langner<sup>1</sup> · Michael Kirsch<sup>1</sup> · Werner Weitschies<sup>3</sup> · Jens-Peter Kühn<sup>1</sup>

Published online: 21 June 2016  
© European Society of Radiology 2016

**Erratum to: Eur Radiol**  
**DOI 10.1007/s00330-016-4418-z**

Unfortunately, the authors have discovered an error in the specified dosage of the contrast agent Gadobutrol. The correct dosage is 0.15 mmol/kg and not 1.5 mmol/kg, as indicated. The authors sincerely apologise for this error.

---

The online version of the original article can be found at <http://dx.doi.org/10.1007/s00330-016-4418-z>.

---

✉ Marie-Luise Kromrey  
[marie-luise.kromrey@uni-greifswald.de](mailto:marie-luise.kromrey@uni-greifswald.de)

<sup>1</sup> Institute of Diagnostic Radiology and Neuroradiology, University Medicine Greifswald, Ferdinand-Sauerbruch-Straße, D-17475 Greifswald, Germany

<sup>2</sup> Institute for Community Medicine, University Medicine Greifswald, Greifswald, Germany

<sup>3</sup> Institute of Biopharmacy and Pharmaceutical Technology, University Greifswald, Greifswald, Germany