



Correction to: Magnetic resonance with diffusion-weighted imaging improves assessment of focal liver lesions in patients with potentially resectable pancreatic cancer on CT

Sun Kyung Jeon^{1,2} · Jeong Min Lee^{1,2,3} · Ijin Joo^{1,2} · Dong Ho Lee^{1,2} · Su Joa Ahn^{1,2} · Hyunsik Woo⁴ · Myoung Seok Lee⁴ · Jin-Young Jang⁵ · Joon Koo Han^{1,2,3}

Published online: 19 April 2018
© European Society of Radiology 2018

Correction to: Eur Radiol

<https://doi.org/10.1007/s00330-017-5258-1>

The original version of this article, published on 19 January 2018, unfortunately contained a mistake. The following correction has therefore been made in the original:

A sentence in the second paragraph of the discussion was worded incorrectly. The correct wording is “Thus, additional MRI with DWI for the assessment of liver metastasis may not be necessary for this group.”

The online version of the original article can be found at <https://doi.org/10.1007/s00330-017-5258-1>

✉ Jeong Min Lee
jmsh@snu.ac.kr

- ¹ Department of Radiology, Seoul National University Hospital, 101 Daehangno, Jongno-gu, Seoul 03080, Korea
- ² Seoul National University College of Medicine, Seoul, Korea
- ³ Institute of Radiation Medicine, Seoul National University Medical Research Center, Seoul, Korea
- ⁴ Department of Radiology, Seoul Metropolitan Government – Seoul National University Boramae Medical Center, Seoul, Korea
- ⁵ Department of Surgery, Seoul National University Hospital, Seoul, Korea