Correction to: Volumetric dynamic oxygen-enhanced MRI (OE-MRI): comparison with CT Brody score and lung function in cystic fibrosis patients

K. Martini^{1,2} · C. M. Gyga² · C. Benden³ · A. R. Morgan^{4,5} · G. J. M. Parker^{4,5} · T. Frauenfelder^{1,2}

© European Society of Radiology 2018

Correction to: Eur Radiol

<https://doi.org/10.1007/s00330-018-5383-5>

The original version of this article, published on 13 April 2018, unfortunately contained a mistake. The following correction has therefore been made in the original: The % predicted FEV1 in the caption of Fig. 1 was incorrect. The correct value should be 98% and not 27%. The corrected figure is given below.

The online version of the original article can be found at <https://doi.org/10.1007/s00330-018-5383-5>

✉ K. Martini
katharina.martini@usz.ch

¹ Institute for Diagnostic and Interventional Radiology, University Hospital Zurich, Rämistrasse 100, 8091 Zurich, Switzerland

² University of Zurich, Zurich, Switzerland

³ Division of Pulmonology, University Hospital Zurich, Zurich, Switzerland

⁴ Bioxydyn Limited, Manchester, UK

⁵ Centre for Imaging Science, The University of Manchester, Manchester, UK

Fig. 1 Dynamic OE-MRI parameter maps (**a–c**) and respective histograms (**d–f**) from a healthy volunteer (a man, 31 years old, FEV₁ %predicted = 98%). ΔPO_{2max} [**a** and **d** (mmHg)], wash-in time [**b** and **e**

(min)] as well as wash-out time [**c** and **f** (min)] are illustrated. The healthy volunteer shows relatively homogeneous parameter maps for ΔPO_{2max} , wash-in and wash-out values