



# Correction to: T1 mapping for liver function evaluation in gadoxetic acid-enhanced MR imaging: comparison of look-locker inversion recovery and B1 inhomogeneity-corrected variable flip angle method

Ji Eun Kim<sup>1</sup> · Hyun Ok Kim<sup>2</sup> · Kyungsoo Bae<sup>1</sup> · Dae Seob Choi<sup>1</sup> · Dominik Nickel<sup>3</sup>

Published online: 29 July 2019  
© European Society of Radiology 2019

**Correction to: European Radiology (2019) 29:3584–3594**  
<https://doi.org/10.1007/s00330-018-5947-4>

The original version of this article, published on 22 March 2019, unfortunately contained a mistake. The following correction has therefore been made in the original: Affiliations 1 and 2 were presented incorrectly. The corrected version is given below.

**Publisher's note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

---

The online version of the original article can be found at <https://doi.org/10.1007/s00330-018-5947-4>

---

✉ Hyun Ok Kim  
galleonok@hanmail.net

<sup>1</sup> Department of Radiology, Gyeongsang National University College of Medicine and Gyeongsang National University Hospital, 79 Gangnam-ro, Jinju 52727, Republic of Korea

<sup>2</sup> Department of Internal Medicine, Gyeongsang National University College of Medicine and Gyeongsang National University Hospital, 79 Gangnam-ro, Jinju 52727, Republic of Korea

<sup>3</sup> Siemens Healthineers, Erlangen, Germany