Correction to: Vessel-selective 4D MRA based on ASL might potentially show better performance than 3D TOF MRA for treatment evaluation in patients with intra-extracranial bypass surgery: a prospective study

Maoxue Wang¹ · Yongbo Yang² · Yi Wang² · Ming Li¹ · Jilei Zhang³ · Bing Zhang^{1,4}

Published online: 10 March 2021

© European Society of Radiology 2021

Correction to: European Radiology

<https://doi.org/10.1007/s00330-020-07503-3>

The original version of this article, published on 02 January 2021, unfortunately contained a mistake. The following correction has therefore been made in the original: The information that Maoxue Wang and Yongbo Yang contributed equally to this work was missing. The original article has been corrected.

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

The online version of the original article can be found at <https://doi.org/10.1007/s00330-020-07503-3>

✉ Bing Zhang
zhangbing_nanjing@nju.edu.cn; zhangbing_nanjing@vip.163.com

¹ Department of Radiology, The Affiliated Drum Tower Hospital of Nanjing University Medical School, 321 Zhongshan Road, Drum Tower District, Nanjing 210002, Jiangsu Province, China

² Department of Neurosurgery, The Affiliated Drum Tower Hospital of Nanjing University Medical School, Nanjing, Jiangsu, China

³ Philips Healthcare, Shanghai, China

⁴ Institute of Brain Science, Nanjing University, Nanjing, Jiangsu, China