



# Membranous urethral length measurement on preoperative MRI to predict incontinence after radical prostatectomy: a literature review towards a proposal for measurement standardization

Thierry N. Boellaard<sup>1</sup> · Margriet C. van Dijk-de Haan<sup>1</sup> · Stijn W. T. P. J. Heijmink<sup>1</sup> · Corinne N. Tillier<sup>2</sup> · Hans Veerman<sup>2,3</sup> · Laura S. Mertens<sup>2</sup> · Henk G. van der Poel<sup>2,3</sup> · Pim J. van Leeuwen<sup>2</sup> · Ivo G. Schoots<sup>1,4</sup>

Received: 26 February 2023 / Revised: 29 May 2023 / Accepted: 7 July 2023 / Published online: 22 September 2023

© The Author(s) 2023

## Abstract

**Objectives** To investigate the membranous urethral length (MUL) measurement and its interobserver agreement, and propose literature-based recommendations to standardize MUL measurement for increasing interobserver agreement. MUL measurements based on prostate MRI scans, for urinary incontinence risk assessment before radical prostatectomy (RP), may influence treatment decision-making in men with localised prostate cancer. Before implementation in clinical practise, MRI-based MUL measurements need standardization to improve observer agreement.

**Methods** Online libraries were searched up to August 5, 2022, on MUL measurements. Two reviewers performed article selection and critical appraisal. Papers reporting on preoperative MUL measurements and urinary continence correlation were selected. Extracted information included measuring procedures, MRI sequences, population mean/median values, and observer agreement.

**Results** Fifty papers were included. Studies that specified the MRI sequence used T2-weighted images and used either coronal images ( $n=13$ ), sagittal images ( $n=18$ ), or both ( $n=12$ ) for MUL measurements. ‘Prostatic apex’ was the most common description of the proximal membranous urethra landmark and ‘level/entry of the urethra into the penile bulb’ was the most common description of the distal landmark. Population mean (median) MUL value range was 10.4–17.1 mm (7.3–17.3 mm), suggesting either population or measurement differences. Detailed measurement technique descriptions for reproducibility were lacking. Recommendations on MRI-based MUL measurement were formulated by using anatomical landmarks and detailed descriptions and illustrations.

**Conclusions** In order to improve on measurement variability, a literature-based measuring method of the MUL was proposed, supported by several illustrative case studies, in an attempt to standardize MRI-based MUL measurements for appropriate urinary incontinence risk preoperatively.

**Clinical relevance statement** Implementation of MUL measurements into clinical practise for personalized post-prostatectomy continence prediction is hampered by lack of standardization and suboptimal interobserver agreement. Our proposed standardized MUL measurement aims to facilitate standardization and to improve the interobserver agreement.

## Key Points

- Variable approaches for membranous urethral length measurement are being used, without detailed description and with substantial differences in length of the membranous urethra, hampering standardization.
- Limited interobserver agreement for membranous urethral length measurement was observed in several studies, while preoperative incontinence risk assessment necessitates high interobserver agreement.
- Literature-based recommendations are proposed to standardize MRI-based membranous urethral length measurement for increasing interobserver agreement and improving preoperative incontinence risk assessment, using anatomical landmarks on sagittal T2-weighted images.

✉ Ivo G. Schoots  
i.schoots@erasmusmc.nl

Thierry N. Boellaard  
tn.boellaard@nki.nl

<sup>1</sup> Department of Radiology, Netherlands Cancer Institute, Plesmanlaan 121, 1066 CX Amsterdam, the Netherlands

<sup>2</sup> Department of Urology, Netherlands Cancer Institute, Amsterdam, the Netherlands

<sup>3</sup> Department of Urology, Amsterdam University Medical Centers, Amsterdam, the Netherlands

<sup>4</sup> Department of Radiology and Nuclear Medicine, Erasmus University Medical Center, Rotterdam, the Netherlands

**Keywords** Urethra · Prostatectomy · Urinary incontinence · Prostatic neoplasms · Magnetic resonance imaging

### Abbreviations

BMI	Body mass index
Cor	Coronal
CPRED	Continence prediction tool
FSE	Fast spin echo
ICC	Intraclass correlation coefficient
LUTS	Lower urinary tract symptoms
MRI	Magnetic resonance imaging
MU	Membranous urethra
MUL	Membranous urethral length
NA	Not available
PI-RADS	Prostate Imaging – Reporting and Data System
PRISMA	Preferred Reporting Items for Systematic Reviews and Meta-analyses
RP	Radical prostatectomy
Sag	Sagittal
TSE	Turbo spin echo

### Introduction

In men with localized prostate cancer, several (curative) treatment options are available, such as radical prostatectomy (RP), external beam radiotherapy, brachytherapy, and active surveillance, all with good oncological outcome [1]. The oncological benefit of each treatment should be carefully weighed against the risk in terms of side effects by both the physician and patient (shared decision-making). The major potential side effects of RP are urinary incontinence and erectile dysfunction, both impacting on quality of life [2]. Counselling patients about these potential side effects is part of the shared decision-making on treatment [1]. Algorithms on individual risk assessment on postoperative urinary incontinence are available, guiding this counselling process [3].

Besides patient-related factors (e.g. age, pre-existing lower urinary tract symptoms (LUTS), and body mass index (BMI)) and surgical factors (e.g. nerve sparing), it was reported that magnetic resonance imaging (MRI)-based anatomical related factors may improve the individual risk assessment of incontinence after RP [4, 5]. The most studied MRI parameter has been the membranous urethral length (MUL). Recent meta-analyses have shown the predictive value of the MRI-based MUL measurement [6, 7] with larger MUL is associated with significantly greater odds for return to continence [7].

The potential impact of pre-treatment incontinence risk assessment for treatment decision-making including the MUL as input parameter is embraced in urological surgical

practices [3]. Several institutions have adopted their own prediction models and have calculated their own threshold for low- and high-risk postoperative (in)continence, including the MUL [3, 8, 9]. However, before implementation into broad clinical practices, there should be agreement on the standardized approach of MUL measurement.

The purpose of this review was to investigate the current literature on the utility of MUL measurement, to identify objective findings regarding MRI acquisition, anatomical landmarks, and measurement definitions, and to propose the first literature-based recommendations on how MUL measurement on pre-treatment MRI scans should be performed.

### Methods

#### Objective

We investigated the literature on published MUL measurements, including measuring approaches, MRI sequences used, population mean/median values, type of observer, and observer agreement. We proposed recommendations to standardize MRI-based MUL measurement using anatomical landmarks, with detailed descriptions and illustrations of MUL measurements and measurement pitfalls.

#### Search strategy

A systematic search was conducted using the Embase, Medline ALL Ovid, Web of Science Core Collection, Cochrane CENTRAL register of trials, and Google Scholar databases up to August 5, 2022, without restrictions regarding publication date or language (supplementary material, appendix 1). The literature search was conducted by a medical librarian. References from selected studies were also screened. This search was also used in a previous publication, but has been updated [6].

#### Inclusion criteria

The study population was limited to men with non-metastasis primary diagnose prostate cancer who underwent RP using any route or approach. Randomized controlled trials and prospective and retrospective cohort studies reporting data on preoperative MRI-based MUL measurements and follow-up data on urinary continence were included. There were no restrictions on follow-up time. We excluded unpublished data, conference abstracts, and review articles. We also excluded studies with the smallest number of patients for published papers using the same data sets (in case of complete overlapping data).

## Data extraction

We followed the Preferred Reporting Items for Systematic Reviews and Meta-analyses (PRISMA) process for reporting study inclusion and exclusion [10]. The abstract and full-text screening and subsequent data extraction were carried out by two researchers independently (M.C.d.H. and T.N.B.). Discrepancies between the reviewers were resolved via discussion (M.C.d.H., T.N.B., and I.G.S.). A data extraction form was developed to collect information on the patient characteristics and study methodology (surgical technique, MRI protocol, questionnaires, and continence follow-up protocols). More detailed data extraction on MUL measurement methodology used (MRI sequence, image orientation, landmarks, agreement) was performed by one researcher (T.N.B.).

## Statistical analysis

This literature review refers to descriptive data; therefore, statistical analysis was not performed.

## Results

### Study parameters

We included 50 papers (Table 1), widely distributed over the world, dominated by South Korea ( $n=16$ ), the USA ( $n=10$ ), and Japan ( $n=9$ ). The studies cover 18,545 men with pre-treatment MRI.

### MRI sequences, anatomical landmarks, and lengths

T2-weighted images for MUL measurement were used in all studies that specified the MRI sequence; either using sagittal images ( $n=18$ ), coronal images ( $n=13$ ), or both ( $n=12$ ) (Table 1).

The anatomical landmark of the proximal end of the membranous urethra (MU) was most commonly described as ‘prostatic apex’.

The anatomical landmark of the distal end of the MU was most commonly described as ‘level of the urethra at the penile bulb’ and ‘entry of the urethra into the penile bulb’. Detailed reproducible measurement technique descriptions were lacking in all studies.

The mean MUL was reported between 10.4 and 17.1 mm and median MUL between 7.3 and 17.3 mm, showing large variations.

Measurements were performed by urologists, radiologists, and trainees.

Articles did not specify the location of the measurement line on sagittal images (e.g. anterior, central, posterior to the

urethra) and exact line orientation. In the provided figures in the articles, the location of the measurement line is variable. Additionally, there is no evidence on how to deal with an anterior membranous urethra (MU) overlapping apex.

### Interobserver agreement

Six studies reported on the interobserver agreement. The intraclass correlation coefficient (ICC) was reported by 5 studies, ranging from 0.34 to 0.89 (Table 2).

### Recommendations based on literature for reproducible MUL measurement

Based on current observations, we suggest to measure the MUL in a way with high interobserver observer agreement [11]. We propose the following recommendations:

- Acquire high-resolution T2-weighted images, according to PI-RADS guidelines [12], preferably on 3-Tesla scanners, in both sagittal and coronal planes.
- Measure the MUL in sagittal T2-weighted images since the coronal images are usually not angulated parallel to the MU.
- Standardize the measurement approach into the following
  - Identify the hyperintense urethral lumen of the MU on one of the midsagittal images, and the dorsal hypointense membranous structure.
  - Place the measurement line just dorsally from and perpendicular to this hyperintense urethral lumen, from the prostate apex to the penile bulb.
  - Identify the upper (cranial) limit, where the measurement line intersects with the prostate apex defined as the lowest border of the peripheral zone at the dorsal prostate. Scroll parasagittally to the left and right to confirm the lowest border of the peripheral zone. When in doubt, crosslink with coronal images.
  - Identify the lower (caudal) limit, where the MU enters the penile bulb. The landmark for the penile bulb is the intersection of the urethra with the bulb of the corpus spongiosum. Scroll parasagittally to left and right to confirm the border of the penile bulb. When in doubt, crosslink with the coronal images.

### Illustrations of proposed measurement technique

The proposed measurement and difference between coronal angulation and MUL measurement direction are shown

**Table 1** Study parameters, MRI sequences, anatomical landmarks and lengths

Author, publication year	Subjects ( <i>n</i> =)	MUL (mm)	Follow-up moments post-prostatectomy	Prostate size	Definition of MUL measurement (description in methods)	MRI sequence	Orientation	Proximal MU landmark	Distal MU landmark	Reader	Country
Cho, 2015 [17]	27	11.3 ± 1.6*	1, 3, 6, 9, and 12 months	39.4 ± 14.2 g	A straight line between the prostatic apex and the penile bulb was drawn in the mid-sagittal plane and the coronal plane	T2 TSE	Sagittal and coronal (mean)	Prostatic apex	Penile bulb	Radiologist	South Korea
Choi, 2015 [18]	158	11.8/11.9*	1, 3, and 6 months	31/33.2 ml	The length from the prostatic apex to the level of the urethra at the penile bulb in the midline sagittal plane	T2	Sagittal	Prostatic apex	Level of the urethra at the penile bulb	NA	South Korea
Coakley, 2002 [19]	180	14 (6–24)*	12 months	NA	The distance from the prostatic apex to the entry of the urethra into the penile bulb	T2 FSE	Coronal	Prostatic apex	Entry of the urethra into the penile bulb	1 of 2 readers	USA
Fukui, 2019 [20]	270	7.3#	1, 3, 6, and 12 months	NA	NA	T2 <sup>i</sup>	Sagittal	NA	NA	NA	Japan
Greenberg, 2022 [21]	251	14 (12–17)# rad 15 (12–18)# uro	Within 6, 12, and 24 months	34 (27–44) cc	From the lowest of the prostatic apex to the entry of the urethra into the penile bulb	T2 <sup>i</sup>	Sagittal (uro)	Prostatic apex	Entry of the urethra into the penile bulb	22 radiologists 1 urologist	USA
Grivas, 2018 [8]	439	12.9 (1.7)/11.5 (1.5)*	6, 12, 18, and 24 months	49.2 cm <sup>3</sup> (68.1)/44.5 cm <sup>3</sup> (19.3)	NA	T2 TSE <sup>a</sup>	Coronal and sagittal	NA	NA	1 urologist 1 radiologist	The Netherlands
Grivas, 2017 [22]	49	16.2 (14.1–18.4)#	6 and 12 months	62 cm <sup>3</sup> (55–78)	NA	T2 TSE <sup>a</sup>	Sagittal <sup>a</sup>	NA	NA	1 urologist 1 radiologist	The Netherlands
Hakimi, 2011 [23]	75	14.6 (8–26)*	1.5, 3, 6, 9, 12, 15, and 18 months	58.4 (28–185) g	The distance from the apex of the prostate to the bulb	T2	Coronal <sup>i</sup>	Apex of the prostate	Bulb	1 of 2 radiologists	USA

**Table 1** (continued)

Author, publication year	Subjects (n=)	MUL (mm)	Follow-up moments post-prostatectomy	Prostate size	Definition of MUL measurement (description in methods)	MRI sequence	Orientation	Proximal MU landmark	Distal MU landmark	Reader	Country
Hikita, 2020 [24]	119	12.1 (8.9–16.1)#	1, 3, 6, 9, and 12 months	26.0 (9.6–66.1) ml	The distance from the prostatic apex to the level of the urethra at the penile bulb	T2	Coronal	Prostatic apex	Level of the urethra at the penile bulb	NA	Japan
Hoeh 2022 [25]	68	14.7 (13.0–16.7)# cor 15.1 (12.8, 16.8)# sag	> 6 months	35 (28–45) ml	NA	T2'	Sagittal and coronal	NA	NA	Specialist in urologic imaging, supervised by a board-certified radiologist	Germany
Hong, 2009 [26]	141	NA	6 months	38.4 (17–88) g	NA	NA	NA	NA	NA	2 radiologists	South Korea
Iacovelli 2022 [27]	100	NA <sup>s</sup>	1, 3, 6, and 12 months	NA <sup>s</sup>	NA	NA	Sagittal	NA	NA	Two radiologists	Italy
Ikarashi, 2018 [28]	204 (10 were incontinent preoperative)	13.1 (4.5–22.9)#	3 (at least), 6, 9, and (max) 12 months	38 (7–94) g	A distance from the prostatic apex to the level of the urethra at the penile bulb	T2	Coronal and sagittal	Prostatic apex	Level of the urethra at the penile bulb	Several urologists and a researcher	Japan
Jeong, 2012 [29]	708	NA <sup>s</sup>	3, 6, 9, 12, 15, 18, 21, and 24 months	NA	NA	NA	NA	NA	NA	NA	South Korea
Jeong, 2013 [30]	731	12.8 (6–23)*	2 weeks, 1, 3, 6, 9, 12, and 24 months	38.5 (16.0–141.0) ml	The distance from the prostatic apex to the entry of the urethra into the penile bulb	T2	Coronal	Prostatic apex	Entry of the urethra into the penile bulb	2 uro-radiologists	South Korea
Jeong, 2014 [31]	1168	NA <sup>s</sup>	1, 3, and 12 months	NA	NA	NA	NA	NA	NA	NA	South Korea
Kadono, 2016 [32]	111	13.6 ± 2.4*	12 months	40.4 ± 9.0 ml	The distance from prostatic apex to the entry of the urethra into the penile bulb	NA	Coronal	Prostatic apex	Entry of the urethra into the penile bulb	NA	Japan

Table 1 (continued)

Author, publication year	Subjects (n=)	MUL (mm)	Follow-up moments continece post-prostatectomy	Prostate size	Definition of MUL measurement (description in methods)	MRI sequence	Orientation	Proximal MU landmark	Distal MU landmark	Reader	Country
Kim, 2011 [33]	763 (3.9% preoperative incontinent)	NA <sup>s</sup>	1, 3, 6, 9, 12, 18, and 24 months	NA	From prostatic apex to the level of the urethra at penile bulb	NA	Sagittal	Prostatic apex	Level of the urethra at penile bulb	NA	South Korea
Kim, 2020 [34]	190	14.6 ± 3.0 cor/14.2 ± 2.7 sag*	8 weeks, 3, 6, 9, and 12 months	36.4 (29.5–45) cm <sup>3</sup>	From the inferior end of the prostatic apex to the level of the penile bulb	T2 <sup>i</sup>	Coronal and sagittal	Inferior end of the prostatic apex	Level of the penile bulb	1 uro-radiologist 1 uro-oncology fellow 1 urologist	UK
Kim, 2019 [35]	529	12.3 ± 4.5* 11.7#	1, 3, 6, and 12 months	35.6 ± 14.6, 32.0	The distance from the posterior prostatic apex to the urethra level at the penile bulb	T2 <sup>i</sup>	Sagittal	Posterior prostatic apex	Urethra level at the penile bulb	NA	South Korea
Kitamura, 2019 [36]	320	10.5 (9.3–11.5)#	1, 3, 6, and 12 months	25.0 (19.0–35.0)	The distance from the prostatic apex to the level of the urethra at the penile bulb	T2	Sagittal	Prostatic apex	Level of the urethra at the penile bulb	1 urologist	Japan
Ko, 2020 [37]	123	NA	1–3 months	NA	The distance from the prostatic apex to the entry of the urethra into the penile bulb	NA	Coronal and sagittal	Prostatic apex	Entry of the urethra into the penile bulb	1 radiologist	USA
Kohjimoto, 2020 [38]	179	17.3 (14.6–19.7)#	3, 6, 12, and 24 months	NA	The distance from prostatic apex to the entry of urethra into penile bulb	T2	Coronal	Prostatic apex	Entry of urethra into penile bulb	NA	Japan

Table 1 (continued)

Author, publication year	Subjects (n=)	MUL (mm)	Follow-up moments post-prostatectomy	Prostate size	Definition of MUL measurement (description in methods)	MRI sequence	Orientation	Proximal MU landmark	Distal MU landmark	Reader	Country
Lamberg, 2022 [39]	589	NA <sup>s</sup>	3, 6, 12, and 24 months	NA <sup>s</sup>	The distance from the prostate apex to the urethral entry into the penile bulb	T2	Sagittal and coronal	Prostate apex	Penile bulb	Three abdominal radiologists One abdominal radiology fellow	USA
Lee, 2013 [40]	249: 92 early recovery (<3 months), 157 late recovery (remaining)	NA <sup>s</sup>	Monthly, at least 3 months	NA	The distance from the prostatic apex to the entry of the urethra into the penile bulb	T2	Coronal	Prostatic apex	Entry of the urethra into the penile bulb	NA	South Korea
Lee, 2006 [15]	156	NA	Within 3 months	NA	NA (only reference)	T2 FSE	NA	NA	NA	2 radiologists	South Korea
Lee, 2014 [16]	1011	NA	1, 3, 6, and 12 months	NA	NA (only reference)	NA	Coronal	NA	NA	2 radiologists	South Korea
Lee, 2020 [41]	2310 (of which 610 aged > 70 years)	NA <sup>s</sup>	3 and 12 months	NA	NA	NA	NA	NA	NA	NA	South Korea
Li, 2020 [42]	156	NA	6, 9, and 12 months	31.0 (23.6–40.3)/ 33.8 (22.1–44.8)	The inferior edge of the prostate apex to the superior margin of the penile bulb	T2 FSE	Sagittal <sup>1</sup>	Prostate apex	Superior margin of the penile bulb	NA	China
Lim, 2012 [43]	94	10.4 ± 3.8*	12 months	29.7 ± 13.5 ml	The most prominent portion of the prostate apex to the level of the urethra at the penile bulb	T2	Sagittal (cross-referenced to coronal)	Most prominent portion of the prostate apex	Level of the urethra at the penile bulb	1 radiologist	South Korea
Lin, 2020 [44]	602	14.6 ± 3.6*	3, 6, and 12 months	42.6 ± 21.2	A straight line between the prostatic apex and the penile bulb	NA	Coronal and sagittal	Prostatic apex	Penile bulb	2 readers	Australia

Table 1 (continued)

Author, publication year	Subjects (n=)	MUL (mm)	Follow-up moments post-prostatectomy	Prostate size	Definition of MUL measurement (description in methods)	MRI sequence	Orientation	Proximal MU landmark	Distal MU landmark	Reader	Country
Matsushita, 2015 [45]	2849: 1899 training, 950 validation	12#	At least 6 months, 1 year	31 cm <sup>3</sup>	NA (only reference)	T2 FSE	NA	NA	NA	1 reader (accuracy confirmed by radiologist)	USA?
Mendoza, 2011 [46]	80	17.1 ± 4.5*	Monthly until 6 months	34.7 ± 17.8 g	From the prostate base to the bulb	NA	Coronal	NA	NA	Radiologists	USA
Nguyen, 2008 [47]	274	14.0*	12 months	37.5 (12.6–175.8) cc	The distance from the prostatic apex to the entry of the urethra into the penile bulb. Urethral sphincter anatomy was studied using 3-cross referenced planes of T2-weighted images, that is on the midline sagittal plane with coronal reference and on the coronal plane with axial reference	T2 FSE	Coronal and sagittal (unclear how combined)	Prostatic apex	Entry of the urethra into the penile bulb	1 reader	USA
Omishi, 2018 [48]	215	12.7 ± 3*	Monthly until patients were continent	40.5 ± 13.8 g	A distance from apex of prostate to the urethra at the level of the penile bulb	T2	Coronal and sagittal (unclear how combined)	Apex of prostate	Urethra at the level of the penile bulb	2 urologists	Japan
Ota, 2021 [49]	50	12.7 (11.2–13.9)/11.3 (10.2–12.8)#	1, 3, 6, and 12 months	27.2 (22.9–37.2) ml/ 22.7 (18.2–29.5) ml	NA	NA	NA	NA	NA	NA	Japan



**Table 1** (continued)

Author, publication year	Subjects (n=)	MUL (mm)	Follow-up moments post-prostatectomy	Prostate size	Definition of MUL measurement (description in methods)	MRI sequence	Orientation	Proximal MU landmark	Distal MU landmark	Reader	Country
Oza, 2022 [50]	NA <sup>s</sup>	NA <sup>s</sup>	12 months	33.50 ml	Distance between the apex of the prostate to the bulb of the penis	T2	Sagittal and coronal	Apex of the prostate	Bulb of the penis	Urologist and uro-radiologist	UK
Paparel, 2009 [51]	64 with pre and post OK MRI	14#	Median 7 months	NA	The distance from the prostatic apex to the level of the urethra at the penile bulb	T2 FSE	Sagittal (cross-referenced to coronal)	Prostatic apex	Level of urethra at the penile bulb	1 radiologist 1 urologist (consensus)	USA
Park, 2021 [9]	166	14.7 (5.1–24.8)*	3 months	44 (19–150) mm <sup>3</sup>	NA	T2	Sagittal <sup>i</sup>	NA	NA	1 radiologist	South Korea
Regis, 2019 [52]	72	NA	1, 6, and 12 months	41 (15–155) cm <sup>3</sup>	NA	T2	Coronal	NA	NA	1 urologist (in consensus with radiologist if doubtful)	Spain
Sadahira, 2019 [53]	70	? (8.7–14.9)#	12 months	? (8.9–103.7) ml	The entry of the urethra into the penile bulb to the prostatic apex	T2 TSE	Sagittal	Prostatic apex	Entry of the urethra into the penile bulb	1 urologist	Japan
Sauer, 2019 [54]	316	10.5 (5.0–25.0)*	6 and 12 months (and 1 week after catheter removal)	47 (20–160) ml	The distance of two horizontals: (1) on the level of the deepest part of the prostate's apex and (2) on the highest visible part of the penile bulbous	T2	Sagittal	Deepest part of prostate apex	Highest visible part of the penile bulbous (between 2 horizontal)	2 radiologists	Germany
Schmid, 2020 [55]	42	NA	6 weeks, 3, 6, and 12 months	NA	The distal prostate apex to the proximal penile bulb in sagittal orientation	T2	Sagittal	Distal prostate apex	Proximal penile bulb	NA	Switzerland

Table 1 (continued)

Author, publication year	Subjects (n=)	MUL (mm)	Follow-up comments post-prostatectomy	Prostate size	Definition of MUL measurement (description in methods)	MRI sequence	Orientation	Proximal MU landmark	Distal MU landmark	Reader	Country
Son, 2013 [56]	258 (PALP vs RRP)	13.1 ± 2.4*	2 weeks, 1, 3, 6, 9, and 12 months	43.3 ± 24.9 ml	The distance from the prostatic apex to the entry of the urethra into the penile bulb	T2	Coronal	Prostatic apex	Entry of the urethra into the penile bulb	NA	South Korea
Song, 2017 [57]	186	15.6 ± 2.7* 15.9 (7.2–22.9)#	1, 3, 6, and 12 months	34.0 ± 15.9, 30.0 (8.0–113.0) ml	A distance from the apex of prostate to the urethra at the level of the penile bulb	T2	Sagittal (cross-referenced to the coronal)	Apex of prostate	Urethra at the level of the penile bulb	2 non-radiologists	South Korea
Tienza, 2018 [58]	746	NA	12 months	NA	NA	T2	Sagittal	NA	NA	NA	Spain
Tutolo 2022 [59]	209	14 (11–16)#	3 months, 3–6 months, 6–12, and from 12 to last-month follow-up	37 (27–51) cm <sup>3</sup>	Distance between the prostate apex and the penile bulb	T2	Sagittal	Prostate apex	Penile bulb	Two urologists (after training by a radiologist)	Belgium Italy
Von Bodman, 2012 [60]	600	13 (11–16)#	6 and 12 months	15.8 (12.4–20.0) cm <sup>3</sup>	From the apex of the prostate to the base of the urethral bulb	T2	Coronal	Apex of the prostate	Base of the urethral bulb	2 raters trained by a radiologist	USA?
Wenzel, 2021 [61]	128	15 (12–17)#	24 h after catheter removal (routinely between 5 and 7 days after surgery)	40 (30–50)	NA	T2 <sup>1</sup>	Sagittal and coronal	NA	NA	Researcher with special training supervised by a radiologist	Germany
Yang, 2020 [62]	150	13.8 ± 3.7*	3, 6, and 12 months	40.1 ± 26.1 ml	A distance from the apex of prostate to the urethra at the level of the penile bulbi	T2	Coronal	Apex of prostate	Urethra at the level of the penile bulbi	NA	China

FSE fast spin echo, MU membranous urethra, NA not available, TSE turbo spin echo. \*Mean, #median, a average, b based on figure in manuscript, c requested from authors, d provided for subgroups only, rad radiologist, uro urologist

**Table 2** Interobserver agreement

Author, publication year	Statistics	Interobserver	Interobserver with training	Intraobserver
Greenberg, 2022 [21]	ICC	0.34 (sag)		
Kim, 2020 [34]	ICC	0.89 (cor) 0.77 (sag)		
Lamberg, 2022 [39]	ICC	0.38 (sag and cor)	0.62 (sag and cor)	
Sauer, 2019 [54]	ICC	0.82 (sag)		
Von Bodman, 2012 [60]	Weighted kappa	0.48 (cor)		
Veerman, 2022 [11] (additional data)	ICC	0.84 (sag)		0.93–0.98

*Cor* coronal, *Sag* sagittal, *ICC* intraclass correlation coefficient

in Fig. 1. The critical steps of our proposed MUL measurement technique are shown in Fig. 2. The identification of the upper limit (lower border of the peripheral zone) is illustrated by Fig. 3. The identification of the lower limit (upper border of the penile bulb) is illustrated by Fig. 4.

### Anatomy and measurement pitfalls

The sphincter is composed of an external rhabdosphincter (skeletal muscle) that is responsible for the active continence and the internal lissosphincter (smooth muscle) that is responsible for the passive continence (Fig. 5f). The rhabdosphincter is the thickest at the level of the MU and has fibres continuous with the anterior fibromuscular stroma. The lissosphincter starts in the bladder neck and continues to the upper border of the penile base (perineal membrane). The MU is the part of the urethra between the prostatic apex and penile bulb (Fig. 5g). Both external and internal sphincter fibres are located at the level of the MU.

There are several pitfalls to consider when measuring the MU, resulting from the complex anatomy shown in Fig. 5a. In Fig. 5b–e, the correlation between the anatomy and the most important MRI pitfalls are shown. In Fig. 5f, g, the concept of the sphincter complex is shown according to Koraitim [13], showing a MUL measurement line in the anatomy image illustrating what is measured on MRI using our proposed technique (Fig. 5g).

Pitfalls include challenging superior limit (abnormal peripheral zone intensity, signal intensity of the retroprostatic part of the rectovesical space similar to the peripheral zone), challenging lower limit (double contour or difficulty to appreciate correct penile bulb contour at midsagittal slice (Fig. 6)), suggestion of rhabdosphincter fibers of MU continuing in the prostate (supplemental Fig. 1), angulated MU (supplemental Fig. 4), and crosslink errors between coronal and sagittal images (supplemental Fig. 5). It is important to have a good understanding of the anatomy of the MU and its surrounding structures. Additional text and illustrations on the anatomy and pitfalls are provided in the supplementary material, appendix 2.

### Discussion


The aim of the review was to investigate the MUL measurement and its interobserver agreement and propose literature-based recommendations to standardize MUL measurement for increasing interobserver agreement. To our knowledge, this is the first review to summarize the literature on MUL measurement methods and also the first to propose a standardized MRI-based MUL measurement approach with detailed landmarks and pitfalls. This could provide guidance for radiologists and urologists that would like to start performing these measurements as part of the preoperative risk assessment of postoperative urinary incontinence in men with localized prostate cancer. Standardization could also help to use externally validated urinary continence prediction tools.

### Populations

We observed that most literature on the MUL is from Asian countries. One study showed the average Asian MUL was significantly smaller than a non-Asian MUL [14]. The exact effect of different MUL size across populations (and whether this variation is associated with body length) should be further studied, as this may influence the continence prediction models suitable for different populations.

### Sequence and orientation

We observed that all studies that specified the image used for MUL measurement made use of T2-weighted images. Although most studies included sagittal images (sagittal only or both coronal and sagittal), a substantial number of publications used solely coronal images. The advantage of coronal images is that it allows easier delineation of upper and lower border. In literature, coronal and sagittal MUL measurements have shown significant correlation with urinary incontinence after RP and some studies showed good correlation between the sagittal and coronal measurements. We, however, recommend the use of sagittal images for




**Fig. 1** Proposal of membranous urethral length (MUL) measurement on midsagittal MR images. **a** Midsagittal T2w image of the prostate. Sagittal MR images are mandatory for appropriate MUL measurements. The proposed MUL measurement (red line) is determined at the dorsal side of the urethra lumen (this MUL was measured 16 mm). The upper border of the MU is determined by the presence of prostatic tissue. Intraprostatic urethra is excluded from the measurement as intraoperatively sparing is not always performed or possible (i.e. apical tumors). The upper border of the MU is determined by the intersection of the urethra with the dorsally located peripheral zone (white line). The lower border of the MU is determined by the intersection of the urethra with the entrance

to penile bulb (green line). **b** Crosslinking of coronal T2w images with sagittal images may help identifying and determining the borders of the anatomical structures related to the MU. Delineation of the peripheral zone (white line) on coronal images informs on the intersection with the MU. **c, d** Crosslinking of sagittal (**c**) and coronal (**d**) T2w images with a cross mark (red X) in PACS viewing software may confirm the correct identification of the lower border of the peripheral zone. The scan direction of the coronal images is illustrated by the blue dashed line. Notice that this coronal scan orientation may not be similar to the MU direction (dashed yellow line), which may lead to an inappropriate MUL measurement when coronal images would have been used

MUL measurements. The angulation of coronal images is often different from the correct MU orientation that is seen in sagittal images. These variations in angulation will lead to different measurements compared with sagittal, causing under- or overestimation. Also, different coronal angulations will lead to different measurements in the same patient.

Another theoretical possibility could be to angulate the coronal images parallel to the MU. However, it is questionable whether one should adjust the angulation and consequently the prostate appearance you are used to, especially for one measurement. Also, this requires training of radiologic technicians to accurately angulate parallel to the MU.



**Fig. 2** Critical steps in membranous urethral length (MUL) measurement on midsagittal MR images. **a, b** A prostate cancer patient with sagittal T2w images on preoperative MRI. In MUL measurement, critical steps need to be distinguished: (1) the hyperintense lumen (yellow arrows) and the hypointense dorsal part (orange arrows) of the membranous urethra need to be identified. (2) The line of the membranous urethra measurement should be placed dorsally and parallel (red line, MUL). (3) The upper border of the measured membranous urethra intersects with the prostate apex, defined as the lower

border of the peripheral zone at the dorsal side (white line). (4) The lower border the measured membranous urethra intersects with the penile bulb, the bulb of the corpus spongiosum (green line). **c, d** Another prostate cancer patient with sagittal T2w images on preoperative MRI. Notice the difference between hyperintense signal of the peripheral zone and retroprostatic part of the rectovesical space (yellow delineation). Also notice base of the penile bulb can be slightly curved (purple arrow), extending the MUL slightly. The measurement (red line) was 20 mm in this case


### Anatomical landmarks and line placement

We have seen that similar landmarks were used for upper and lower border of the measurement (‘prostatic apex’ and level/entry of the urethra at the penile bulb). However, the exact measurement descriptions in literature lacked details and are therefore poorly reproducible. For example, the exact measurement line location and orientation were not described and it was not mentioned how it was dealt with different apex types. All these factors can influence the MUL length. The transitional zone may be overlapping anteriorly [15] and it is unclear if measured towards prostate apex dorsally or anteriorly. The apical shape of the prostate is variable and may influence the predicted

incontinence [16]. For reproducibility purposes, we suggest a standard measurement at the dorsal side of the MU towards the peripheral zone. To our knowledge, it is unknown whether measurement towards an apical protruding transitional zone is better for the predictive power of MUL measurements and intra- and interobserver agreement. In our experience, the dorsal side is easier to measure than central or anterior and parallel to the urethra would seem a rational approach.

### Measurement variations

The large variation in mean and median population MUL (median 7.3 to 17.3 mm) is suggesting large variation



**Fig. 3** Challenges in membranous urethral length (MUL) measurement — the upper border, intersecting with the peripheral zone (1). Identifying the lower border of the peripheral zone may be challenging. Scrolling through the sagittal T2w images may help in determining the upper border of the MU, by better identifying the lower border of the peripheral zone. At the upper border of the membranous urethra, the retroprostatic part of the rectovesical space surrounding the peripheral zone in the prostate apex could be difficult to distinguish from the peripheral zone. These structures have similar signal intensities to the peripheral zone on T2-weighted images, especially when peripheral zone is less hyper-

intense on T2w images due to inflammation, fibrosis, blood products, or cancer. Scrolling 1 or 2 slices right (**a**; red arrow) and left (**c**; blue arrow) from the midsagittal view (**b**; white arrow) may improve the determination of the peripheral zone, and subsequently the upper border of the MU. On midsagittal images of the prostate, the dark tissue in the membranous urethra lumen may extend intraprostatic. This intraprostatic tissue will most likely be resected and should therefore be excluded from measurement. **d** Coronal T2 image shows the sagittal orientation of the lower borders of the peripheral zone on right parasagittal image (red arrow), midsagittal image (white arrow), and left parasagittal image (blue arrow)

in measurement method or population. The large difference between these specific studies may be measurement method related, since both studies are from Japan.

### Observer agreement


Few studies reported on interobserver agreement variable results from fair to high agreement. In a recent agreement study from our group, we have seen high inter- and intra-observer agreement results using our defined landmarks [11]. It is important to obtain the highest possible intra- and

interobserver agreement as a variation of several millimetres in MUL measurement results in substantially different percentage-predicted continence after RP.

### Imaging technique

We believe MRI is the technique of choice to use for the MUL measurements. It is possible to measure the MUL with other techniques, such as ultrasound and retrograde urethrography. However, the MRI is already made for targeting biopsy and/or staging and is able to visualize the anatomy very well.

**Fig. 4** Challenges in membranous urethral length (MUL) measurement—the lower border, intersecting with the penile bulb (1). Scrolling through the sagittal T2w images may help in determining the lower border of the MU, by better identifying the upper border of the penile bulb. The lower border of the membranous urethra is determined by the intersection with the upper border of the penile bulb. Scrolling 1 or 2 slices right (a; red arrow) and left (c; blue arrow) from the midsagittal view (b; white arrow) may improve the determination of the penile bulb, and subsequently the lower border of the MU. On midsagittal images of the prostate, the hypointense tissue surrounding the membranous urethra lumen may continue into the penile bulb (purple in b). This intrabulbic part should be excluded from measuring. d Coronal T2 image shows the sagittal orientation of the upper borders of the penile bulb on right parasagittal image (red arrow), midsagittal image (white arrow), and left parasagittal image (blue arrow), and the intrabulbic continuation of hypointense tissue (purple line)



## Our recommendations

For some of our measurement recommendations, there will be little discussion. The use of T2-weighted images is standard practice and the landmarks used are very similar in literature. Other recommendations may be a cause for more discussion. For example, the measurement on the midsagittal T2-weighted images, measuring dorsally along the urethra and towards the peripheral zone. Given the lack of evidence, we made these recommendations based on rationale and experience; this is a limitation.

## Other limitations


We did not study how interobserver agreement is of MUL measurements performed by readers (outside our institution) using our proposed measurement technique. Furthermore, the scanner type,

coil type, and scan protocol may affect the image quality and appearance and therefore may influence the MUL measurement.


## Integration of MUL measurements in incontinence risk assessment following surgery

Because of the predictive power of the MUL, the authors believe that institutions are justified to implement the MUL measurements in clinical practise. The radiologists can provide the measurement in their standardized report, providing that the urologist knows how to interpretate the results and the radiologist is skilled in performing the measurement. Although the predictive power of the MUL has been proven in meta-analyses, the best way for the urologist to implement the measurement in decision-making can be debated. It is possible to use risk nomograms for personalized urinary incontinence risk to use for shared decision-making [3, 8].

**Fig. 5** Complex anatomy of the region of the membranous urethra, correlation between MRI pitfalls, and the anatomy and the concept of the male urethral sphincter complex. The complex anatomy of the membranous urethra region in the midsagittal plane with the anatomical names (a). The areas on MRI that are responsible for the most important measurement pitfalls (c–e) with the corresponding area shown in the anatomical illustration (b). The pitfalls shown are fibres of the rhabdosphincter that appear to continue into the prostate (blue, c), dark tissue surrounding the urethra in the penile base (green, d), and the signal intensity of the retroprostatic part of the rectovesical space that can be similar to the peripheral zone (yellow, e). Concept of male urethral sphincter according to Koraitim with names (f, g). The MUL measurement with our proposed technique is indicated by the line (ruler) in g







**Fig. 6** Pitfall in membranous urethral length (MUL) measurement—the lower border, intersecting with the penile bulb (2). The upper contour of the penile bulb may sometimes be difficult to determine at the midsagittal image. Peribulbic tissue surrounding the membranous urethra has various low T2 signal intensities. This tissue contains the perineal body, Cowper glands, and deep transverse perineal muscle. These structures are difficult to appreciate separately from the rhabdosphincter and similar dark signal intensity may continue several millimetres into the penile bulb. This may decrease the accurate demarcation of the lower border of the MU, resulting in extended MUL measurement with poor reproducibility. Scrolling 1 or 2 slices parasagittal (and when difficult) crosslinking with the coronal will help to identify the lower limit of the membranous urethra correctly. **a** A prostate cancer patient with sagittal T2w images on preoperative MRI. The proposed MUL measurement (red line) was challenging at

the lower border, at the intersection with the penile bulb (green line). A ‘double contour’ appearance was suggested on the midsagittal image, as a result of partial volume effects (purple line). **b** Scrolling through the sagittal images left and right (not shown) and crosslinking with coronal images determined the appropriate upper contour of the penile bulb (green) and the intrabulbic hypointense tissue (purple line). **c** A prostate cancer patient with sagittal T2w images on preoperative MRI. The proposed MUL measurement (red line) was challenging at the lower border, at the intersection with the penile bulb (green line), due to the intrabulbic continuation hypointense tissue surrounding the urethra at the midsagittal image. **d** Scrolling through the sagittal images left and right (not shown) and crosslinking with coronal images determined the appropriate upper contour of the penile bulb (green) and the intrabulbic continuation of hypointense tissue surrounding the urethra (purple line)

Other methods could be to stratify patients into two or three categories of MUL size (e.g. high, intermediate, and low risk). Using these categories, small interobserver variations would lead to the same category. Although in some cases, a 1-mm difference may lead to a different risk category. At our own institution, we use the continence prediction tool (CPRED) which is based on the preoperative MRI-measured MUL, inner levator muscle distance (ILD), and the

estimated extent of fascia preservation (i.e. nerve sparing) during RP [8]. However, the ILD is not as extensively studied as the MUL and the predictive power seems less compared with the MUL.

The recommended standardized MUL measurement needs to be validated and consensus among experts needs to be encouraged, including expert opinions.

## Conclusions

In order to improve measurement variability, a literature-based method for measuring the MUL was proposed, supported by several illustrative case studies, in an attempt to standardize MRI-based MUL measurements for appropriate urinary incontinence risk assessment following radical surgery.

**Supplementary information** The online version contains supplementary material available at <https://doi.org/10.1007/s00330-023-10180-7>.

**Funding** The authors state that this work has not received any funding.

## Declarations

**Guarantor** The scientific guarantor of this publication is Dr. Ivo Schoots.

**Conflict of interest** The authors of this manuscript declare no relationships with any companies, whose products or services may be related to the subject matter of the article.

**Statistics and biometry** No complex statistical methods were necessary for this paper.

**Informed consent** Written informed consent was not required for this study because it is a review.

**Ethical approval** Institutional review board approval was not required because it is a review.

**Study subjects or cohorts overlap** Not applicable.

## Methodology

- Review

**Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

## References

- Mottet N, van den Bergh RCN, Briers E et al (2021) EAU-EANM-ESTRO-ESUR-SIOG Guidelines on Prostate Cancer-2020 update. Part 1: Screening, diagnosis, and local treatment with curative intent. *Eur Urol* 79:243–262
- Chen RC, Basak R, Meyer AM et al (2017) Association between choice of radical prostatectomy, external beam radiotherapy, brachytherapy, or active surveillance and patient-reported quality of life among men with localized prostate cancer. *JAMA* 317:1141–1150
- Tillier CN, Vromans RD, Boekhout AH et al (2021) Individual risk prediction of urinary incontinence after prostatectomy and impact on treatment choice in patients with localized prostate cancer. *Neurourol Urodyn* 40:1550–1558
- Lardas M, Grivas N, Debray TPA et al (2022) Patient- and tumour-related prognostic factors for urinary incontinence after radical prostatectomy for nonmetastatic prostate cancer: a systematic review and meta-analysis. *Eur Urol Focus* 8:674–689
- Heesakkers J, Farag F, Bauer RM, Sandhu J, De Ridder D, Stenzl A (2017) Pathophysiology and contributing factors in postprostatectomy incontinence: a review. *Eur Urol* 71:936–944
- van Dijk-de Haan MC, Boellaard TN, Tissier R et al (2022) Value of different magnetic resonance imaging-based measurements of anatomical structures on preoperative prostate imaging in predicting urinary continence after radical prostatectomy in men with prostate cancer: a systematic review and meta-analysis. *Eur Urol Focus* 8:1211–1225
- Mungovan SF, Sandhu JS, Akin O, Smart NA, Graham PL, Patel MI (2017) Preoperative membranous urethral length measurement and continence recovery following radical prostatectomy: a systematic review and meta-analysis. *Eur Urol* 71:368–378
- Grivas N, van der Roest R, Schouten D et al (2018) Quantitative assessment of fascia preservation improves the prediction of membranous urethral length and inner levator distance on continence outcome after robot-assisted radical prostatectomy. *Neurourol Urodyn* 37:417–425
- Park S, Byun J (2021) A study of predictive models for early outcomes of post-prostatectomy incontinence: machine learning approach vs. logistic regression analysis approach. *Appl Sci* 11:6225
- Moher D, Liberati A, Tetzlaff J, Altman DG, Group P (2009) Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *PLoS Med* 6:e1000097
- Veerman H, Hagens M, Hoeks C et al (2022) A standardized method to measure the membranous urethral length (MUL) on MRI of the prostate with high inter- and intra-observer agreement. *Eur Radiol* 33:3295–3302
- Turkbey B, Rosenkrantz AB, Haider MA et al (2019) Prostate imaging reporting and data system version 2.1: 2019 update of prostate imaging reporting and data system version 2. *Eur Urol* 76:340–351
- Koraitim MM (2008) The male urethral sphincter complex revisited: an anatomical concept and its physiological correlate. *J Urol* 179:1683–1689
- Basourakos SP, Ramaswamy A, Yu M, Margolis DJ, Hu JC (2021) Racial variation in membranous urethral length and postprostatectomy urinary function. *Eur Urol Open Sci* 27:61–64
- Lee SE, Byun SS, Lee HJ et al (2006) Impact of variations in prostatic apex shape on early recovery of urinary continence after radical retropubic prostatectomy. *Urology* 68:137–141
- Lee H, Kim K, Hwang SI et al (2014) Impact of prostatic apical shape and protrusion on early recovery of continence after robot-assisted radical prostatectomy. *Urology* 84:844–849
- Cho DS, Lee EJ, Kim SJ, Kim SI (2015) The influence of membranous stretched urethral length and urethral circumference on postoperative recovery of continence after radical prostatectomy: a pilot study. *Can Urol Assoc J* 9:E262–E266
- Choi SK, Park S, Ahn H (2015) Randomized clinical trial of a bladder neck plication stitch during robot-assisted radical prostatectomy. *Asian J Androl* 17:304–308
- Coakley FV, Eberhardt S, Kattan MW, Wei DC, Scardino PT, Hricak H (2002) Urinary continence after radical retropubic prostatectomy: relationship with membranous urethral length on preoperative endorectal magnetic resonance imaging. *J Urol* 168:1032–1035
- Fukui S, Kagebayashi Y, Iemura Y, Matsumura Y, Samma S (2019) Preoperative MRI parameters predict urinary continence after robot-assisted laparoscopic prostatectomy in prostatic cancer patients. *Diagnostics (Basel)* 9:102
- Greenberg SA, Cowan JE, Lonergan PE et al (2022) The effect of preoperative membranous urethral length on likelihood of postoperative urinary incontinence after robot-assisted radical prostatectomy. *Prostate Cancer Prostatic Dis* 25:344–350

22. Grivas N, van der Roest R, Tillier C et al (2017) Patterns of benign prostate hyperplasia based on magnetic resonance imaging are correlated with lower urinary tract symptoms and continence in men undergoing a robot-assisted radical prostatectomy for prostate cancer. *Urology* 107:196–201
23. Hakimi AA, Faleck DM, Agalliu I, Rozenblit AM, Chernyak V, Ghavamian R (2011) Preoperative and intraoperative measurements of urethral length as predictors of continence after robot-assisted radical prostatectomy. *J Endourol* 25:1025–1030
24. Hikita K, Honda M, Teraoka S et al (2019) Intravesical prostatic protrusion may affect early postoperative continence in men undergoing robot-assisted radical prostatectomy for prostate cancer. *Neurourol Urodyn* 38:S374–S375
25. Hoeh B, Wenzel M, Muller M et al (2022) Urethral sphincter length but not prostatic apex shape in preoperative MRI is associated with mid-term continence rates after radical prostatectomy. *Diagnostics (Basel)* 12:701
26. Hong SK, Lee ST, Kim SS et al (2009) Effect of bony pelvic dimensions measured by preoperative magnetic resonance imaging on performing robot-assisted laparoscopic prostatectomy. *BJU Int* 104:664–668
27. Iacovelli V, Carilli M, Sandri M et al (2022) The role of preoperative prostatic shape in the recovery of urinary continence after robotic radical prostatectomy: a single cohort analysis. *Prostate Cancer Prostatic Dis* 26(2):374–378. <https://doi.org/10.1038/s41391-022-00563-0>
28. Ikarashi D, Kato Y, Kanehira M et al (2018) Appropriate preoperative membranous urethral length predicts recovery of urinary continence after robot-assisted laparoscopic prostatectomy. *World J Surg Oncol* 16:224
29. Jeong SJ, Kim HJ, Kim JH et al (2012) Urinary continence after radical prostatectomy: predictive factors of recovery after 1 year of surgery. *Int J Urol* 19:1091–1098
30. Jeong CW, Oh JJ, Jeong SJ et al (2013) Effect of dorsal vascular complex size on the recovery of continence after radical prostatectomy. *World J Urol* 1:383–388
31. Jeong SJ, Yeon JS, Lee JK et al (2014) Development and validation of nomograms to predict the recovery of urinary continence after radical prostatectomy: comparisons between immediate, early, and late continence. *World J Urol* 32:437–444
32. Kadono Y, Ueno S, Kadomoto S et al (2016) Use of preoperative factors including urodynamic evaluations and nerve-sparing status for predicting urinary continence recovery after robot-assisted radical prostatectomy: nerve-sparing technique contributes to the reduction of postprostatectomy incontinence. *Neurourol Urodyn* 35:1034–1039
33. Kim SC, Song C, Kim W et al (2011) Factors determining functional outcomes after radical prostatectomy: robot-assisted versus retropubic. *Eur Urol* 60:413–419
34. Kim LHC, Patel A, Kinsella N, Sharabiani MTA, Ap Dafydd D, Cahill D (2019) Association between preoperative magnetic resonance imaging-based urethral parameters and continence recovery following robot-assisted radical prostatectomy. *Eur Urol Focus* 6:1013–1020
35. Kim M, Park M, Pak S et al (2019) Integrity of the urethral sphincter complex, nerve-sparing, and long-term continence status after robotic-assisted radical prostatectomy. *Eur Urol Focus* 5:823–830
36. Kitamura K, China T, Kanayama M et al (2019) Significant association between urethral length measured by magnetic resonance imaging and urinary continence recovery after robot-assisted radical prostatectomy. *Prostate Int* 7:54–59
37. Ko YH, Huynh LM, See K, Lall C, Skarecky D, Ahlering TE (2020) Impact of surgically maximized versus native membranous urethral length on 30-day and long-term pad-free continence after robot-assisted radical prostatectomy. *Prostate Int* 8:55–61
38. Kohjimoto Y, Yamashita S, Kikkawa K, Iba A, Matsumura N, Hara I (2020) The association of length of the resected membranous urethra with urinary incontinence after radical prostatectomy. *Urol J* 17:146–151
39. Lamberg H, Shankar PR, Singh K et al (2022) Preoperative prostate MRI predictors of urinary continence following radical prostatectomy. *Radiology* 303:99–109
40. Lee S, Yoon CJ, Park HJ, Lee JZ, Ha HK (2013) The surgical procedure is the most important factor affecting continence recovery after laparoscopic radical prostatectomy. *World J Men's Health* 31:163–169
41. Lee YJ, Jung JW, Lee S et al (2020) Contemporary trends in radical prostatectomy and predictors of recovery of urinary continence in men aged over 70 years: comparisons between cohorts aged over 70 and less than 70 years. *Asian J Androl* 22:280–286
42. Li Y, Li W, Lu W et al (2020) Association of preoperative urethral parameters on magnetic resonance imaging and immediate recovery of continence following Retzius-sparing robot-assisted radical prostatectomy. *Transl Androl Urol* 9:501–509
43. Lim TJ, Lee JH, Lim JW, Moon SK, Jeon SH, Chang SG (2012) Preoperative factors predictive of continence recovery after radical retropubic prostatectomy. *Korean J Urol* 53:524–530
44. Lin D, O'Callaghan M, David R et al (2020) Does urethral length affect continence outcomes following robot assisted laparoscopic radical prostatectomy (RALP)? *BMC Urol* 20:8
45. Matsushita K, Kent MT, Vickers AJ et al (2015) Preoperative predictive model of recovery of urinary continence after radical prostatectomy. *BJU Int* 116:577–583
46. Mendoza PJ, Stern JM, Li AY et al (2011) Pelvic anatomy on preoperative magnetic resonance imaging can predict early continence after robot-assisted radical prostatectomy. *J Endourol* 25:51–55
47. Nguyen L, Jhaveri J, Tewari A (2008) Surgical technique to overcome anatomical shortcoming: balancing post-prostatectomy continence outcomes of urethral sphincter lengths on preoperative magnetic resonance imaging. *J Urol* 179:1907–11
48. Onishi T, Sekito S, Terabe T, Shibahara T (2018) A combination of findings obtained from pre- and postoperative imaging predict recovery of urinary continence after non-nerve-sparing laparoscopic radical prostatectomy. *Anticancer Res* 38:5525–5530
49. Ota Y, Hamamoto S, Matsuyama N et al (2021) Pelvic anatomical features after Retzius-sparing robot-assisted radical prostatectomy intended for early recovery of urinary symptoms. *J Endourol* 35:296–304
50. Oza P, Walker NF, Rottenberg G et al (2022) Pre-prostatectomy membranous urethral length as a predictive factor of post prostatectomy incontinence requiring surgical intervention with an artificial urinary sphincter or a male sling. *Neurourol Urodyn* 41:973–979
51. Paparel P, Akin O, Sandhu JS et al (2009) Recovery of urinary continence after radical prostatectomy: association with urethral length and urethral fibrosis measured by preoperative and postoperative endorectal magnetic resonance imaging. *Eur Urol* 55:629–639
52. Regis L, Salazar A, Cuadras M et al (2019) Preoperative magnetic resonance imaging in predicting early continence recovery after robotic radical prostatectomy. *Actas Urol Esp (Engl Ed)* 43:137–142
53. Sadahira T, Mitsui Y, Araki M et al (2019) Pelvic magnetic resonance imaging parameters predict urinary incontinence after robot-assisted radical prostatectomy. *Lower Urin Tract Symptoms* 11:122–6
54. Sauer M, Tennstedt P, Berliner C et al (2019) Predictors of short and long term urinary incontinence after radical prostatectomy in prostate MRI: significance and reliability of standardized measurements. *Eur J Radiol* 120:108668
55. Schmid FA, Wettstein MS, Kessler TM et al (2019) Contrast media kinetics in multiparametric magnetic resonance imaging before radical prostatectomy predicts the probability of postoperative incontinence. *World J Urol* 38:1741–1748
56. Son SJ, Lee SC, Jeong CW, Jeong SJ, Byun SS, Lee SE (2013) Comparison of continence recovery between robot-assisted laparoscopic prostatectomy and open radical retro public prostatectomy: a single surgeon experience. *Korean J Urol* 54:598–602. <https://doi.org/10.4111/kju.2013.54.9.598>
57. Song W, Kim CK, Park BK et al (2017) Impact of preoperative and postoperative membranous urethral length measured by 3 Tesla

- magnetic resonance imaging on urinary continence recovery after robotic-assisted radical prostatectomy. *Can Urol Assoc J* 11:E93–E99
58. Tienza A, Robles JE, Hevia M, Algarra R, Diez-Caballero F, Pascual JI (2018) Prevalence analysis of urinary incontinence after radical prostatectomy and influential preoperative factors in a single institution. *Aging Male* 21:24–30
  59. Tutolo M, Rosiello G, Stabile G et al (2022) The key role of levator ani thickness for early urinary continence recovery in patients undergoing robot-assisted radical prostatectomy: a multi-institutional study. *Neurourol Urodyn* 41:1563–1572
  60. Von Bodman C, Matsushita K, Savage C et al (2012) Recovery of urinary function after radical prostatectomy: predictors of urinary function on preoperative prostate magnetic resonance imaging. *J Urol* 187:945–950
  61. Wenzel M, Preisser F, Mueller M et al (2021) Effect of prostatic apex shape (Lee types) and urethral sphincter length in preoperative MRI on very early continence rates after radical prostatectomy. *Int Urol Nephrol* 53:1297–1303
  62. Yang B, Zhang F, Xiao C, Lu J, Ma L, Huang Y (2020) Impact of preoperative magnetic resonance imaging anatomic features on urinary continence recovery after laparoscopic radical prostatectomy. *Urol Int* 104:239–246

**Publisher's note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.