

Erratum to: Regulation of Phytohormone Biosynthesis and Accumulation in *Arabidopsis* Following Treatment with Commercial Extract from the Marine Macroalga *Ascophyllum nodosum*

Owen S. D. Wally · Alan T. Critchley · David Hiltz · James S. Craigie · Xiumei Han · L. Irina Zaharia · Suzanne R. Abrams · Balakrishnan Prithiviraj

Published online: 15 December 2012
© Springer Science+Business Media New York 2012

Erratum to: J Plant Growth Regul
DOI 10.1007/s00344-012-9301-9

There are errors in the published version of Table 1 due to an inadvertent mistake in the calculation. The change made in the table does not alter the results and conclusions drawn in the paper. Please see overleaf for the corrected table.

The online version of the original article can be found under doi:
[10.1007/s00344-012-9301-9](https://doi.org/10.1007/s00344-012-9301-9).

O. S. D. Wally
Department of Environmental Sciences, Nova Scotia Agriculture
College, Truro, NS B2N 5E3, Canada

A. T. Critchley · D. Hiltz
Acadian Seaplants Limited, Dartmouth, NS B3B 1X8, Canada

J. S. Craigie
Institute for Marine Biosciences, National Research Council of
Canada, Halifax, NS B3H 3Z1, Canada

X. Han · L. I. Zaharia · S. R. Abrams
Plant Biotechnology Institute, National Research Council of
Canada, Saskatoon, SK S7N 0W9, Canada

B. Prithiviraj (✉)
Department of Environmental Sciences, Faculty of Agriculture,
Dalhousie University, Truro, NS B2N 5E3, Canada
e-mail: bprithiviraj@dal.ca

Table 1 Concentration of plant growth regulators in various commercial seaweed extracts (ng g⁻¹ dry weight)

Seaweed extract	Auxins										ABA and ABA metabolites							Cytokinins				
	IAA	IAA-ala	IAA-asp	IAA-glu	IAA-leu	IBA	ABA	DPA	PA	Neo-PA	t-ABA	c-ZOG	t-Z	c-Z	2iP	iPA						
	IAA	IAA-ala	IAA-asp	IAA-glu	IAA-leu	IBA	ABA	DPA	PA	Neo-PA	t-ABA	c-ZOG	t-Z	c-Z	2iP	iPA						
Canadian Atlantic ANE 7/21/08	20	12	50	5	2	nd	2.0	nd	nd	nd	nd	3	nd	nd	7	0.3						
Canadian Atlantic ANE 2/6/09	35	10	50	5	2	nd	1.0	nd	nd	nd	nd	2	nd	nd	8	0.5						
Canadian Atlantic ANE 4/10/09	20	10	50	5	3	nd	1.5	nd	nd	nd	nd	2	nd	nd	8	2						
French ANE	3	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	1.0	nd						
American Atlantic ANE	33	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	1.0	3.0						
Irish ANE	nd	5.0	nd	nd	1.0	nd	nd	nd	nd	nd	nd	nd	nd	nd	2.0	25.0						
Norwegian ANE	130	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	2.0	10.0						
S. African <i>Ecklonia</i> extract	nd	4.0	nd	6	nd	20	nd	nd	nd	nd	nd	4	nd	nd	nd	nd						
S. African <i>Ecklonia</i> extract	80	4.0	nd	nd	nd	nd	4	nd	nd	nd	nd	nd	4	nd	4.0	4.0						
N. Pacific <i>Macrocystis</i> extract	nd	4.0	nd	nd	0	40	4	nd	nd	nd	216	nd	nd	1.7	24.0	6.0						
South Pacific <i>Durvillea</i>	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	28.0	1.0						
Chinese <i>Sargassum</i> extract	nd	5	nd	nd	nd	222	2	nd	nd	nd	nd	nd	2.0	nd	2.0	2.0						

Approximately 1 g of seaweed extracts from random samples were dried and pooled for analysis. Canadian Atlantic ANE harvested from three distinct batches were analyzed separated and presented above

IAA indole acetic acid, IAA-ala N-(Indole-3-yl-acetyl)-alanine, IAA-asp N-(Indole-3-yl-acetyl)-aspartic acid, IAA-glu N-(Indole-3-yl-acetyl)-glutamic acid, IAA-leu N-(Indole-3-yl-acetyl)-leucine, IBA indole-3-butyrac acid, ABA cis-Abscisic acid, DPA dihydrophaseic acid, PA phaseic acid, t-ABA trans-ABA, c-ZOG cis-Zeatin-O-glucoside, t-Z trans-Zeatin, 2iP isopentenyladenine, iPA isopentenyladenosine, nd not-detected