

Erratum to: Time-resolved OCT- μ PIV: a new microscopic PIV technique for noninvasive depth-resolved pulsatile flow profile acquisition

Chia-Yuan Chen · Prahlad G. Menon ·
William Kowalski · Kerem Pekkan

Published online: 10 February 2013
© Springer-Verlag Berlin Heidelberg 2013

Erratum to: Exp Fluids (2013) 54:1426
DOI 10.1007/s00348-012-1426-x

The authors would like to correct errors in Table 1 of the original publication of the article.

The corrected Table 1 is given below:

Table 1 Optical data of various μ PIV techniques for high Hct (>10 %) flow measurements

PIV method	Depth resolution	Max. imaging depth
Stereo μ PIV	$10^1 \mu\text{m}^{\text{a}}$	$10^1 \mu\text{m}^{\text{a}}$
Confocal- μ PIV	$10^0 \mu\text{m}^{\text{b}}$	$10^1 \mu\text{m}^{\text{c}}$
OCT- μ PIV	$10^1 \mu\text{m}^{\text{d}}$	$10^3 \mu\text{m}^{\text{d}}$

^a Data were acquired using the same optical set-up and objective as described in Lindken et al. (2006)

^b Patrick et al. (2011), Lima et al. (2006)

^c Patrick et al. (2011)

^d Davis et al. (2009)

The online version of the original article can be found under doi:[10.1007/s00348-012-1426-x](https://doi.org/10.1007/s00348-012-1426-x).

C.-Y. Chen
Department of Mechanical Engineering, National Taiwan
University of Science and Technology, Taipei 10607, Taiwan

P. G. Menon · W. Kowalski · K. Pekkan
Departments of Biomedical and Mechanical Engineering,
Carnegie Mellon University, 700 Technology Drive,
Pittsburgh, PA 15219, USA

K. Pekkan (✉)
Department of Mechanical Engineering, Koc University,
Rumelifeneri, Istanbul, Turkey
e-mail: kpekk@andrew.cmu.edu; kpekk@ku.edu.tr