



A lower bound for the dimension of tetrahedral splines in large degree

Michael DiPasquale¹ · Nelly Villamizar²

Received: 3 March 2021 / Revised: 26 June 2022 / Accepted: 7 July 2022 / Published online: 12 June 2023
© The Author(s) 2023

Abstract

Splines are piecewise polynomial functions which are continuously differentiable to some order r . For a fixed integer d the space of splines of degree at most d is a finite dimensional vector space, and a largely open problem in numerical analysis is to determine its dimension. While considerable attention has been given to this problem in the bivariate setting, the literature on trivariate splines is less conclusive. In particular, the dimension of generic trivariate splines is not known even in large degree when $r > 1$. In this paper we use a bound we previously derived for splines on vertex stars to compute a new lower bound on the dimension of trivariate splines in large enough degree. We illustrate in several examples that our formula gives the exact dimension of the spline space in large enough degree if vertex positions are generic. In contrast, for splines continuously differentiable of order $r > 1$, every lower bound in the literature diverges (often significantly) in large degree from the dimension of the spline space in these examples. We derive the bound using commutative and homological algebra.

Keywords Trivariate spline spaces · Tetrahedral partitions · Dimension of spline spaces

Mathematics Subject Classification 65D07 · 41A15 · 13D02

Communicated by Larry L. Schumaker.

✉ Nelly Villamizar
n.y.villamizar@swansea.ac.uk
<https://sites.google.com/site/nvillami>

Michael DiPasquale
mdipasquale@southalabama.edu
<https://midipasq.github.io>

¹ Department of Mathematics and Statistics, University of South Alabama, Mobile, USA

² Department of Mathematics, Swansea University, Swansea, UK

1 Introduction

A multivariate spline is a piecewise polynomial function on a partition Δ of some domain $\Omega \subset \mathbb{R}^n$ which is continuously differentiable to order r for some integer $r \geq 0$. Multivariate splines play an important role in many areas such as finite elements, computer-aided geometric design, isogeometric analysis, and data fitting [13, 24]. Splines on both triangulations and tetrahedral partitions have been used to solve boundary value problems by the finite element method; some early references are [14, 34, 35], see also [24] and the references therein. For quite recent applications in isogeometric analysis, in [18, 19], Engvall and Evans outline frameworks to parametrize volumes for isogeometric analysis using triangular and tetrahedral Bézier elements. While Engvall and Evans in [19] focus on C^0 elements, C^r tetrahedral Bézier elements are also used for isogeometric analysis—see Xia and Qiang [38]. In these applications it is important to construct a basis, often with prescribed properties, for splines of bounded total degree. Thus it is important to compute the dimension of the space of multivariate splines of bounded degree on a fixed partition. We write $\mathcal{S}_d^r(\Delta)$ for the vector space of piecewise polynomial functions of degree at most d on the partition Δ which are continuously differentiable of order r . By an abuse of notation we identify Δ with its embedding in \mathbb{R}^n .

A formula for the dimension of C^1 splines on triangulations was proposed by Strang [34] and proved for *generic* triangulations by Billera [9]. Subsequently the problem of computing the dimension of planar splines on triangulations has received considerable attention using a wide variety of techniques, see [3, 4, 9, 11, 22, 23, 31–33, 36, 37]. Alfeld and Schumaker show in [4] that the dimension of $\mathcal{S}_d^r(\Delta)$, for (most) planar triangulations Δ and $d \geq 3r + 1$, is given by a quadratic polynomial in d whose coefficients are determined from simple data of the triangulation. The computation of $\dim \mathcal{S}_d^r(\Delta)$ for planar Δ when $r + 1 \leq d \leq 3r$ remains an open problem, although Whiteley has shown that there are only trivial splines on Δ in degrees at most $\frac{3r+1}{2}$ if Δ is generic with a triangular boundary [36]. (This result of Whiteley is an essential ingredient of our lower bound for trivariate splines.)

The literature on computing the dimension of trivariate splines on tetrahedral partitions is much less conclusive. The dimension has been computed if $r = 0$ (see [6] or [10]), and also if $r = 1$, $d \geq 8$, and Δ is generic by Alfeld, Schumaker, and Whiteley [7]. For $r > 1$ bounds on $\dim \mathcal{S}_d^r(\Delta)$ have been computed in [1, 5, 25, 28]. A major difficulty is that computing $\dim \mathcal{S}_d^r(\Delta)$ exactly in large degree for arbitrary tetrahedral partitions cannot be done without computing the dimension of splines on planar triangulations exactly in *all* degrees (see [7, Remark 65]). More precisely, to compute $\dim \mathcal{S}_d^r(\Delta)$ exactly for $d \gg 0$, we must be able to compute the space of *homogeneous* splines $\dim \mathcal{H}_d^r(\Delta_\gamma)$ exactly in all degrees, where γ is a vertex of Δ and Δ_γ is the *star* of γ (that is, Δ_γ consists of all tetrahedra having γ as a vertex). The computation of such spline spaces has only been made for $r \leq 1$; for $r = 1$, the partition Δ is required to be generic [7]. For this crucial computation we rely on our previous paper [17], where we establish a lower bound on the dimension of homogeneous splines on vertex stars.




Fig. 1 A three-dimensional version of the Morgan–Scott triangulation, the star of the boundary vertex γ' (center), and the star of the interior vertex γ (right)

In our main result, Theorem 2.6, we establish a formula which is a lower bound on the dimension of the spline space on most tetrahedral partitions of interest (any triangulation of a compact three-manifold with boundary) in large enough degree. While we have no proof of what degree is large enough, empirical evidence suggests that, for generic Δ , our formula begins to be a lower bound in degrees close to the *initial degree* of $S^r(\Delta)$; by the initial degree of $S^r(\Delta)$ we mean the smallest degree d in which $S_d^r(\Delta)$ admits a spline which is not globally polynomial. If Δ is generic, our formula gives the exact dimension of $S_d^r(\Delta)$ beginning at the initial degree of $S_d^r(\Delta)$ for several tetrahedral partitions considered by Alfeld and Schumaker [5] (see Example 1.1 and Sect. 5). It is worth noting that none of the lower bounds in the literature [5, 25, 28] give the exact dimension of the generic spline space (even in large degree) on these tetrahedral partitions for $r \geq 2$. Below is an example, discussed in detail later in the paper, comparing our results to work by Alfeld and Schumaker in [5] and Mourrain and Villamizar in [28].

Example 1.1 Let Δ be the tetrahedral partition on the left of Figure 1. In Table 1 we record the values of the lower bounds on $\dim S_d^r(\Delta)$ for order of smoothness $2 \leq r \leq 4$. In column 3 we give the dimension of the space of polynomials of degree at most d (this is $\binom{d+3}{3}$), in columns 4–6 the bounds are obtained by applying the formulas proved in [28, Theorem 5.1], [5, Example 8.2], and our new lower bound $LB(d)$ (proved in Theorem 2.6 below), respectively. The last column records the value for the exact dimension for the given order of continuity r , degree d , and generic vertex positions. The bolded entry in the d column indicates the initial degree of $S_d^r(\Delta)$.

The paper is organized as follows. In Sect. 2 we explicitly state our lower bound in purely numerical terms allowing a straightforward application of the formula and illustrate in an example. In Sect. 3 we set up notation and give relevant homological background, and in Sect. 4 we prove the bound of Theorem 2.6. Section 5 is devoted to illustrating our bounds in a number of examples and comparing them to the bounds in [5, 28]. Finally, we give some concluding remarks in Sect. 6. We draw special attention to Remark 6.2, as we think it likely that work of Alfeld, Schumaker, and Sirvent [6] implies that our formula is a lower bound in degrees at least $8r + 1$. Our methods are sufficiently different from [6] that we do not attempt to prove this here.

Table 1 Lower bounds on $\dim S_d^r(\Delta)$, where Δ is the three dimensional Morgan–Scott partition in Fig. 1; see Example 5.1. The initial degree is bolded

r	d	$\binom{d+3}{3}$	LB[28]	LB[5]	LB(d)	gendim
2	5	56	56	56	48	56
2	6	84	84	84	72	84
2	7	120	120	120	132	132
2	8	165	165	207	243	243
2	9	220	320	384	420	420
3	8	165	165	165	137	165
3	9	220	220	220	208	220
3	10	286	286	286	332	332
3	11	364	364	364	524	524
3	12	455	591	593	799	799
3	13	560	964	948	1172	1172
4	11	364	364	364	308	364
4	12	455	455	455	439	455
4	13	560	560	560	640	640
4	14	680	680	680	926	926
4	15	680	896	832	1312	1312

2 The lower bound

Throughout we let Δ be a tetrahedral partition. We are more precise in Sect. 3; for now it is sufficient for the reader to think of a tetrahedral partition as a triangulation of a three-dimensional polytope. We use Δ_i and Δ_i° to denote the i -faces and interior i -faces (respectively) of Δ . We put $f_i(\Delta) = |\Delta_i|$ and $f_i^\circ(\Delta) = |\Delta_i^\circ|$ (if Δ is clear we simply write f_i and f_i°). We define the following data for each edge (Notation 2.1) and for each vertex of Δ (Notation 2.3).

Notation 2.1 [Data attached to edges] For a given $r \geq 0$ and $\tau \in \Delta_1$,

- we define $t_\tau = \min\{n_\tau, r + 2\}$, where $n_\tau = \#\{\sigma \in \Delta_2 : \tau \subset \sigma\}$ is the number of two-dimensional faces having τ as an edge;
- and the constants

$$q_\tau = \left\lfloor \frac{t_\tau(r+1)}{t_\tau-1} \right\rfloor, \quad a_\tau = t_\tau(r+1) - (t_\tau-1)q_\tau, \quad \text{and} \quad b_\tau = t_\tau - 1 - a_\tau.$$

Notice that $t_\tau(r+1) = q_\tau(t_\tau-1) + a_\tau$; i.e., q_τ and a_τ are, respectively, the quotient and remainder obtained when dividing $t_\tau(r+1)$ by $t_\tau-1$.

Given a vertex $\gamma \in \Delta$, we call the set of tetrahedra of Δ which contain γ the *star of γ* and we denote this tetrahedral partition by Δ_γ . If γ is an interior vertex of Δ , so γ is completely surrounded by tetrahedra, then we call Δ_γ a *closed* vertex star. If γ is a boundary vertex of Δ , so γ is not completely surrounded by tetrahedra, then we call Δ_γ an *open* vertex star.

The following convention for binomial coefficients is crucial in all our formulas.

Convention 2.2 For binomial coefficients we always put $\binom{n}{k} = 0$ when $n < k$.

We make the following definitions following Notation 2.1 and Convention 2.2.

Notation 2.3 [Data attached to vertices] For given integers $d \geq r \geq 0$ and $\gamma \in \Delta_0$,

- if Δ_γ is a closed star i.e., $\gamma \in \Delta_0^\circ$, we define

$$D_\gamma := \begin{cases} 2r & \text{if } f_1^\circ(\Delta_\gamma) = 4 \\ \lfloor (5r + 2)/3 \rfloor & \text{if } f_1^\circ(\Delta_\gamma) = 5 \\ \lfloor (3r + 1)/2 \rfloor & \text{if } f_1^\circ(\Delta_\gamma) \geq 6, \end{cases} \tag{1}$$

and

$$\begin{aligned} \text{LB}^\star(\Delta_\gamma, d, r) := & 2 \binom{d+2}{2} + \left(f_2^\circ(\Delta_\gamma) - \sum_{\tau \in (\Delta_\gamma)_1^\circ} t_\tau \right) \binom{d+1-r}{2} \\ & + \sum_{\tau \in (\Delta_\gamma)_1^\circ} \left(a_\tau \binom{d+1-q_\tau}{2} + b_\tau \binom{d+2-q_\tau}{2} \right). \end{aligned} \tag{2}$$

We write $\text{LB}^\star(d)$ instead of $\text{LB}^\star(\Delta, d, r)$ if Δ and r are understood.

- If Δ_γ is an open vertex star i.e., if $\gamma \in \Delta_0 \setminus \Delta_0^\circ$, we define

$$\begin{aligned} \text{LB}^\star(\Delta_\gamma, d, r) := & \binom{d+2}{2} + \left(f_2^\circ(\Delta_\gamma) - \sum_{\tau \in (\Delta_\gamma)_1^\circ} t_\tau \right) \binom{d+1-r}{2} \\ & + \sum_{\tau \in (\Delta_\gamma)_1^\circ} \left(a_\tau \binom{d+1-q_\tau}{2} + b_\tau \binom{d+2-q_\tau}{2} \right). \end{aligned} \tag{3}$$

Again we write $\text{LB}^\star(d)$ if Δ and r are understood.

- For each vertex $\gamma \in \Delta_0$ we define the constant N_γ as follows

$$N_\gamma = \begin{cases} \sum_{d=r+1}^{D_\gamma} \left[\binom{d+2}{2} - \text{LB}^\star(d) \right] + \sum_{d=D_\gamma+1}^{3r+1} \left[\binom{d+2}{2} - \text{LB}^\star(d) \right]_+ & \text{if } \gamma \in \Delta_0^\circ \\ \sum_{d=r+1}^{3r+1} \left[\binom{d+2}{2} - \text{LB}^\star(d) \right]_+ & \text{if } \gamma \in \Delta_0 \setminus \Delta_0^\circ \end{cases} \tag{4}$$

where for a real number m , we put $[m]_+ = \max\{m, 0\}$. The constants $D_\gamma, \text{LB}^\star(d)$, and $\text{LB}^\star(d)$ are those defined in Equations (1), (2), and (3), respectively.

Remark 2.4 In [17] we show $\text{LB}^\star(\Delta, d, r)$ is a lower bound for homogeneous splines on a generic closed vertex star for $d > D_\gamma$ and [2] shows there is equality for $d \geq 3r + 2$. In [2] it is shown that $\text{LB}^\star(\Delta, d, r)$ is a lower bound for homogeneous splines on a generic open vertex star, with equality if $d \geq 3r + 2$.

Remark 2.5 When $\gamma \in \Delta_0^\circ$ and $r + 1 \leq d \leq D_\gamma$, notice that the contribution to N_γ can be negative, while if $d > D_\gamma$, only positive contributions are counted. This is a crucial difference between the contributions from interior vertices and the contributions from boundary vertices.

Theorem 2.6 [Lower bound in large degree for tetrahedral partitions] Suppose Δ is a tetrahedral partition. If $d \gg 0$ then $\dim S_d^r(\Delta) \geq \text{LB}(\Delta, d, r)$, where

$$\begin{aligned} \text{LB}(\Delta, d, r) := & (f_3 - f_2^\circ + f_1^\circ) \binom{d+3}{3} + \left(f_2^\circ - \sum_{\tau \in \Delta_1^\circ} t_\tau \right) \binom{d+2-r}{3} \\ & + \sum_{\tau \in \Delta_1^\circ} \left(a_\tau \binom{d+2-q_\tau}{3} + b_\tau \binom{d+3-q_\tau}{3} \right) \\ & - f_0^\circ \binom{r+3}{3} + \sum_{\gamma \in \Delta_0} N_\gamma. \end{aligned} \quad (5)$$

If Δ and r are understood then we abbreviate $\text{LB}(\Delta, d, r)$ to $\text{LB}(d)$.

2.1 Example

We illustrate Theorem 2.6 for C^2 splines on the tetrahedral partition in Fig. 1, which is a three-dimensional analog of the Morgan–Scott triangulation [27]. If γ is an interior vertex then Δ_γ is the triangulated octahedron on the right in Fig. 1. We have $f_0^\circ(\Delta_\gamma) = 1$, $f_1^\circ(\Delta_\gamma) = 6$, and $f_2^\circ(\Delta_\gamma) = 12$. For every $\tau \in (\Delta_\gamma)_1^\circ$, we have $n_\tau = 4$ and hence $t_\tau = \min\{n_\tau, r + 2\} = 4$. We compute $q_\tau = 4$, $a_\tau = 0$, and $b_\tau = 3$, hence by Equation (2),

$$\text{LB}^\star(\Delta_\gamma, d, 2) = 2 \binom{d+2}{2} - 12 \binom{d-1}{2} + 18 \binom{d-2}{2}.$$

If γ' is a boundary vertex, then $\Delta_{\gamma'}$ is the cone over the Morgan–Scott triangulation (see the star of vertex γ' in Fig. 1). We have $f_0^\circ(\Delta_{\gamma'}) = 0$, $f_1^\circ(\Delta_{\gamma'}) = 3$, and $f_2^\circ(\Delta_{\gamma'}) = 9$. For every $\tau \in (\Delta_{\gamma'})_1^\circ$, we have $n_\tau = 4$ and hence $t_\tau = \min\{n_\tau, r + 2\} = 4$. Again we have $q_\tau = 4$, $a_\tau = 0$, and $b_\tau = 3$. Thus, following Equation (3),

$$\text{LB}^\star(\Delta_{\gamma'}, d, 2) = \binom{d+2}{2} - 3 \binom{d-1}{2} + 9 \binom{d-2}{2}.$$

In Table 2 we record the values of $\text{LB}^\star(\Delta_\gamma, d, 2)$, $\text{LB}^\star(\Delta_{\gamma'}, d, 2)$, and $\binom{d+2}{2}$ where γ is an interior vertex of Δ and γ' is a boundary vertex of Δ .

Now we turn to computing the bound $\text{LB}(\Delta, d, 2)$ in Theorem 2.6 for $\dim S_d^2(\Delta)$, where Δ is the full simplicial complex depicted in Fig. 1. If γ is a boundary vertex then $N_\gamma = 3$ (corresponding to the one difference in degree 3 in Table 2). If γ is an interior

Table 2 Lower bounds for the star of an interior (γ) and boundary (γ') vertex of the simplicial complex in Fig. 1

d	3	4	5	6	7	8	9	10
$\binom{d+2}{2}$	10	15	21	28	36	45	55	66
$\text{LB}^\star(\Delta_\gamma, d, 2)$	8	12	24	44	72	108	152	204
$\text{LB}^{\star\star}(\Delta_{\gamma'}, d, 2)$	7	15	30	52	81	117	160	210

Table 3 Illustrating Theorem 2.6 for the tetrahedral partition in Fig. 1

d	0	1	2	3	4	5	6	7	8	9	10
$\binom{d+3}{3}$	1	4	10	20	35	56	84	120	165	220	286
$\text{LB}(\Delta, d, 2)$	-57	12	42	48	45	48	72	132	243	420	678
$\dim S_d^2(\Delta)$	1	4	10	20	35	56	84	132	243	420	678

vertex then $D_\gamma = 3$. Reading down each column in the first two rows of Table 2 we get $N_\gamma = (10 - 8) + (15 - 12) = 5$. Thus $\sum_{\gamma \in \Delta_0} N_\gamma = 4 \cdot 3 + 4 \cdot (5) = 32$.

For the remaining statistics we have $f_0^\circ = 4, f_1^\circ = 18, f_2^\circ = 28, \text{ and } f_3 = 15$. For each interior 1-face τ we have $n_\tau = t_\tau = 4, d_\tau = 4, a_\tau = 0, b_\tau = 3$. Thus, by Theorem 2.6,

$$\text{LB}(\Delta, d, 2) = 5 \binom{d+3}{3} - 44 \binom{d}{3} + 54 \binom{d-1}{3} - 8 = \frac{5}{2}d^3 - 27d^2 + \frac{187}{2}d - 57,$$

where the second equality holds as long as $d \geq 1$. Table 3 compares the values of $\text{LB}(\Delta, d, 2)$ and $\dim S_d^2(\Delta)$ for generic positions of the vertices of Δ . Notice that while $\text{LB}(\Delta, d, 2)$ is neither an upper or lower bound for $d \leq 6$, it predicts the correct dimension of the generic spline space for $d \geq 7$. Incidentally, $d = 7$ is the initial degree of $S^2(\Delta)$; that is, the first non-trivial splines appear in degree 7. We computed the exact dimension of the spline space for generic vertex positions using the Algebraic Splines package in Macaulay2 [20]. Furthermore, a computation in Macaulay2 shows that $\dim S_d^2(\Delta) = \frac{5}{2}d^3 - 27d^2 + \frac{187}{2}d - 57$ for $d \gg 0$, so our lower bound gives the exact dimension of the spline space for $r = 2$ when $d \geq 7$. Code to compute all examples in this paper can be found on the first author’s website under the Research tab: <https://midipasq.github.io/>.

3 Background and homological methods

In this section we introduce the homological methods of Billera [9] and Schenck and Stillman [32]. A *simplex* in \mathbb{R}^n is the convex hull of $i \leq n + 1$ vertices which are in linearly general position (no three on a line, no four on a plane, etc.). A *face* of a simplex is the convex hull of any subset of the vertices which define it (thus a face of a simplex is a simplex). An i -simplex (or i -face) is the convex hull of $i + 1$ vertices in linearly general position; i is the *dimension* of the i -simplex or i -face.

Definition 3.1 A *simplicial complex* Δ is a collection of simplices in \mathbb{R}^n satisfying:

- If $\beta \in \Delta$ then so are all of its faces.
- If $\beta_1, \beta_2 \in \Delta$ then $\beta_1 \cap \beta_2$ is either empty or a proper face of both β_1 and β_2 .

We also refer to the simplices of Δ as *faces* of Δ . The *dimension* of Δ is the dimension of a maximal simplex of Δ under inclusion. If all maximal simplices have equal dimension we say that Δ is *pure*.

In this paper we only consider *finite* simplicial complexes. If β is a face of Δ of dimension i we call β an i -face. Denote by Δ_i and Δ_i° the set i -faces of Δ and interior i -faces of Δ , respectively. We write $f_i(\Delta)$ and $f_i^\circ(\Delta)$ for the number of i -faces and interior i -faces, respectively (we write f_i and f_i° if Δ is understood). By an abuse of notation, we will identify Δ with its underlying space $\bigcup_{\beta \in \Delta} \beta \subset \mathbb{R}^n$.

Definition 3.2 If Δ is a simplicial complex and β is a face of Δ , then the *link* of β is the set of all simplices γ in Δ so that $\beta \cap \gamma = \emptyset$ and $\beta \cup \gamma$ is a face of Δ . The *star* of β is the union of the link of β with the set of all simplices which contain β (including β). We denote the star of β by Δ_β .

If γ is a vertex of a simplicial complex Δ so that all maximal simplices of Δ contain γ (so $\Delta_\gamma = \Delta$), then we call Δ the *star* of γ and we say Δ is a *vertex star*. If γ is an *interior* vertex we call Δ a closed vertex star and if γ is a boundary vertex then we call Δ an open vertex star.

We refer to the set of points in \mathbb{R}^{n+1} of unit norm as the n -sphere, and the set of points in \mathbb{R}^n with norm at most one as the n -disk. A *homeomorphism* $f : X \rightarrow Y$ between two sets is a continuous bijection; if such an f exists we say X and Y are *homeomorphic*.

Definition 3.3 (Simplicial n -manifold with boundary) If Δ is a finite simplicial complex in \mathbb{R}^n , we say it is a *simplicial n -manifold* with boundary if it satisfies the conditions:

- Δ is pure n -dimensional,
- the link of every vertex of Δ is homeomorphic to an $(n - 1)$ -sphere (if the vertex is *interior*) or an $(n - 1)$ -disk (if the vertex is on the boundary),
- and every $(n - 1)$ -simplex of Δ is either the intersection of two n -simplices of Δ or it is on the boundary of Δ and so contained in only one n -simplex of Δ .

Example 3.4 Consider the simplicial complex in Fig. 1, which is a simplicial 3-manifold with boundary homeomorphic to the 3-disk. The star of the interior vertex γ is shown in the center of Fig. 1; the link of the vertex γ is obtained from the star of γ by removing γ and all simplices which contain it. The link of γ is homeomorphic to a 2-sphere. Likewise, the star of the boundary vertex γ' is shown on the right in Fig. 1; the link of the vertex γ' is obtained from it by removing the vertex γ' and all simplices which contain it. The link of γ' is the usual planar Morgan–Scott configuration [27], and is homeomorphic to a 2-disk.

Throughout this paper we abuse notation by referring to a simplicial n -manifold with boundary simply as a simplicial complex. We refer to a simplicial 2-manifold

with boundary as a *triangulation* and a simplicial 3-manifold with boundary as a *tetrahedral partition*.

Write $S = \mathbb{R}[x_1, \dots, x_n]$ for the polynomial ring in n variables and $S_{\leq d}$ for the \mathbb{R} -vector space of polynomials of total degree most d , and S_d for the \mathbb{R} -vector space of polynomials which are homogeneous of degree exactly d . For a fixed integer r , we denote by $C^r(\Delta)$ the set of all functions $F : \Delta \rightarrow \mathbb{R}$ which are continuously differentiable of order r .

Definition 3.5 Let $\Delta \subset \mathbb{R}^n$ be an n -dimensional simplicial complex. We denote by

$$S^r(\Delta) := \{F \in C^r(\Delta) : F|_\iota \in S \text{ for all } \iota \in \Delta_n\}$$

the vector space of splines which are continuously differentiable of order r , by

$$S_d^r(\Delta) := \{F \in C^r(\Delta) : F|_\iota \in S_{\leq d} \text{ for all } \iota \in \Delta_n\}$$

the subspace of $S^r(\Delta)$ consisting of splines of degree at most d , and by

$$\mathcal{H}_d^r(\Delta) := \{F \in C^r(\Delta) : F|_\iota \in S_d \text{ for all } \iota \in \Delta_n\}$$

the subspace of $S^r(\Delta)$ consisting of splines whose restriction to each n -dimensional simplex is a homogeneous polynomial of degree d . We call splines in $\mathcal{H}_d^r(\Delta)$ *homogeneous splines*.

Suppose Δ is the star of a vertex. By changing coordinates, we will assume that this vertex is the origin in \mathbb{R}^n . Then one can show that

$$S^r(\Delta) \cong \bigoplus_{i \geq 0} \mathcal{H}_i^r(\Delta), \text{ and } S_d^r(\Delta) \cong \bigoplus_{i=0}^d \mathcal{H}_i^r(\Delta), \tag{6}$$

where the isomorphism is as \mathbb{R} -vector spaces. We refer to the first isomorphism in (6) as the *graded structure* of $S^r(\Delta)$. If Δ is not the star of a vertex, then (6) does not hold for $S^r(\Delta)$; we summarize a coning construction of Billera and Rose under which (6) will still be valid.

Construction 3.6 Let \mathbb{R}^n have coordinates x_1, \dots, x_n , \mathbb{R}^{n+1} have coordinates x_0, \dots, x_n , and define $\phi : \mathbb{R}^n \rightarrow \mathbb{R}^{n+1}$ by $\phi(x_1, \dots, x_n) = (1, x_1, \dots, x_n)$. If σ is a simplex in \mathbb{R}^n , the *cone over* σ , denoted $\widehat{\sigma}$, is the simplex in \mathbb{R}^{n+1} which is the convex hull of the origin in \mathbb{R}^{n+1} and $\phi(\sigma)$. If $\Delta \subset \mathbb{R}^n$ is a simplicial complex, the *cone over* Δ , denoted $\widehat{\Delta}$, is the simplicial complex consisting of the simplices $\{\widehat{\beta} : \beta \in \Delta\}$ along with the origin in \mathbb{R}^{n+1} , which is called the *cone vertex*. We denote the polynomial ring $\mathbb{R}[x_0, x_1, \dots, x_n]$ associated to $\widehat{\Delta}$ by \widehat{S} .

For any simplicial complex $\Delta \subset \mathbb{R}^n$, the simplicial complex $\widehat{\Delta} \subset \mathbb{R}^{n+1}$ is an (open) vertex star of the cone vertex. Thus (6) yields $S^r(\widehat{\Delta}) \cong \bigoplus_{i \geq 0} \mathcal{H}_i^r(\widehat{\Delta})$ and $S_d^r(\widehat{\Delta}) \cong$

$\bigoplus_{i=0}^d \mathcal{H}_i^r(\widehat{\Delta})$. Moreover, Billera and Rose show that

Theorem 3.7 [11, Theorem 2.6] $S_d^r(\Delta) \cong \mathcal{H}_d^r(\widehat{\Delta})$.

Thus the study of spline spaces reduces to the study of homogeneous spline spaces.

Definition 3.8 A subset $\mathbb{I} \subset S$ is called an *ideal* if, for every $f, g \in \mathbb{I}$ and $h \in S$, $f + g \in \mathbb{I}$ and $hf \in \mathbb{I}$. If $f_1, \dots, f_k \in S$ are polynomials, we write $\langle f_i \rangle$ for the vector space of all polynomial multiples of f_i ($i = 1, \dots, k$) and $\langle f_1, \dots, f_k \rangle := \sum_{i=1}^k \langle f_i \rangle$. This is called the *ideal generated by* f_1, \dots, f_k . We typically only use its vector space structure.

Definition 3.9 Suppose $\Delta \subset \mathbb{R}^n$ is an n -dimensional simplicial complex. If $\beta \in \Delta_n$ we define $\mathcal{J}(\beta) = 0$. If $\sigma \in \Delta_{n-1}$, let ℓ_σ be a choice of linear form vanishing on σ . We define $\mathcal{J}(\sigma) = \langle \ell_\sigma^{r+1} \rangle$. For any face $\beta \in \Delta_i$ where $i < n$ we define

$$\mathcal{J}(\beta) := \sum_{\sigma \supseteq \beta} \mathcal{J}(\sigma) = \langle \ell_\sigma^{r+1} : \beta \subseteq \sigma \rangle.$$

Billera and Rose show that if Δ is *hereditary* (a hypothesis which is implied by ours) then

Proposition 3.10 [11, Proposition 1.2] $F \in S^r(\Delta)$ if and only if

$$F|_\iota - F|_{\iota'} \in \mathcal{J}(\sigma) \text{ for every } \iota, \iota' \in \Delta_n \text{ satisfying } \iota \cap \iota' = \sigma \in \Delta_{n-1}.$$

3.1 Chain complexes

If C_0, \dots, C_k are vector spaces and $\partial_i : C_i \rightarrow C_{i-1}$ ($i = 1, \dots, k$) are linear maps satisfying $\partial_{i-1} \circ \partial_i = 0$ (for $i = 2, \dots, k$), then the collection of this data is called a *chain complex*; this is typically recorded as

$$C : 0 \rightarrow C_k \xrightarrow{\partial_k} C_{k-1} \xrightarrow{\partial_{k-1}} \dots \xrightarrow{\partial_1} C_0 \rightarrow 0.$$

We call the subscript i of C_i the *homological index* and refer to C_i as the vector space of C in homological index i . The *homologies* of the chain complex are the quotient vector spaces $H_i(C) = \ker(\partial_i) / \text{im}(\partial_{i+1})$ for $i = 0, \dots, k$. (We put $H_0(C) = C_0 / \text{im}(\partial_1)$ and $H_k(C) = \ker(\partial_k)$.) Often $H_*(C)$ is used to denote the entire set of homology groups $H_0(C), \dots, H_k(C)$. We are primarily concerned with a topological construction of chain complexes; see [21, Chapter 2] for a standard reference.

We now define the chain complex introduced by Billera [9] and refined by Schenck and Stillman [32]. Let S^{Δ_i} ($i = 0, \dots, n$) denote the direct sum $\bigoplus_{\beta \in \Delta_i} S[\beta]$, where $[\beta]$ is a formal basis symbol corresponding to the i -face β . Fix an ordering $\gamma_1, \dots, \gamma_{f_0}$ of the vertices of Δ . Each i -face $\beta \in \Delta_i$ can be represented as an *ordered list* $\beta = (\gamma_{j_0}, \dots, \gamma_{j_i})$ of $i + 1$ vertices. We define the *simplicial boundary map* ∂_i (for $i = 1, 2, 3$) on the formal symbol $[\beta] = [\gamma_{j_0}, \dots, \gamma_{j_i}]$ by $\partial_i([\beta]) = \partial_i([\gamma_{j_0}, \dots, \gamma_{j_i}]) = \sum_{k=0}^i (-1)^k [\gamma_{j_0}, \dots, \hat{\gamma}_{j_k}, \dots, \gamma_{j_i}]$, where $\hat{\gamma}_{j_k}$ means that the vertex γ_{j_k} is omitted from the list. We extend this map linearly to $\bigoplus_{\beta \in \Delta_i} S[\beta]$.

It is straightforward to verify that $\partial_{i-1} \circ \partial_i = 0$ for $i = 2, \dots, n$ (this only needs to be checked on the basis symbols $[\beta]$). Clearly the simplicial boundary map ∂_i can be restricted to a map $\partial_i : S^{\Delta_i^\circ} \rightarrow S^{\Delta_{i-1}^\circ}$ where all formal symbols corresponding to faces on the boundary of Δ are dropped. We denote by $\mathcal{R}[\Delta]$ the chain complex

$$\mathcal{R}[\Delta]: 0 \longrightarrow S^{\Delta_n} \xrightarrow{\partial_n} S^{\Delta_{n-1}^\circ} \xrightarrow{\partial_{n-1}} \dots \xrightarrow{\partial_2} S^{\Delta_1^\circ} \xrightarrow{\partial_1} S^{\Delta_0^\circ} \longrightarrow 0.$$

(This is the simplicial chain complex of Δ relative to its boundary $\partial\Delta$ with coefficients in S —see [21, Chapter 2.1]).

We now put the vector spaces $\mathcal{J}(\beta)$ together to make a sub-chain complex of $\mathcal{R}[\Delta]$

$$\begin{aligned} \mathcal{J}[\Delta]: 0 \longrightarrow \bigoplus_{\iota \in \Delta_n} \mathcal{J}(\iota) = 0 &\longrightarrow \bigoplus_{\sigma \in \Delta_{n-1}^\circ} \mathcal{J}(\sigma) \xrightarrow{\partial_{n-1}} \dots \xrightarrow{\partial_2} \\ \bigoplus_{\tau \in \Delta_1^\circ} \mathcal{J}(\tau) \xrightarrow{\partial_1} \bigoplus_{\gamma \in \Delta_0^\circ} \mathcal{J}(\gamma) &\longrightarrow 0. \end{aligned}$$

The *Billera-Schenck-Stillman chain complex* is the quotient of $\mathcal{R}[\Delta]$ by $\mathcal{J}[\Delta]$, namely

$$\begin{aligned} \mathcal{R}/\mathcal{J}[\Delta]: 0 \longrightarrow \bigoplus_{\iota \in \Delta_n} S \xrightarrow{\bar{\partial}_n} \bigoplus_{\sigma \in \Delta_{n-1}^\circ} \frac{S}{\mathcal{J}(\sigma)} \xrightarrow{\bar{\partial}_{n-1}} \dots \xrightarrow{\bar{\partial}_2} \\ \bigoplus_{\tau \in \Delta_1^\circ} \frac{S}{\mathcal{J}(\tau)} \xrightarrow{\bar{\partial}_1} \bigoplus_{\gamma \in \Delta_0^\circ} \frac{S}{\mathcal{J}(\gamma)} \longrightarrow 0. \end{aligned}$$

Remark 3.11 If the simplicial complex Δ is fixed, we simply write \mathcal{J}, \mathcal{R} , and \mathcal{R}/\mathcal{J} for the chain complexes $\mathcal{J}[\Delta], \mathcal{R}[\Delta]$, and $\mathcal{R}/\mathcal{J}[\Delta]$, respectively.

Notation 3.12 We introduce a natural abuse of notation regarding the coning construction 3.6. If Δ is a simplicial complex and $\widehat{\Delta}$ is the cone over Δ , then $\widehat{\Delta}$ is an open vertex star. Hence there is no interior vertex of $\widehat{\Delta}$ and thus the vector space of homological index 0 in $\mathcal{J}[\widehat{\Delta}], \mathcal{R}[\widehat{\Delta}]$, and $\mathcal{R}/\mathcal{J}[\widehat{\Delta}]$ is just zero. We thus decrease the homological index by one of each of the vector spaces in $\mathcal{J}[\widehat{\Delta}], \mathcal{R}[\widehat{\Delta}]$, and $\mathcal{R}/\mathcal{J}[\widehat{\Delta}]$. Hence if $\Delta \subset \mathbb{R}^n$ and thus $\widehat{\Delta} \subset \mathbb{R}^{n+1}$, $H_n(\mathcal{R}/\mathcal{J}[\widehat{\Delta}])$ is the top homology of the chain complex $\mathcal{R}/\mathcal{J}[\widehat{\Delta}]$, not $H_{n+1}(\mathcal{R}/\mathcal{J}[\widehat{\Delta}])$ (and likewise for lower indices). Thus the vector space in homological index i ($0 \leq i \leq n$) in $\mathcal{R}/\mathcal{J}[\widehat{\Delta}]$ corresponds to the homological index i in $\mathcal{R}/\mathcal{J}[\Delta]$, so its summands are indexed by Δ_i° .

The crucial observation of Billera is that $H_n(\mathcal{R}/\mathcal{J}[\Delta]) \cong S^r(\Delta)$; this follows from the criterion of Proposition 3.10.

3.2 Graded structure

The vector space $\mathcal{J}(\beta)$ is infinite-dimensional for each face $\beta \in \Delta$ which is not a tetrahedron. Thus the constituents of the chain complexes $\mathcal{J}[\Delta], \mathcal{R}[\Delta]$, and $\mathcal{R}/\mathcal{J}[\Delta]$

are also infinite-dimensional. In order to get a chain complex of finite dimensional vector spaces to relate to the fundamental spaces of interest ($\mathcal{S}_d^r(\Delta)$ and $\mathcal{H}_d^r(\Delta)$), we make use of a *graded structure*.

Definition 3.13 Let V be a real vector space and suppose V_i is a finite-dimensional vector subspace of V for every integer $i \geq 0$. If $V \cong \bigoplus_{i \geq 0} V_i$, then we refer to this isomorphism as a *graded structure* of V and we call V a *graded vector space*. In particular, if $\mathcal{J} \subset \mathcal{S}$ is an ideal (c.f. Definition 3.8), then we write \mathcal{J}_d for the vector space of homogeneous polynomials of degree d in \mathcal{J} . If $\mathcal{J} \cong \bigoplus_{d \geq 0} \mathcal{J}_d$ then we call \mathcal{J} a *graded ideal* of \mathcal{S} .

Definition 3.14 If $\mathcal{C} : 0 \rightarrow C_n \xrightarrow{\partial_n} \dots \xrightarrow{\partial_1} C_0 \rightarrow 0$ is a chain complex of vector spaces so that

- (1) The vector space C_j has a graded structure $C_j \cong \bigoplus_{i \geq 0} (C_j)_i$ for $j = 0, \dots, n$ and
- (2) The map $\partial_j : C_j \rightarrow C_{j-1}$ satisfies $\partial_j((C_j)_i) \subset (C_{j-1})_i$ for $j = 1, \dots, n$,

then $\mathcal{C}_d := 0 \rightarrow (C_n)_d \xrightarrow{\partial_n} (C_{n-1})_d \xrightarrow{\partial_{n-1}} \dots \xrightarrow{\partial_1} (C_0)_d \rightarrow 0$ is a chain complex which we call the *degree d strand* of \mathcal{C} . In this case we say \mathcal{C} is *graded with graded structure* $\mathcal{C} \cong \bigoplus_{d \geq 0} \mathcal{C}_d$.

If a chain complex \mathcal{C} has a graded structure $\mathcal{C} \cong \bigoplus_{d \geq 0} \mathcal{C}_d$, it is straightforward to see that the homologies of \mathcal{C} also have the graded structure $H_i(\mathcal{C}) \cong \bigoplus_{d \geq 0} H_i(\mathcal{C})_d$, where $H_i(\mathcal{C})_d := H_i(\mathcal{C}_d)$ is the i th homology of the degree d strand.

Remark 3.15 The isomorphisms (6) show that $\mathcal{S}^r(\Delta)$ has a graded structure if Δ is the star of a vertex where the vertex is at the origin in \mathbb{R}^n .

If Δ is a vertex star of γ (which we can assume to be located at the origin, if necessary, by changing coordinates) and $\gamma \in \beta$, then the linear forms whose powers generate $\mathcal{J}(\beta)$ have no constant term and $\mathcal{J}(\beta)$ is a graded ideal. It is straightforward to see that the simplicial boundary map respects this graded structure (i.e. property (2) of Definition 3.14 is satisfied), so if Δ is a vertex star then the chain complexes $\mathcal{J}[\Delta]$, $\mathcal{R}[\Delta]$, and $\mathcal{R}/\mathcal{J}[\Delta]$ also have a graded structure, along with their homologies. In particular, $\mathcal{H}_d^r(\Delta) \cong H_n(\mathcal{R}/\mathcal{J}[\Delta])_d$ if $\Delta \subset \mathbb{R}^n$ is a vertex star. If Δ is not necessarily a vertex star, we use the coning construction $\Delta \rightarrow \widehat{\Delta}$. Then $\widehat{\Delta}$ is a vertex star (whose vertex is at the origin) and so $\mathcal{S}^r(\widehat{\Delta})$, along with $\mathcal{J}[\widehat{\Delta}]$, $\mathcal{R}[\widehat{\Delta}]$, and $\mathcal{R}/\mathcal{J}[\widehat{\Delta}]$, all have a graded structure. Keeping in mind Theorem 3.7 and Notation 3.12, we have $\mathcal{S}_d^r(\Delta) \cong \mathcal{H}_d^r(\widehat{\Delta}) \cong \ker(\bar{\partial}_n)_d \cong H_n(\mathcal{R}/\mathcal{J}[\widehat{\Delta}])_d$.

3.3 Euler characteristic and dimension formulas

If $\mathcal{C} : 0 \rightarrow C_n \rightarrow C_{n-1} \rightarrow \dots \rightarrow C_0 \rightarrow 0$ is a chain complex with a graded structure, we write $\chi(\mathcal{C}, d) = \sum_{i=0}^n (-1)^{n-i} \dim(C_i)_d$. This is the *Euler-Poincaré characteristic* of \mathcal{C}_d . The rank-nullity theorem yields:

$$\chi(\mathcal{C}, d) = \sum_{i=0}^n (-1)^{n-i} \dim H_i(\mathcal{C})_d. \tag{7}$$

The three chain complexes \mathcal{J} , \mathcal{R} , and \mathcal{R}/\mathcal{J} fit into the short exact sequence of chain complexes $0 \rightarrow \mathcal{J} \rightarrow \mathcal{R} \rightarrow \mathcal{R}/\mathcal{J} \rightarrow 0$. Correspondingly there is the long exact sequence:

$$0 \rightarrow H_n(\mathcal{J}) \rightarrow H_n(\mathcal{R}) \rightarrow \dots \rightarrow H_1(\mathcal{R}/\mathcal{J}) \rightarrow H_0(\mathcal{J}) \rightarrow H_0(\mathcal{R}) \rightarrow H_0(\mathcal{R}/\mathcal{J}) \rightarrow 0.$$

The short exact sequence $0 \rightarrow \mathcal{J} \rightarrow \mathcal{R} \rightarrow \mathcal{R}/\mathcal{J} \rightarrow 0$ also yields

$$\chi(\mathcal{R}/\mathcal{J}, d) = \chi(\mathcal{R}, d) + \chi(\mathcal{J}, d). \tag{8}$$

There is a sum instead of a difference on the right hand side of Equation (8) because the first non-zero term in the chain complex \mathcal{J} has homological degree $n - 1$ instead of n .

Proposition 3.16 *For an n -dimensional simplicial complex Δ in \mathbb{R}^n , $H_n(\mathcal{R}/\mathcal{J}[\widehat{\Delta}])_d \cong S_d^r(\Delta)$ and $H_0(\mathcal{R}/\mathcal{J}[\Delta]) = 0$. If Δ is a vertex star, $H_n(\mathcal{R}/\mathcal{J}[\Delta])_d \cong \mathcal{H}_d^r(\Delta)$. If Δ is connected, then $H_0(\mathcal{R}/\mathcal{J}[\Delta]) = 0$. If Δ is a vertex star whose link is homeomorphic to an $(n - 1)$ -sphere or an $(n - 1)$ -disk, then $S^r(\Delta) \cong H_n(\mathcal{R}/\mathcal{J}) \cong S \oplus H_{n-1}(\mathcal{J})$ and $H_i(\mathcal{R}/\mathcal{J}) \cong H_{i-1}(\mathcal{J})$ for $i = 1, \dots, n - 1$.*

Proof By Theorem 3.7 and Proposition 3.10, $S_d^r(\Delta) \cong \mathcal{H}_d^r(\widehat{\Delta}) \cong H_n(\mathcal{R}/\mathcal{J}[\Delta])_d$. Since every vertex can be connected to the boundary of Δ by a path consisting of interior edges, $\partial_1 : S^{f_1} \rightarrow S^{f_0}$ is surjective and thus $H_0(\mathcal{R}[\Delta]) = 0$, hence $H_0(\mathcal{R}/\mathcal{J}[\Delta]) = 0$ by the long exact sequence associated to $0 \rightarrow \mathcal{J} \rightarrow \mathcal{R} \rightarrow \mathcal{R}/\mathcal{J} \rightarrow 0$.

The hypothesis that Δ is a vertex star whose link is homeomorphic to an $(n - 1)$ -sphere or an $(n - 1)$ -disk implies that $H_i(\mathcal{R}[\Delta]) = 0$ for $0 \leq i < n$ and $H_n(\mathcal{R}[\Delta]) \cong S$ (by excision [21, Proposition 2.22], the homology of Δ relative to its boundary coincides with the homology of the n -sphere, which gives the claimed homologies). Then the last result follows from the long exact sequence associated to $0 \rightarrow \mathcal{J} \rightarrow \mathcal{R} \rightarrow \mathcal{R}/\mathcal{J} \rightarrow 0$. □

Remark 3.17 If Δ is homeomorphic to an n -disk, then the copy of S in $S^r(\Delta) \cong S \oplus H_{n-1}(\mathcal{J})$ corresponds to the globally polynomial splines, while the so-called *smoothing cofactors* are encoded by the map

$$\bigoplus_{\sigma \in \Delta_{n-1}^\circ} \mathcal{J}(\sigma) \xrightarrow{\partial_{n-1}} \bigoplus_{\tau \in \Delta_{n-2}^\circ} \mathcal{J}(\beta).$$

Proposition 3.18 *If Δ is a tetrahedral partition then*

$$\begin{aligned} \dim S_d^r(\Delta) &= (f_3 - f_2^\circ + f_1^\circ - f_0^\circ) \dim \widehat{S}_d + \chi(\mathcal{J}[\widehat{\Delta}], d) \\ &+ \dim H_2(\mathcal{R}/\mathcal{J}[\widehat{\Delta}])_d - \dim H_1(\mathcal{R}/\mathcal{J}[\widehat{\Delta}])_d. \end{aligned} \tag{9}$$

If Δ is a tetrahedral vertex star whose link is homeomorphic to a 2-sphere or a 2-disk then

$$\dim \mathcal{H}_d^r(\Delta) = \dim S_d + \chi(\mathcal{J}[\Delta], d) + \dim H_1(\mathcal{J}[\Delta])_d. \tag{10}$$

Proof First we make use of the identifications $\mathcal{S}_d^r(\Delta) \cong \mathcal{H}_d^r(\widehat{\Delta})$ and $H_3(\mathcal{R}/\mathcal{J}[\widehat{\Delta}])_d \cong \mathcal{H}_d^r(\widehat{\Delta})$ of Theorems 3.7 and Proposition 3.16 (using Notation 3.12 for the second isomorphism). The identity (7) applied to the Euler-Poincaré characteristic of $\mathcal{R}/\mathcal{J}[\widehat{\Delta}]$, coupled with Proposition 3.16, gives

$$\dim \mathcal{S}_d^r(\Delta) = \chi(\mathcal{R}/\mathcal{J}[\widehat{\Delta}], d) + \dim H_2(\mathcal{R}/\mathcal{J}[\widehat{\Delta}])_d - \dim H_1(\mathcal{R}/\mathcal{J}[\widehat{\Delta}])_d.$$

To get Equation (9), note that \mathcal{R} has the form $0 \rightarrow S^{f_3} \rightarrow S^{f_2} \rightarrow S^{f_1} \rightarrow S^{f_0} \rightarrow 0$; taking the Euler characteristic in degree d and using Equation (8) yields Equation (9). For Equation (10), Proposition 3.16 implies that $\dim \mathcal{H}_d^r(\Delta) = \dim S_d + \dim H_2(\mathcal{J}[\Delta])_d$. Taking the Euler-Poincaré characteristic of $\mathcal{J}[\Delta]$ gives

$$\dim H_2(\mathcal{J}[\Delta])_d = \chi(\mathcal{J}[\Delta], d) + \dim H_1(\mathcal{J}[\Delta])_d - \dim H_0(\mathcal{J}[\Delta])_d.$$

It is straightforward to show that $H_0(\mathcal{J}[\Delta]) = 0$; putting together the above two equations yields Equation (10). □

3.4 Generic simplicial complexes

It is well-known that, for a fixed r and d , there is an open set in $(\mathbb{R}^n)^{f_0}$ of vertex coordinates of Δ for which $\dim \mathcal{S}_d^r(\Delta)$ is constant.

Definition 3.19 Suppose Δ has vertex coordinates so that $\dim \mathcal{S}_d^r(\Delta) \leq \dim \mathcal{S}_d^r(\Delta')$ for all simplicial complexes Δ' obtained from Δ by a small perturbation of the vertex coordinates. Then we say Δ is *generic* with respect to r and d , or simply *generic* if r and d are understood.

Hence, for the purposes of obtaining a lower bound on $\dim \mathcal{S}_d^r(\Delta)$, it suffices to obtain a lower bound on $\dim \mathcal{S}_d^r(\Delta)$ when Δ is generic.

4 Proof of Theorem 2.6: a lower bound in large degree

To prove Theorem 2.6 we use Equation (9) from Proposition 3.18, so we first describe how to compute the terms which appear in $\chi(\mathcal{J}[\widehat{\Delta}], d)$. From the discussion in Sect. 3.4, it suffices to consider *generic* tetrahedral partitions. First, the Euler characteristic of $\mathcal{J}[\widehat{\Delta}]$ has the form

$$\chi(\mathcal{J}[\widehat{\Delta}], d) = \sum_{\sigma \in \Delta_2^\circ} \dim \mathcal{J}(\widehat{\sigma})_d - \sum_{\tau \in \Delta_1^\circ} \dim \mathcal{J}(\widehat{\tau})_d + \sum_{\gamma \in \Delta_0^\circ} \dim \mathcal{J}(\widehat{\gamma})_d.$$

If Δ is a vertex star with γ placed at the origin, we describe the effect which coning has on the vector spaces $\mathcal{J}(\beta)$, where β is an i -face of Δ . The vector spaces $\mathcal{J}(\beta) \subset \mathcal{S}$ and $\mathcal{J}(\widehat{\beta}) \subset \widehat{\mathcal{S}}$ are related by tensor product. Explicitly, $\mathcal{J}(\widehat{\beta}) \cong \mathcal{J}(\beta) \otimes_{\mathbb{R}} \mathbb{R}[x_0]$ and

$$\dim \mathcal{J}(\widehat{\beta})_d = \sum_{i \leq d} \dim \mathcal{J}(\beta)_i. \tag{11}$$

Hence to compute $\dim \mathcal{J}(\widehat{\beta})_d$ it is necessary and sufficient to compute $\dim \mathcal{J}(\beta)_i$ for every $0 \leq i \leq d$. Since these dimensions are invariant under a translation of \mathbb{R}^3 , we assume β contains the origin and thus $\mathcal{J}(\beta)$ is graded.

Proposition 4.1 *Suppose $\Delta \subset \mathbb{R}^3$ is a tetrahedral partition, $r \geq 0$ is an integer, and $\tau \in \Delta_1$. With t_τ, a_τ , and b_τ as in Notation 2.1, we have*

$$\dim \mathcal{J}(\tau)_d \leq t_\tau \binom{d+1-r}{2} - a_\tau \binom{d+1-q_\tau}{2} - b_\tau \binom{d+2-q_\tau}{2},$$

with equality if every triangle σ containing τ has a distinct linear span (in particular, there is equality if Δ is generic).

Proof This is one of the fundamental computations for planar splines, originally due to Schumaker. In its stated form, this formula was presented by Schenck [16, Theorem 3.1]. □

The following proposition is one of our main results from [17], which gives a degree bound after which the vertex contributions stabilize.

Proposition 4.2 [17, Corollary 3.18] *Suppose $\Delta \subset \mathbb{R}^3$ is a generic closed vertex star with interior vertex γ , $r \geq 0$ is an integer, and D_γ is the integer defined in (1). Then $\dim \mathcal{J}(\gamma)_d \leq \binom{d+2}{2}$, with equality for $d > D_\gamma$.*

Remark 4.3 The ideal $\mathcal{J}(\gamma)$ in Proposition 4.2 is related, via inverse systems, to an ideal of so-called *fat points* in the projective plane which are dual to the linear forms generating $\mathcal{J}(\gamma)$. Such ideals are the subject of much interest in algebraic geometry; in particular the celebrated Segre-Harbourne-Gimigliano-Hirschowitz conjecture concerns the dimension of the graded pieces of fat point ideals for general points in the projective plane (see [30] for a survey on this connection).

The smallest degree d for which $\dim \mathcal{J}(\gamma)_d = \binom{d+2}{2}$ is called the *(Castelnuovo-Mumford) regularity* of $\mathcal{J}(\gamma)$, written $\text{reg}(\mathcal{J}(\gamma))$. Our main observation in [17] is that $\text{reg}(\mathcal{J}(\gamma))$ can be bounded by the *Waldschmidt constant* of the ideal of the dual set of points. When Δ is a generic vertex star, we take advantage of the fact that the dual set of points is covered by relatively few lines corresponding to the interior edges of Δ . For such point sets, we use work of Cooper, Harbourne, and Teitler [12] to bound the Waldschmidt constant, obtaining Proposition 4.2.

For a tetrahedral partition Δ and vertex $\gamma \in \Delta_0$, we now relate the formulas $\text{LB}^\star(\Delta_\gamma, d, r)$ and $\text{LB}^{\star\star}(\Delta_\gamma, d, r)$ from Equations (2) and (3) (respectively) to the Euler characteristic of $\mathcal{J}[\Delta_\gamma]$.

Proposition 4.4 *Let Δ be a generic tetrahedral partition. If $\gamma \in \Delta_0^\circ$ then*

$$\begin{aligned} \text{LB}^\star(\Delta_\gamma, d, r) &= 2 \binom{d+2}{2} + \chi(\mathcal{J}[\Delta_\gamma], d) - \dim \mathcal{J}(\gamma)_d \\ &= \binom{d+2}{2} + \chi(\mathcal{J}[\Delta_\gamma], d) \text{ if } d > D_\gamma, \end{aligned}$$

where $\text{LB}^\star(\Delta_\gamma, d, r)$ is defined in Equation (2). If γ is a boundary vertex of Δ then

$$\text{LB}^\star(\Delta_\gamma, d) = \binom{d+2}{2} + \chi(\mathcal{J}[\Delta_\gamma], d) \text{ for all } d \geq 0,$$

where $\text{LB}^\star(\Delta_\gamma, d, r)$ is defined in Equation (3).

Proof If γ is an interior vertex then Δ_γ is a closed vertex star $\mathcal{J}[\Delta_\gamma]$ has the form $0 \rightarrow \bigoplus_{\sigma \in \Delta_2^\circ} \mathcal{J}(\sigma) \rightarrow \bigoplus_{\tau \in \Delta_1^\circ} \mathcal{J}(\tau) \rightarrow \mathcal{J}(\gamma) \rightarrow 0$. Taking the graded Euler characteristic, the first equation now follows from the fact that $\dim \mathcal{J}(\sigma)_d = \binom{d+1-r}{2}$, Proposition 4.1, and Proposition 4.2. If γ is a boundary vertex then Δ_γ is an open vertex star and $\mathcal{J}[\Delta_\gamma]$ has the form $0 \rightarrow \bigoplus_{\sigma \in \Delta_2^\circ} \mathcal{J}(\sigma) \rightarrow \bigoplus_{\tau \in \Delta_1^\circ} \mathcal{J}(\tau) \rightarrow 0$. Taking the graded Euler characteristic, the first equation now follows from the fact that $\dim \mathcal{J}(\sigma)_d = \binom{d+1-r}{2}$ and Proposition 4.1. \square

Proposition 4.5 *Suppose $\Delta \subset \mathbb{R}^3$ is a tetrahedral partition, $r \geq 0$ is an integer, $\sigma \in \Delta_2$, $\tau \in \Delta_1$, and $\gamma \in \Delta_0^\circ$. Then*

$$\dim \widehat{\mathcal{S}}_d = \binom{d+3}{3}, \quad \dim \mathcal{J}(\widehat{\sigma})_d = \binom{d+2-r}{3}, \tag{12}$$

$$\begin{aligned} \dim \mathcal{J}(\widehat{\tau})_d &\leq t_\tau \binom{d+2-r}{3} - a_\tau \binom{d+2-q_\tau}{3} \\ &\quad - b_\tau \binom{d+3-q_\tau}{3}, \end{aligned} \tag{13}$$

$$\dim \mathcal{J}(\widehat{\gamma})_d = \binom{d+3}{3} + \sum_{i=0}^{D_\gamma} \left(\dim \mathcal{J}(\gamma)_i - \binom{d+2}{2} \right), \tag{14}$$

$\dim H_2(\mathcal{R}/\mathcal{J}[\widehat{\Delta}])_d = C$ for some positive integer C and $d \gg 0$, and $\dim H_1(\mathcal{R}/\mathcal{J}[\widehat{\Delta}])_d = 0$ for $d \gg 0$. If Δ is generic then (13) is an equality.

Proof Equations (12) are straightforward to derive. Equations (13) and (14) follow from Propositions 4.1 and 4.2, respectively, using Equation (11). It follows from [29, Lemma 3.1] that $H_1(\mathcal{R}/\mathcal{J}[\widehat{\Delta}])$ vanishes in large degree and $\dim H_2(\mathcal{R}/\mathcal{J}[\widehat{\Delta}])_d = C$ for a positive integer C . \square

We now provide a lower bound on the integer C satisfying $\dim H_2(\mathcal{R}/\mathcal{J}[\widehat{\Delta}])_d = C$ for $d \gg 0$ (see Proposition 4.5). The key is to describe the effect of the coning construction $\Delta \rightarrow \widehat{\Delta}$ on the homology module $H_2(\mathcal{R}/\mathcal{J}[\Delta])$ in large degree.

Proposition 4.6 *Let $\Delta \subset \mathbb{R}^3$ be a tetrahedral partition. Then, for $d \gg 0$,*

$$\begin{aligned} \dim H_2(\mathcal{R}/\mathcal{J}[\widehat{\Delta}])_d &= \sum_{\gamma \in \Delta_0} \sum_{i \geq 0} (\dim \mathcal{H}_i^r(\Delta_\gamma) - \chi(\mathcal{R}/\mathcal{J}[\Delta_\gamma], i)) \\ &= \sum_{\gamma \in \Delta_0} \sum_{i=0}^{3r+1} \left(\dim \mathcal{H}_i^r(\Delta_\gamma) - \binom{i+2}{2} - \chi(\mathcal{J}[\Delta_\gamma], i) \right) \\ &\geq \sum_{\gamma \in \Delta_0} \sum_{i=0}^{3r+1} [-\chi(\mathcal{J}[\Delta_\gamma], i)]_+ \end{aligned}$$

Proof The first equality is [15, Corollary 9.2]. For the second equality,

$$\dim H_1(\mathcal{J}[\Delta_\gamma])_i = \dim \mathcal{H}_i^r(\Delta) - \binom{i+2}{2} - \chi(\mathcal{J}[\Delta_\gamma], i)$$

is an immediate consequence of Equation (10). It follows from the main result of [2] (see also [16]) that $\dim \mathcal{H}_i^r(\Delta_\gamma) = \binom{i+2}{2} + \chi(\mathcal{J}[\Delta_\gamma], i)$ for $i \geq 3r + 2$. In other words, $H_1(\mathcal{J}[\Delta_\gamma])_i = 0$ for $i \geq 3r + 2$. The final inequality follows from the fact that $\mathcal{H}_i^r(\Delta_\gamma)$ always contains the space of global homogeneous polynomials of degree i , which has dimension $\binom{i+2}{2}$. □

We prove in [17] the following slight modification of a result of Whiteley [36].

Theorem 4.7 [17, Theorem 1.3] *If Δ is a generic closed star with interior vertex γ , then $\dim \mathcal{H}_d^r(\Delta) = \binom{d+2}{2}$ for $d \leq D_\gamma$.*

Corollary 4.8 *If Δ is a generic closed star with interior vertex γ , then $\dim H_1(\mathcal{J}[\Delta])_d = -\chi(\mathcal{J}[\Delta], d)$ for $d \leq D_\gamma$.*

Proof Immediate from Equation (10) and Theorem 4.7. □

Proof of Theorem 2.6 Since $H_0(\mathcal{J}[\widehat{\Delta}])_d = 0$ for $d \gg 0$ by Proposition 4.5, then (9) implies that for $d \gg 0$,

$$\dim S_d^r(\Delta) = (f_3 - f_2^\circ + f_1^\circ - f_0^\circ) \binom{d+3}{3} + \chi(\mathcal{J}[\widehat{\Delta}], d) + \dim H_2(\mathcal{R}/\mathcal{J}[\widehat{\Delta}])_d. \tag{15}$$

By Proposition 4.5, there is a constant C so that $\dim H_2(\mathcal{R}/\mathcal{J}[\widehat{\Delta}])_d = C$ when $d \gg 0$. Hence it suffices to prove that $\text{LB}(\Delta, d, r) \leq (f_3 - f_2^\circ + f_1^\circ - f_0^\circ) \binom{d+3}{3} +$

$\chi(\mathcal{J}[\widehat{\Delta}], d) + C \cdot \text{Put}$

$$\chi'(d) := \left(f_2^\circ - \sum_{\tau \in \Delta_1^\circ} t_\tau \right) \binom{d+2-r}{3} + \sum_{\tau \in \Delta_1^\circ} \left(a_\tau \binom{d+2-q_\tau}{3} + b_\tau \binom{d+3-q_\tau}{3} \right),$$

so $\chi(\mathcal{J}[\widehat{\Delta}], d) = \chi'(d) + \sum_{\gamma \in \Delta_0^\circ} \dim \mathcal{J}(\widehat{\gamma})_d$ by Equation (4) and Proposition 4.5. Another application of Proposition 4.5 gives

$$\chi(\mathcal{J}[\widehat{\Delta}], d) = \chi'(d) + f_0^\circ \binom{d+3}{3} + \sum_{\gamma \in \Delta_0^\circ} \sum_{i=0}^{D_\gamma} \left(\dim \mathcal{J}(\gamma)_i - \binom{i+2}{2} \right). \tag{16}$$

Now, by Proposition 4.6, $\dim H_1(\mathcal{J}[\widehat{\Delta}])_d \geq \sum_{\gamma \in \Delta_0} \sum_{i=0}^{3r+1} [-\chi(\mathcal{J}[\Delta_\gamma], i)]_+$ for $d \gg 0$.

Corollary 4.8 allows us to remove the $+$ from the summation for interior vertices in the range $0 \leq i \leq D_\gamma$:

$$\begin{aligned} \dim H_1(\mathcal{J}[\widehat{\Delta}])_d &\geq \sum_{\gamma \in \Delta_0^\circ} \left(\sum_{i=0}^{D_\gamma} [-\chi(\mathcal{J}[\Delta_\gamma], i)] + \sum_{i=D_\gamma+1}^{3r+1} [-\chi(\mathcal{J}[\Delta_\gamma], i)]_+ \right) \\ &\quad + \sum_{\gamma \in \Delta_0 \setminus \Delta_0^\circ} \sum_{i=0}^{3r+1} [-\chi(\mathcal{J}[\Delta_\gamma], i)]_+, \end{aligned} \tag{17}$$

for $d \gg 0$. Combining (16) and (17) with (15) yields

$$\begin{aligned} \dim S_d^r(\Delta) &\geq (f_3 - f_2^\circ + f_1^\circ) \binom{d+3}{3} + \chi'(d) \\ &\quad + \sum_{\gamma \in \Delta_0^\circ} \left(\sum_{i=0}^{D_\gamma} \left[\dim \mathcal{J}(\gamma)_i - \binom{i+2}{2} - \chi(\mathcal{J}[\Delta_\gamma], i) \right] \right. \\ &\quad \left. + \sum_{i=D_\gamma+1}^{3r+1} [-\chi(\mathcal{J}[\Delta_\gamma], i)]_+ \right) \\ &\quad + \sum_{\gamma \in \Delta_0 \setminus \Delta_0^\circ} \sum_{i=0}^{3r+1} [-\chi(\mathcal{J}[\Delta_\gamma], i)]_+ \end{aligned}$$

for $d \gg 0$. By Proposition 4.4, if $\gamma \in \Delta_0^\circ$,

$$\begin{aligned} \binom{i+2}{2} - \text{LB}^\star(\Delta_\gamma, i, r) &= \dim \mathcal{J}(\gamma)_i - \binom{i+2}{2} - \chi(\mathcal{J}[\Delta_\gamma], i), \text{ and} \\ \binom{i+2}{2} - \text{LB}^\star(\Delta_\gamma, i, r) &= -\chi(\mathcal{J}[\Delta_\gamma], i) \text{ for } i > D_\gamma. \end{aligned}$$

Also by Proposition 4.4, if $\gamma \in \Delta_0 \setminus \Delta_0^\circ$ then $\binom{i+2}{2} - \text{LB}^\star(\Delta_\gamma, i, r) = -\chi(\mathcal{J}[\Delta_\gamma], i)$. Thus,

$$\begin{aligned} \dim S'_d(\Delta) &\geq (f_3 - f_2^\circ + f_1^\circ) \binom{d+3}{3} + \chi'(d) \\ &\quad + \sum_{\gamma \in \Delta_0^\circ} \left(\sum_{i=0}^{D_\gamma} \left[\binom{i+2}{2} - \text{LB}^\star(\Delta_\gamma, i, r) \right] \right. \\ &\quad \left. + \sum_{i=D_\gamma+1}^{3r+1} \left[\binom{i+2}{2} - \text{LB}^\star(\Delta_\gamma, i, r) \right]_+ \right) \\ &\quad + \sum_{\gamma \in \Delta_0 \setminus \Delta_0^\circ} \sum_{i=0}^{3r+1} \left[\binom{i+2}{2} - \text{LB}^\star(\Delta_\gamma, i, r) \right]_+ \\ &= (f_3 - f_2^\circ + f_1^\circ) \binom{d+3}{3} + \chi'(d) - f_0^\circ \binom{r+3}{3} \\ &\quad + \sum_{\gamma \in \Delta_0^\circ} \left(\sum_{i=r+1}^{D_\gamma} \left[\binom{i+2}{2} - \text{LB}^\star(\Delta_\gamma, i, r) \right] \right. \\ &\quad \left. + \sum_{i=D_\gamma+1}^{3r+1} \left[\binom{i+2}{2} - \text{LB}^\star(\Delta_\gamma, i, r) \right]_+ \right) \\ &\quad + \sum_{\gamma \in \Delta_0 \setminus \Delta_0^\circ} \sum_{i=r+1}^{3r+1} \left[\binom{i+2}{2} - \text{LB}^\star(\Delta_\gamma, i, r) \right]_+ \\ &= (f_3 - f_2^\circ + f_1^\circ) \binom{d+3}{3} + \chi'(d) - f_0^\circ \binom{r+3}{3} \\ &\quad + \sum_{\gamma \in \Delta_0} N_\gamma = \text{LB}(\Delta, d, r), \end{aligned}$$

where N_γ is defined in (4) and $\text{LB}(\Delta, d, r)$ is defined in (5). □

5 Examples

In this section we compare our lower bounds with those by Alfeld and Schumaker in [5] and Murrain and Villamizar [28]. Except for the non-simplicial partition in

Table 4 Lower bounds for the Morgan-Scott with cavity partition in Example 5.2. The initial degree is bolded

r	d	$\binom{d+3}{3}$	LB [5]	LB(d)	gendim
1	2	10	10	8	10
1	3	20	20	16	20
1	4	35	46	46	46
1	5	56	112	112	112
2	4	35	35	34	35
2	5	56	56	40	56
2	6	84	84	72	84
2	7	120	120	144	144
2	8	165	242	270	270
2	9	220	436	464	464
3	6	84	84	84	84
3	7	120	120	96	120
3	8	165	165	138	165
3	9	220	220	224	224
3	10	286	286	368	368
3	11	364	428	584	584
4	8	165	165	162	165
4	9	220	220	176	220
4	10	286	286	224	286
4	11	364	364	320	364
4	12	455	455	478	478
4	13	560	560	712	712

Example 5.4, the other examples appear in [5]. It is well-known that for $d \gg 0$, $\dim S_d^r(\Delta)$ is a *polynomial* function. That is, there is a polynomial in d with rational coefficients, which we denote by $P_d^r(\Delta)$, so that $\dim S_d^r(\Delta) = P_d^r(\Delta)$ for $d \gg 0$. (In commutative algebra this is called the *Hilbert polynomial* of $S^r(\widehat{\Delta})$ —see Remark 6.1.) We can compute both the exact dimension $\dim S_d^r(\Delta)$ and the polynomial $P_d^r(\Delta)$ in Macaulay2 [20] using the Algebraic Splines package. We give the computations of $P_d^r(\Delta)$ in Sects. 5.1, 5.2, 5.3, and 5.4 for generic vertex positions of the examples. The exact generic dimension $\dim S_d^r(\Delta)$ for our examples is shown in the column labeled ‘gendim’ in Tables 1, 4, 5 and 6. The lower bound from Theorem 2.6 is in the column labeled LB(d), and lower bounds from the literature appear in columns labeled LB with an appropriate citation.

5.1 Three dimensional Morgan–Scott

Let Δ be the simplicial complex in Fig. 1 from Examples 1.1 and 2.1. For order of smoothness $r = 3$ and $r = 4$, the lower bounds obtained by applying Theorem 2.6 are recorded in column 6 in Table 1. For $d \gg 0$, the lower bounds can be computed

Table 5 Lower bounds for the Square-shaped torus partition in Example 5.3. The initial degree is bolded

r	d	$\binom{d+3}{3}$	LB[5]	LB(d)	gendim
1	2	10	10	8	10
1	3	20	48	48	48
1	4	35	144	144	144
1	5	56	320	320	320
2	4	35	35	24	35
2	5	56	93	96	96
2	6	84	237	240	240
2	7	120	477	480	480
2	8	165	837	840	840
2	9	220	1341	1344	1344
3	6	84	84	60	84
3	7	120	151	176	176
3	8	165	351	380	380
3	9	220	663	696	696
3	10	286	1111	1148	1148
3	11	364	1719	1760	1760
4	8	165	165	120	165
4	9	220	220	288	288
4	10	286	483	560	560
4	11	364	875	960	960
4	12	455	1419	1512	1512
4	13	560	2139	2240	2240

as in Example 2.1 and are given by

$$\begin{aligned}
 \text{LB}(\Delta, d, 3) &= \frac{5}{3}d^3 - 41d^2 + \frac{451}{2}d - 323 \quad \text{and} \quad \text{LB}(\Delta, d, 4) \\
 &= \frac{5}{2}d^3 - 55d^2 + \frac{807}{2}d - 803.
 \end{aligned}$$

These coincide with the polynomials $P_d^3(\Delta)$ and $P_d^4(\Delta)$, respectively.

5.2 Morgan–Scott with a cavity

We consider Δ as the partition obtained by removing the central tetrahedron in Fig. 1. In Table 4a we list the values of the lower bound in Theorem 2.6 applied for $r = 1, \dots, 4$ along with those presented in [5, Example 8.4]. For this partition we have $f_3 = 14$ tetrahedra, $f_2^\circ = 24$, $f_2^\circ = 12$, and $f_0^\circ = 0$. Applying (5) in Theorem 2.6 we get

$$\text{LB}(\Delta, d, 1) = \frac{7}{3}d^3 - 10d^2 + \frac{41}{3}d + 2, \quad \text{LB}(\Delta, d, 2) = \frac{7}{3}d^3 - 22d^2 + \frac{185}{3}d - 10,$$

Table 6 Bounds for the non-simplicial partition Δ in Example 5.4, Fig. 2 (right); The bolded entry in the d column indicates the initial degree of $S_d^r(\Delta)$

r	d	$\binom{d+3}{3}$	LB[28]	LB(d)	gendim
1	3	20	20	-8	20
1	4	35	35	1	35
1	5	56	56	60	60
1	6	84	160	196	196
2	8	165	165	79	165
2	9	220	220	268	268
2	10	286	352	586	586
2	11	364	826	1060	1060
2	12	455	1483	1717	1717
2	13	560	2350	2584	2584
2	14	680	3454	3688	3688
3	11	364	364	148	364
3	12	455	455	425	455
3	13	560	560	856	856
3	14	680	988	1468	1468
3	15	816	1808	2288	2288
3	16	969	2863	3343	3343

$$\begin{aligned} \text{LB}(\Delta, d, 3) &= \frac{7}{3}d^3 - 34d^2 + \frac{473}{3}d - 142, \text{ and } \text{LB}(\Delta, d, 4) \\ &= \frac{7}{3}d^3 - 46d^2 + \frac{869}{3}d - 406. \end{aligned}$$

As shown in Table 4a, for $r = 1, \dots, 4$, the bound $\text{LB}(\Delta, d, r)$ gives the exact dimension of $S_d^r(\Delta)$ beginning at the initial degree of $S^r(\Delta)$. Hence the polynomials $\text{LB}(\Delta, d, r)$ coincide with the polynomials $P_d^r(\Delta)$ for $r = 1, 2, 3, 4$.

5.3 Square-shaped torus

We consider the tetrahedral decomposition of the square-shaped torus depicted on the left in Fig. 2. This is composed of four three-dimensional ‘trapezoids,’ each of which is split into six tetrahedra along an interior diagonal. We have $f_3 = 24$, $f_2^\circ = 32$, $f_1^\circ = 8$, and $f_0^\circ = 0$. An explicit set of faces and coordinates is provided in [5, Example 8.3]. In Table 4, 5 and 6 we list the values of the lower bound of Theorem 2.6 applied for $r = 1, \dots, 4$ along with those presented in [5, Example 8.3]. We have,

$$\begin{aligned} \text{LB}(\Delta, d, 1) &= 4d^3 - 8d^2 + 4d, \quad \text{LB}(\Delta, d, 2) = 4d^3 - 24d^2 + 44d - 24, \\ \text{LB}(\Delta, d, 3) &= 4d^3 - 40d^2 + 128d - 132, \text{ and } \text{LB}(\Delta, d, 4) = \frac{7}{3}d^3 \\ &- 46d^2 + \frac{869}{3}d - 406 = 4d^3 - 56d^2 + 25d - 360. \end{aligned}$$

Again, the polynomials $\text{LB}(\Delta, d, r)$ coincide with $P_d^r(\Delta)$ for $r = 1, 2, 3, 4$.




Fig. 2 Square-shaped torus in Example 5.3 (left), and the non-simplicial polyhedral partition in Example 5.4 (right)

5.4 Non-simplicial partition

Upon examining the setup for Theorem 2.6 in Sect. 2, it appears that there is only one quantity that depends on the simplicial nature of the subdivision. That is the upper limit in the definition of N_γ , which is $3r + 1$. This upper limit depends on the fact that $\dim \mathcal{H}_d^r(\Delta_\gamma) = \text{LB}^\star(\Delta_\gamma, d, r)$ for $d \geq 3r + 2$, proven in [2]. It is likely that some modification of our lower bound applies for polytopal partitions. The most immediate modification is that one should not stop the sum in the definition of N_γ at degree $3r + 1$, but should continue until all positive contributions are accounted for. In [16] a bound is given that could be used as the upper limit of this sum, but in practice one should stop as soon as the contributions switch from positive to negative.

For reasons which we elaborate on in Remark 6.6, we will not attempt to prove that some appropriate modification of our lower bound works for polytopal subdivisions in this paper. Instead, we illustrate an application of the bound of Theorem 2.6 to a polytopal analog of the three-dimensional Morgan–Scott partition shown in Fig. 2. The subdivision consists of a cube inside of which we place its dual polytope (the octahedron). Then the partition consists of the interior octahedron along with the convex hull of pairs of dual faces. For example, each vertex of the inner octahedron is paired with a dual square face of the cube and their convex hull is a square pyramid. The number of interior vertices is $f_0^\circ = 6$, the number of interior edges is $f_1^\circ = 36$, and the number of interior two-faces is $f_2^\circ = 56$. Each interior vertex γ is connected by an edge to eight vertices i.e., $f_1^\circ(\Delta_\gamma) = 8$ and $f_2^\circ(\Delta_\gamma) = 16$ in the star Δ_γ . Thus, $D_\gamma = \lfloor \frac{3r+1}{2} \rfloor$ for all $\gamma \in \Delta_0^\circ$. There are eight vertices γ' on the boundary, for each of them we have $f_2^\circ(\Delta_{\gamma'}) = 9$, and $f_1^\circ(\Delta_{\gamma'}) = 3$ in the open stars $\Delta_{\gamma'}$. Applying (2) and (3) yields

$$\begin{aligned} \text{LB}^\star(\Delta_\gamma, d, r) &= 2 \binom{d+2}{2} + (16 - 8t_\tau) \binom{d+1-r}{2} \\ &\quad + 8a_\tau \binom{d+1-q_\tau}{2} + 8b_\tau \binom{d+2-q_\tau}{2}, \end{aligned}$$

and

$$\begin{aligned} \text{LB}^{\star}(\Delta_{\gamma'}, d, r) &= \binom{d+2}{2} + (9 - 3t_{\tau}) \binom{d+1-r}{2} \\ &\quad + 3a_{\tau} \binom{d+1-q_{\tau}}{2} + 3b_{\tau} \binom{d+2-q_{\tau}}{2}. \end{aligned}$$

By Theorem 2.6 the dimension of the spline space then $\dim \mathcal{S}_d^r(\Delta) \geq \text{LB}(d)$ for $d \gg 0$, where

$$\begin{aligned} \text{LB}(\Delta, d, r) &= 7 \binom{d+3}{3} + (56 - 36 \cdot 2) \binom{d+2-r}{3} \\ &\quad + 36 \binom{d+1}{3} - 6 \binom{r+3}{3} + 6N_{\gamma} + 8N_{\gamma'}. \end{aligned} \tag{18}$$

Every edge $\tau \in \Delta_1^{\circ}$ is in four two-dimensional faces i.e., $n_{\tau} = 4$. This leads to three values of t_{τ} : if $r = 0$ then $t_{\tau} = 2$; if $r = 1$ then $t_{\tau} = 2$; if $r \geq 2$ then $t_{\tau} = 4$.

Case 1. Let $r = 0$, then $t_{\tau} = 2$, $q_{\tau} = 2$, $a_{\tau} = 0$, and $b_{\tau} = 1$ for all $\tau \in \Delta_1^{\circ}$, and $D_{\gamma} = 0$ for all $\gamma \in \Delta_0^{\circ}$. It follows,

$$\begin{aligned} \text{LB}^{\star}(\Delta_{\gamma}, d, 0) &= 2 \binom{d+2}{2} + 8 \binom{d}{2}, \text{ and} \\ \text{LB}^{\star}(\Delta_{\gamma'}, d, 0) &= \binom{d+2}{2} + (9 - 3 \cdot 2) \binom{d+1}{2} + 3 \binom{d}{2}. \end{aligned}$$

From (4), we have $N_{\gamma} = N_{\gamma'} = 0$. Therefore,

$$\begin{aligned} \text{LB}(\Delta, d, 0) &= 7 \binom{d+3}{3} + (56 - 36 \cdot 2) \binom{d+2}{3} + 36 \binom{d+1}{3} - 6 \\ &= \frac{9}{2}d^3 - d^2 + \frac{3}{2}d + 1. \end{aligned} \tag{19}$$

Case 2. If $r = 1$, then $t_{\tau} = 3$, $q_{\tau} = 3$, $a_{\tau} = 0$, and $b_{\tau} = 2$ for all $\tau \in \Delta_1^{\circ}$, and $D_{\gamma} = 2$ for all $\gamma \in \Delta_0^{\circ}$. It follows,

$$\begin{aligned} \text{LB}^{\star}(\Delta_{\gamma}, d, 1) &= 2 \binom{d+2}{2} + (16 - 8 \cdot 3) \binom{d}{2} + 8 \cdot 2 \binom{d-1}{2}, \text{ and} \\ \text{LB}^{\star}(\Delta_{\gamma'}, d, 1) &= \binom{d+2}{2} + 3 \cdot 2 \binom{d-1}{2}. \end{aligned}$$

From (4) we have $N_\gamma = 2$ and $N_{\gamma'} = 0$. Therefore,

$$\begin{aligned} \text{LB}(\Delta, d, 1) &= 7\binom{d+3}{3} + (56 - 36 \cdot 3)\binom{d+1}{3} + 36 \cdot 2\binom{d}{3} - 6 \cdot 4 + 6 \cdot 2 \\ &= \frac{9}{2}d^3 - 29d^2 + \frac{91}{2}d - 5. \end{aligned} \tag{20}$$

Case 3. For every $r \geq 2$, we have $t_\tau = 4$. We write the explicit formula for $r = 2$, the other cases follow similarly. We have $q_\tau = 4$, $a_\tau = 0$ and $b_\tau = 3$ for all $\tau \in \Delta_1^\circ$, and $D_\gamma = 3$. Then,

$$\begin{aligned} \text{LB}^\star(\Delta_\gamma, d, 2) &= 2\binom{d+2}{2} + (16 - 8 \cdot 4)\binom{d-1}{2} + 8 \cdot 3\binom{d-2}{2}, \quad \text{and} \\ \text{LB}^\star(\Delta_{\gamma'}, d, 2) &= \binom{d+2}{2} + (9 - 3 \cdot 4)\binom{d-1}{2} + 3 \cdot 3\binom{d-2}{2}. \end{aligned}$$

From (4) we have $N_\gamma = 18$ and $N_{\gamma'} = 3$. Therefore,

$$\begin{aligned} \text{LB}(\Delta, d, 2) &= 7\binom{d+3}{3} + (56 - 36 \cdot 4)\binom{d}{3} \\ &\quad + 36 \cdot 3\binom{d-1}{3} - 6 \cdot 10 + 6 \cdot 18 + 8 \cdot 3 \\ &= \frac{9}{2}d^3 - 57d^2 + \frac{363}{2}d - 29. \end{aligned} \tag{21}$$

The bounds (19), (20), and (21) are the polynomials $P_d^1(\Delta)$, $P_d^2(\Delta)$, and $P_d^3(\Delta)$, respectively. In Table 6 we record the values for $\text{LB}(\Delta, d, r)$ along with the lower bound obtained in [28].

The final column in Table 6 bears closer examination. What do we mean by the ‘generic’ dimension in this example? In a simplicial complex, it is clear that small changes in vertex coordinates do not change the overall structure of the simplicial subdivision. However, if we modify the coordinates of a vertex in a non-simplicial face of a polytope, it is most likely that we have taken it out of the plane determined by the other vertices of the face, and so we no longer have the same polytope. So we must be careful about what we mean. In making the final column in Table 6 we have in fact cheated somewhat, as follows. Notice that in Example 5.4, the polytopal subdivision has only seven 3-polytopes which are not tetrahedra (the central octahedron and the six pyramids with square bases, each of which share one vertex with the central octahedron). Furthermore, every single 2-polytope is a triangle except for the squares which form the boundary of the outer cube. This allows a great deal of freedom for moving the vertices without destroying the polytopes in the subdivision. In fact, we can perturb the vertices of the central octahedron without destroying it, since all its 2-faces are triangles. We cannot perturb the eight boundary vertices as we wish without destroying the square faces of the cube. However, notice that if we perturb the coordinates of the five vertices of one of the square pyramids and take the convex

hull of the resulting five points, we obtain a bipyramid over a triangle and the original square base of the pyramid splits into two triangles which are no longer coplanar. Since this only changes the boundary of the domain, an arbitrary perturbation of the vertices of the outer cube will result in a viable polytopal complex, with twice as many two-dimensional boundary faces as the original polytopal complex. In the column labeled *gendim* in Table 6, we have recorded the dimension of splines on this modified polytopal complex obtained via a random perturbation of the vertex coordinates from the symmetric polytopal complex pictured on the right in Figure 2.

6 Concluding remarks

Remark 6.1 The dimension $\dim S_d^r(\Delta)$ of splines on Δ is a polynomial in d when $d \gg 0$; this polynomial is known as the *Hilbert polynomial* of $S^r(\widehat{\Delta})$ in algebraic geometry. Theorem 2.6 gives a lower bound on the Hilbert polynomial of $S^r(\widehat{\Delta})$. For some value of d , $\dim S_d^r(\Delta)$ will begin to agree with the Hilbert polynomial. In algebraic geometry there is an integer which bounds when $\dim S_d^r(\Delta)$ becomes polynomial, known as the *Castelnuovo-Mumford regularity* of $S^r(\Delta)$. It would be interesting to bound the regularity of $S^r(\widehat{\Delta})$ for tetrahedral partitions, perhaps by extending methods from [16].

Remark 6.2 We suspect that our formula in Theorem 2.6 is a lower bound on $\dim S_d^r(\Delta)$ for $d \geq 8r + 1$ by the following reasoning. In [6, Theorem 24], Alfeld, Schumaker, and Sirvent prove that $\dim S_d^r(\Delta) = \sum_{\beta \in \Delta} |\mathcal{D}(\beta)|$ for $d \geq 8r + 1$, where the sum runs across all simplices $\beta \in \Delta$ and $\mathcal{D}(\beta)$ is a *minimal determining set* for the simplex β . Counting the size of the sets $|\mathcal{D}(\beta)|$ gives rise to expressions using binomial coefficients using the same Convention 2.2. For $r = 1$ these are counted explicitly in [7], while counts for more general r (with supersmoothness) may be found in [8]. We expect that for a fixed r and $d \geq 8r + 1$, $|\mathcal{D}(\beta)|$ is a polynomial of degree $\dim \beta$ for all $\beta \in \Delta$. If so, then $\sum_{\beta \in \Delta} |\mathcal{D}(\beta)|$ is a polynomial for $d \geq 8r + 1$, and this is the Hilbert polynomial of $S^r(\widehat{\Delta})$. Since the formula in Theorem 2.6 is a lower bound on the Hilbert polynomial of $S^r(\widehat{\Delta})$ (see Remark 6.1), it would follow that it is a lower bound on $\dim S_d^r(\Delta)$ for $d \geq 8r + 1$. It would also be interesting to know if [6] has implications for the regularity of $S^r(\widehat{\Delta})$ (discussed in Remark 6.1).

Remark 6.3 Building on Remarks 6.1 and 6.2, we have observed in all the examples of Sects. 2.1 and 5 that $\text{LB}(\Delta, d, r) = \dim S_d^r(\Delta)$ (when Δ is generic) for d at least the *initial degree* of $S_d^r(\Delta)$; that is, the bound begins to give the exact dimension of the spline space as soon as there are non-trivial splines. To prove this one would have to know (1) that $\text{LB}(\Delta, d, r)$ agrees with $\dim S_d^r(\Delta)$ for $d \gg 0$ and (2) that the regularity of $S^r(\widehat{\Delta})$ (see Remark 6.1) is very close to the initial degree of $S^r(\Delta)$. We discuss (1) in Remark 6.4. We expect (2) to be quite difficult; a similar statement is not even known for generic triangulations, although we expect it to be true as we indicate in Remark 6.4.

Remark 6.4 In all of the examples in Sects. 2.1 and 5, if $d \gg 0$ and Δ is generic we have $\text{LB}(\Delta, d, r) = \dim S_d^r(\Delta)$; in other words $\text{LB}(\Delta, d, r)$ is the Hilbert polynomial

of $\mathcal{S}^r(\widehat{\Delta})$ when Δ is generic. This is not always the case, although it is only possible for $\text{LB}(\Delta, d, r)$ to differ from $\dim \mathcal{S}_d^r(\Delta)$ by a constant in large degree. In fact, the only term in which we can have error is the approximation provided by Proposition 4.6 to the constant C which is equal to $\dim H_2(\mathcal{R}/\mathcal{J}[\widehat{\Delta}])$ for $d \gg 0$. If γ is a boundary vertex, we see from Proposition 4.6 that its contribution to C is $\sum_{i=0}^{3r+1} \dim \mathcal{H}_i^r(\Delta_\gamma) - \text{LB}^{\star}(\Delta, i, r)$. If $\dim \mathcal{H}_i^r(\Delta_\gamma) = \max \left\{ \binom{i+2}{2}, \text{LB}^{\star}(\Delta, i, r) \right\}$ for $0 \leq i \leq 3r + 1$, then this contribution coincides exactly with

$$\sum_{i=0}^{3r+1} \dim \left[\binom{i+2}{2} - \text{LB}^{\star}(\Delta, i, r) \right]_+ = \sum_{i=0}^{3r+1} \dim [-\chi(\mathcal{J}[\Delta], i)]_+$$

and we capture the entire contribution of the boundary vertex γ to C .

If γ is an interior vertex, the proof of Theorem 4.7 in Sect. 4 shows that its contributions to C in degree $d \leq D_\gamma$ can be accounted for; in particular the term $\dim \mathcal{J}(\gamma)_d$ for $d \leq D_\gamma$ appears both in C and in the Euler characteristic of \mathcal{J} with opposite signs, and so it cancels. By Propositions 4.6 and 4.4, the contribution of γ to C in degrees $d > D_\gamma$ is $\sum_{i=D_\gamma+1}^{3r+1} \dim \mathcal{H}_i^r(\Delta_\gamma) - \text{LB}^{\star}(\Delta, i, r)$. If $\dim \mathcal{H}_i^r(\Delta_\gamma) = \max \left\{ \binom{i+2}{2}, \text{LB}^{\star}(\Delta, i, r) \right\}$ for $i > D_\gamma$, then we again capture all of the contribution of the interior vertex γ to C .

This leads us to Questions 7.1 and 7.2 in [17], namely, is it typically true that

$$\dim \mathcal{H}_d^r(\Delta) = \max \left\{ \binom{d+2}{2}, \text{LB}^{\star}(\Delta, d, r) \right\} \tag{22}$$

when Δ is a generic open vertex star, and that for $d > D_\gamma$ and Δ a generic closed vertex star

$$\dim \mathcal{H}_d^r(\Delta) = \max \left\{ \binom{d+2}{2}, \text{LB}^{\star}(\Delta, d, r) \right\}? \tag{23}$$

(Theorem 4.7 shows that $\dim \mathcal{H}_d^r(\Delta) = \binom{d+2}{2}$ when $d \leq D_\gamma$ and Δ is a generic closed vertex star.) There are configurations for open vertex stars, discussed in [17], where it is *not* true that $\dim \mathcal{H}_d^r(\Delta) = \max \left\{ \binom{d+2}{2}, \text{LB}^{\star}(\Delta, d, r) \right\}$ even for generic vertex positions. If such a configuration is present as the star of a boundary vertex inside of a larger tetrahedral partition, then our lower bound will *not* give the exact dimension in large degree. We do raise the possibility in Question 7.2 of [17] that there could be finitely many *sub-configurations* which serve as obstructions to the correctness of Equation (22) when Δ is generic. We are not aware of any configurations where Equation (23) fails for generic vertex positions when $d > D_\gamma$.

Remark 6.5 The formula we give in Theorem 2.6 is a lower bound for $\dim \mathcal{S}_d^r(\Delta)$ for $d \gg 0$ when Δ is *generic*. That is, it only depends on purely combinatorial

information of Δ such as how many triangular faces are incident upon a given edge, and not on geometric information such as whether the linear span of these triangular faces coincide. It is well-known that such coincident linear spans cause a jump in the dimension of $S_d^r(\Delta)$. As we indicate in [17, Example 6.2], our techniques can sometimes be adjusted to improve the lower bound $\text{LB}(\Delta, d, r)$ for these types of special positions. We leave this as a future research direction. Other special positions, such as the special positions of the Morgan–Scott configuration, may depend on global geometry which is invisible to our methods.

Remark 6.6 As we discuss and illustrate in Sect. 5.4, it is possible that our bound in Theorem 2.6 holds for polytopal subdivisions as long as some appropriate modifications are made. We comment on a few subtleties that arise in the polytopal case. First, our arguments in this paper are simplified by the existence of a *generic* dimension for $S_d^r(\Delta)$ when Δ is simplicial—see Sect. 3.4. The lower bound we present in Theorem 2.6 is a lower bound for this generic dimension. On the other hand, it is not entirely clear what a *generic* polytopal subdivision is, as we pointed out in the final paragraph of Sect. 5.4. Thus to verify Theorem 2.6 for polytopal subdivisions we would need to take care to define what *generic* dimension is, or find an argument that avoids this notion.

A related complication that arises in the non-simplicial case is the presence of ‘unexpected’ associated primes for $H_2(\mathcal{R}/\mathcal{J}[\widehat{\Delta}])$, where $\Delta \subset \mathbb{R}^2$ is a polytopal complex. By ‘unexpected,’ we mean that the associated prime is not the ideal of a face of Δ . McDonald and Schenck first observed these in [26] for planar polytopal splines, where they prove that such associated primes contribute a vertex-like term to the dimension of the spline space. The associated primes of the homologies of $\mathcal{R}/\mathcal{J}[\widehat{\Delta}]$ were further analyzed in [15] for Δ a higher dimensional polytopal subdivision. The non-simplicial example we analyze in Example 5.4 also appears in [15, Example 6.4], where it is shown that there are ‘unexpected’ associated primes for certain vertex positions. At these vertex positions, Proposition 4.5 falls apart because $\dim H_2(\mathcal{R}/\mathcal{J}[\widehat{\Delta}])_d$ is eventually linear in d and $\dim H_1(\mathcal{R}/\mathcal{J}[\widehat{\Delta}])_d$ is eventually constant. It is not too difficult to see that these unexpected associated primes do not show up for sufficiently general vertex positions in Example 5.4. However, in order to extend our Theorem 2.6 to polytopal subdivisions, we would need a guarantee that such ‘unexpected’ associated primes do not appear for a ‘generic’ polytopal complex (whatever this may mean!). A cautionary example is provided by Barnette’s first diagram [39, Example 5.11]; here a coincidence of four lines (at a point which is not a vertex of the polytopal complex) is imposed simply by the combinatorial structure—that is, the partially ordered set of inclusions among the faces. Thus it may not be possible to exclude ‘unexpected’ associated primes even for ‘generic’ polytopal complexes.

For these reasons, we do not attempt in this paper to extend Theorem 2.6 to polytopal subdivisions, but leave this as a future avenue of research.

Funding N. Villamizar was supported by the UK Engineering and Physical Sciences Research Council (EPSRC) New Investigator Award EP/V012835/1.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included

in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

1. Alfeld, P.: Upper and lower bounds on the dimension of multivariate spline spaces. *SIAM J. Numer. Anal.* **33**(2), 571–588 (1996)
2. Alfeld, P., Neamtu, M., Schumaker, L.: Dimension and local bases of homogeneous spline spaces. *SIAM J. Math. Anal.* **27**(5), 1482–1501 (1996)
3. Alfeld, P., Schumaker, L.: The dimension of bivariate spline spaces of smoothness r for degree $d \geq 4r + 1$. *Constr. Approx.* **3**(2), 189–197 (1987)
4. Alfeld, P., Schumaker, L.: On the dimension of bivariate spline spaces of smoothness r and degree $d = 3r + 1$. *Numer. Math.* **57**(6–7), 651–661 (1990)
5. Alfeld, P., Schumaker, L.: Bounds on the dimensions of trivariate spline spaces. *Adv. Comput. Math.* **29**(4), 315–335 (2008)
6. Alfeld, P., Schumaker, L., Sirvent, M.: On dimension and existence of local bases for multivariate spline spaces. *J. Approx. Theory* **70**(2), 243–264 (1992)
7. Alfeld, P., Schumaker, L., Whiteley, W.: The generic dimension of the space of C^1 splines of degree $d \geq 8$ on tetrahedral decompositions. *SIAM J. Numer. Anal.* **30**(3), 889–920 (1993)
8. Alfeld, P., Sirvent, M.: A recursion formula for the dimension of super spline spaces of smoothness r and degree $d > r2^k$. In: *Multivariate approximation theory, IV* (Oberwolfach, 1989), *Internat. Ser. Numer. Math.*, vol. 90, pp. 1–8. Birkhäuser, Basel (1989)
9. Billera, L.: Homology of smooth splines: generic triangulations and a conjecture of Strang. *Trans. Am. Math. Soc.* **310**(1), 325–340 (1988)
10. Billera, L.: The algebra of continuous piecewise polynomials. *Adv. Math.* **76**(2), 170–183 (1989)
11. Billera, L., Rose, L.: A dimension series for multivariate splines. *Discret. Comput. Geom.* **6**(2), 107–128 (1991)
12. Cooper, S., Harbourne, B., Teitler, Z.: Combinatorial bounds on Hilbert functions of fat points in projective space. *J. Pure Appl. Algebra* **215**(9), 2165–2179 (2011)
13. Cottrell, J., Hughes, T., Bazilevs, Y.: *Isogeometric Analysis: Toward Integration of CAD and FEA*. John Wiley & sons, Ltd (2009)
14. Courant, R.: Variational methods for the solution of problems of equilibrium and vibrations. *Bull. Am. Math. Soc.* **49**, 1–23 (1943)
15. DiPasquale, M.: Associated primes of spline complexes. *J. Symb. Comput.* **76**, 158–199 (2016)
16. DiPasquale, M.: Dimension of mixed splines on polytopal cells. *Math. Comp.* **87**(310), 905–939 (2018)
17. DiPasquale, Michael; Villamizar, Nelly: A lower bound for splines on tetrahedral vertex stars. *SIAM J. Appl. Algebra Geom.* **5**(2), 250–277 (2021)
18. Engvall, L., Evans, J.: Isogeometric triangular Bernstein-Bézier discretizations: automatic mesh generation and geometrically exact finite element analysis. *Comput. Methods Appl. Mech. Eng.* **304**, 378–407 (2016)
19. Engvall, L., Evans, J.: Isogeometric unstructured tetrahedral and mixed-element Bernstein-Bézier discretizations. *Comput. Methods Appl. Mech. Eng.* **319**, 83–123 (2017)
20. Grayson, D., Stillman, M.: *Macaulay2, a software system for research in algebraic geometry*. Available at <http://www.math.uiuc.edu/Macaulay2/>
21. Hatcher, A.: *Algebraic Topol.* Cambridge University Press, Cambridge (2002)
22. Hong, D.: Spaces of bivariate spline functions over triangulation. *Approx. Theory Appl.* **7**(1), 56–75 (1991)
23. Ibrahim, A., Schumaker, L.: Super spline spaces of smoothness r and degree $d \geq 3r + 2$. *Constr. Approx.* **7**(3), 401–423 (1991)
24. Lai, M., Schumaker, L.: Spline functions on triangulations. *Encyclopedia of Mathematics and its Applications*, vol. 110. Cambridge University Press, Cambridge (2007)
25. Lau, W.: A lower bound for the dimension of trivariate spline spaces. *Constr. Approx.* **23**(1), 23–31 (2006)
26. McDonald, T., Schenck, H.: Piecewise polynomials on polyhedral complexes. *Adv. in Appl. Math.* **42**(1), 82–93 (2009)

27. Morgan, J., Scott, R.: A nodal basis for C^1 piecewise polynomials of degree $n \geq 5$. *Math. Comput.* **29**, 736–740 (1975)
28. Mourrain, B., Villamizar, N.: Bounds on the dimension of trivariate spline spaces: a homological approach. *Math. Comput. Sci.* **8**(2), 157–174 (2014)
29. Schenck, H.: A spectral sequence for splines. *Adv. Appl. Math.* **19**(2), 183–199 (1997)
30. Schenck, H.: Algebraic methods in approximation theory. *Comput. Aided Geom. Des.* **45**, 14–31 (2016)
31. Schenck, H., Stillman, M.: A family of ideals of minimal regularity and the Hilbert series of $C^r(\hat{\Delta})$. *Adv. Appl. Math.* **19**(2), 169–182 (1997)
32. Schenck, H., Stillman, M.: Local cohomology of bivariate splines. *J. Pure Appl. Algebra* **117**(118), 535–548 (1997)
33. Schumaker, L.: Bounds on the dimension of spaces of multivariate piecewise polynomials. *Rocky Mt. J. Math.* **14**(1), 251–264 (1984)
34. Strang, G.: Piecewise polynomials and the finite element method. *Bull. Am. Math. Soc.* **79**, 1128–1137 (1973)
35. Ženčšek, A.: Polynomial approximation on tetrahedrons in the finite element method. *J. Approx. Theory* **7**, 334–351 (1973)
36. Whiteley, W.: The combinatorics of bivariate splines. In: *Applied geometry and discrete mathematics*, DIMACS Ser. Discrete Math. Theoret. Comput. Sci., vol. 4, pp. 587–608. Amer. Math. Soc. Providence, RI (1991)
37. Whiteley, W.: A matrix for splines. In: *Progress in approximation theory*, pp. 821–828. Academic Press, Boston, MA (1991)
38. Xia, S., Qian, X.: Isogeometric analysis with Bézier tetrahedra. *Comput. Methods Appl. Mech. Eng.* **316**, 782–816 (2017)
39. Ziegler, G.: *Lectures on polytopes*. Graduate Texts in Mathematics, vol. 152. Springer-Verlag, New York (1995)

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.