



On Weak Flexibility in Planar Graphs

Bernard Lidický¹ · Tomáš Masařík^{2,3} · Kyle Murphy^{1,4} · Shira Zerbib¹

Received: 20 September 2020 / Revised: 29 August 2022 / Accepted: 2 September 2022 /
Published online: 27 October 2022
© The Author(s) 2022

Abstract

Recently, Dvořák, Norin, and Postle introduced flexibility as an extension of list coloring on graphs (J Graph Theory 92(3):191–206, 2019, <https://doi.org/10.1002/jgt.22447>). In this new setting, each vertex v in some subset of $V(G)$ has a request for a certain color $r(v)$ in its list of colors $L(v)$. The goal is to find an L coloring satisfying many, but not necessarily all, of the requests. The main studied question is whether there exists a universal constant $\varepsilon > 0$ such that any graph G in some graph class \mathcal{C} satisfies at least ε proportion of the requests. More formally, for $k > 0$ the goal is to prove that for any graph $G \in \mathcal{C}$ on vertex set V , with any list assignment L of size k for each vertex, and for every $R \subseteq V$ and a request vector $(r(v) : v \in R, r(v) \in L(v))$, there exists an L -coloring of G satisfying at least $\varepsilon|R|$ requests. If this is true, then \mathcal{C} is called ε -flexible for lists of size k . Choi, Clemen, Ferrara, Horn, Ma, and Masařík (Discrete Appl Math 306:20–132, 2022, <https://doi.org/10.1016/j.dam.2021.09.021>) introduced the notion of *weak flexibility*, where $R = V$. We further develop this direction by introducing a tool to handle weak flexibility. We demonstrate this new tool by showing that for every positive integer b there exists $\varepsilon(b) > 0$ so that the class of planar graphs without K_4, C_5, C_6, C_7, B_b is weakly $\varepsilon(b)$ -flexible for lists of size 4 (here K_n, C_n and B_n are the complete graph, a cycle, and a book on n vertices, respectively). We also show that the class of planar graphs without K_4, C_5, C_6, C_7, B_5 is ε -flexible for lists of size 4. The results are tight as these graph classes are not even 3-colorable.

Keywords List coloring · Flexibility · Discharging

Mathematics Subject Classification 05C15

✉ Tomáš Masařík
masarik@mimuw.edu.pl

Extended author information available on the last page of the article

1 Introduction

A k -coloring of a graph G is a function $f : V(G) \rightarrow S$, where $|S| = k$. The elements of S are often called *colors*. A k -coloring of G is called *proper* if adjacent vertices are assigned different colors. Suppose that for each vertex v in G , we gave v a list $L(v)$ of available colors. A *list coloring* of a graph G is a proper coloring of G where each vertex v is assigned a color from $L(v)$. In particular, for two distinct vertices u and v , $L(u)$ and $L(v)$ might be different. A graph is *k -choosable* if every assignment L of at least k colors to each vertex guarantees an L -coloring. The *choosability* of a graph G is the minimum k such that G is k -choosable.

In many applications of list coloring, such as scheduling, some vertices may have preferences which are not directly captured by the lists themselves. For example, a professor may be willing to teach classes X,Y, or Z but prefers to teach X. Ideally, the scheduler can satisfy the specific requests of each professor, but it is often the case that they cannot. The goal is then to satisfy as many requests as possible. This idea motivates the following definitions.

A *weighted request* is a function w that assigns a nonnegative real number to each pair (v, c) where $v \in V(G)$ and $c \in L(v)$. For $\epsilon > 0$, we say that w is ϵ -satisfiable if there exists an L -coloring φ of G such that

$$\sum_{v \in V(G)} w(v, \varphi(v)) \geq \epsilon \cdot \sum_{v \in V(G), c \in L(v)} w(v, c).$$

The unweighted variant is defined as follows. A *request* for a graph G with a list assignment L is a function r with domain $\text{dom}(r) \subseteq V(G)$ such that $r(v) \in L(v)$ for all $v \in \text{dom}(r)$. In the special case that each vertex requests a color, i.e., $\text{dom}(r) = V(G)$, we call such a request *widespread*. Analogously, for $\epsilon > 0$, a request r is ϵ -satisfiable if there exists an L -coloring φ of G such that at least $\epsilon|\text{dom}(r)|$ vertices v in $\text{dom}(r)$ receive color $r(v)$. We say that a graph G with list assignment L is ϵ -flexible, weakly ϵ -flexible, or *weighted ϵ -flexible* if every request, widespread request, or weighted request, respectively, is ϵ -satisfiable. Note that weak flexibility does not make sense in the weighted setting since one can set some weights to 0 to turn off the requests for these vertices. If G is (weighted/weakly) ϵ -flexible for every list assignment with lists of length k , we say that G is (weighted/weakly) ϵ -flexible for lists of size k . Note that for k -colorable graphs, if the lists are exactly the same the problem becomes trivial as by permuting the colors we can achieve $\frac{1}{k}$ -flexibility [6].

The concept of ϵ -flexibility was introduced by Dvořák, Norin, and Postle [6]. Subsequently, it was studied for various sub-classes of planar graphs, e.g., triangle-free [5], girth six [4], or C_4 -free [10]. Graphs of bounded maximum degree were subsequently characterized in terms of flexibility [1].

A central notion in graph coloring is that of *reducible configurations*, which are local subgraphs that cannot appear in a smallest counterexample because their presence implies that the graph can be colored from a smaller subgraph by induction. Reducible configurations for flexibility are slightly more delicate as we explain in Sect. 2. Recently, Choi, Clemen, Ferrara, Horn, Ma, and Masařík [2] proposed a strengthened tool (see Lemma 1 below) for designing reducible configurations for flexibility. The authors of [2] also introduced the notion of weak flexibility defined above.

They demonstrated that the weak setting allows one to create stronger reducible configurations. We further develop this direction by strengthening the tools for handling weak flexibility; see Lemma 2 in Sect. 3. We exhibit our new tool by showing the following results for subclasses of planar graphs.

For an integer $n \geq 3$ let B_n denote the book on n vertices, i.e., the graph consisting of $n - 2$ triangles sharing an edge. Let C_n and K_n denote a cycle and a clique on n vertices, respectively. Given a set of graphs \mathcal{F} and a graph H , we say that H is \mathcal{F} -free if there is no subgraph of H isomorphic to any of the graphs in \mathcal{F} .

Theorem 1 *There exists $\epsilon > 0$ such that every planar $\{K_4, C_5, C_6, C_7, B_5\}$ -free graph is weighted ϵ -flexible for lists of size 4.*

Theorem 2 *There exists $\epsilon = \epsilon(b) > 0$ such that every planar $\{K_4, C_5, C_6, C_7, B_b\}$ -free graph is weakly ϵ -flexible for lists of size 4.*

The results in Theorems 1 and 2 are tight as in general such graphs are not even 3-colorable. This is exemplified by the construction in Fig. 1. This construction implies:

Observation 3 *For every $\ell, b \geq 5$ exists a $\{K_4, B_b\}$ -free planar graph G that does not contain any cycle C_k of length $5 \leq k \leq \ell$, such that G is a not 3-colorable.*

Furthermore, our results follow a recent line of research trying to narrow the gap between known degeneracy upper-bounds and choosability lower-bounds, in particular on subclasses of planar graphs, as is described below. We say that a graph G is d -degenerate if each induced subgraph of G contains a vertex of degree at most d . It is easy to observe that d -degenerate graphs are $(d + 1)$ -choosable. A similar statement holds for flexibility as well: in [6] it was proved that d -degenerate graphs with lists of size $d + 2$ are weighted ϵ -flexible. Therefore, as C_5 -free planar graphs are 3-degenerate [11], they are ϵ -flexible for lists of size 5. The same is true for




Fig. 1 A construction proving Observation 3 with an attempt for a 3-coloring that fails

C_6 -free planar graphs [7]. For C_3 -free graphs, Dvořák, Masařík, Musílek, and Pángrác [5] showed that they are weighted ε -flexible for lists of size 4 and that this result is tight. Surprisingly, the discharging proof in [5] is quite involved compared to the easy observation that C_3 -free planar graphs are 3-degenerate, which implies 4-choosability. An analogous result holds for $\{C_3, C_4, C_5\}$ -free graphs, where list of size 3 are sufficient for weighted ε -flexibility and the result is tight [4].

When only C_4 is forbidden, Masařík [10] proved that lists of size 5 are sufficient for weighted ε -flexibility. However, it is unknown whether the result is tight as those graphs are 4-choosable [8] (but not necessarily 3-degenerate). There were attempts to bring down the list size to 4 but so far only partial results are known in this direction: planar graphs that do not contain C_4 and C_3 at distance at most 1 [2] or $\{C_4, C_5\}$ -free planar graphs [12]. See [2, Table 1] for a comprehensive overview of known results for various subclasses of planar graphs. Our results aim to improve this narrow gap as they show that lists of size 4 are sufficient even for planar graphs in which some copies of C_3 and C_4 are allowed.

2 Methods—Informal Discussion

The purpose of this section is to informally describe some of the difficulties one faces when trying to extend a list-coloring proof to a flexibility proof. This discussion serves as the intuition behind the formal definitions in the next section.

As in previous related papers mentioned above, we use the discharging method to obtain our results. For an introduction to the discharging method see [3]. A typical discharging proof that a graph G is L -list-colorable gives a list of unavoidable *reducible configurations*, which are subgraphs of G that cannot appear in a minimal counterexample. The goal is to decompose G into subgraphs R_1, \dots, R_N such that R_i is a reducible configuration in $G[R_i \cup \dots \cup R_N]$ (this will be defined as a *resolution* later), so that any L -coloring of $G[R_{i+1} \cup \dots \cup R_N]$ can be extended to an L -coloring of $G[R_i \cup \dots \cup R_N]$. Extending the coloring in a descending order from R_N to R_1 gives an L -coloring of G .

When requests are introduced, this method becomes more difficult. To explain this, assume for simplicity that every vertex has a request. If we manage to accommodate one request from each R_i and each R_i has at most b vertices, then we would satisfy n/b requests, showing that G is ε -flexible for $\varepsilon = 1/b$, and our job would be done. However, this is not necessarily possible. Indeed, suppose $v \in R_i$ has some request $r(v)$ and let φ be an L -coloring of $G[R_{i+1} \cup \dots \cup R_N]$. Suppose further that v has one neighbor u in $R_{i+1} \cup \dots \cup R_N$. If $\varphi(u) = r(v)$, there is no way to simply extend φ and accommodate the request of v . Thus more changes to φ , such as recoloring u , would have to occur to accommodate $r(v)$. In addition to this issue, it may also be the case that $r(v)$ cannot be satisfied because R_i itself prevents it.

This means that reducible configurations for flexibility need to provide slightly more freedom in the colorings they allow. The easier problem to deal with is that $r(v)$ cannot be satisfied because R_i itself prevents it. This can be patched by adding a requirement that for any one vertex x in R_i , the coloring φ extends to R_i even if x has a list of size 1 after removing the colors of already colored neighbors of x in

$R_{i+1} \cup \dots \cup R_N$. For v , this would be used in case $L(v) = \{r(v)\}$ and $\varphi(u) \neq r(v)$. This requirement will be called (FIX) in the formal definitions.

The problem occurring when $r(v)$ cannot be satisfied because its neighbor u in $R_{i+1} \cup \dots \cup R_N$ is already colored $r(v)$ is more complicated to solve. The idea is the following. Instead of constructing just one L -coloring φ , one needs to construct L -colorings $\varphi_1, \dots, \varphi_\ell$ and in some of them, u gets colored by a color different than $r(v)$. Then $\varphi_1, \dots, \varphi_\ell$ can be extended to $\varphi'_1, \dots, \varphi'_\ell$, where $r(v)$ is satisfied in some of them. At the end, this process gives a set of L -colorings of G and at least one of them satisfies a positive fraction of the requests. Formally, this is done by creating a probability distribution on L -colorings of G .

In order to make this idea work, there must be a sufficient variety of proper colorings for each reducible configuration. In our example, if we want to color v by $r(v)$, we cannot use $r(v)$ on u . We need to address this when we are coloring u and remove $r(v)$ from its list. Further, we would need to do this for each neighbor of v in $R_{i+1} \cup \dots \cup R_N$. This is achieved in the following way. When we are L -coloring R_i , we look at all subsets $I \subseteq V(R_i)$ of vertices that could form a neighborhood of a vertex in $R_1 \cup \dots \cup R_{i-1}$, i.e., in the set of not yet colored vertices. Individually for each I , we show that any proper L -coloring φ of $R_{i+1} \cup \dots \cup R_N$ can still be extended to R_i even if we decrease the sizes of the lists of vertices in I by 1. This will be called (FORB) in the formal definitions.

To summarize, the reducible configurations for flexibility must have size bounded by a constant, any one vertex can be precolored (FIX), and for different subsets of vertices, reducing their lists sizes by 1 does not break the extendability of the coloring (FORB).

The main feature of weak flexibility is that instead of demanding in (FIX) that “any one vertex in R_i can be precolored”, it is enough to ask for “at least one vertex in R_i can be precolored”. It is then easier to satisfy this version of (FIX).

We introduce an additional trick, where we ask “at least one vertex in R_i with few external neighbors can be precolored”. This makes satisfying (FIX) more difficult since the reducible configurations need a vertex of small degree but it helps a lot with checking (FORB).

3 Methods—Definitions and Lemmas

We use some of the notation and tools introduced in [2, 4–6]. In particular, our definitions are quite similar to those used in [2].

Let 1_I denote the characteristic function of I , i.e., $1_I(v) = 1$ if $v \in I$ and $1_I(v) = 0$ otherwise. Let G be a graph. Given a function $f : V(G) \rightarrow \mathbb{Z}$ and a vertex $v \in V(G)$, let $f \downarrow v$ denote the function satisfying $(f \downarrow v)(w) = f(w)$ for $w \neq v$ and $(f \downarrow v)(v) = 1$. We will use $f \downarrow v$ to indicate that the list size at vertex v has been reduced to 1. In other words, $f \downarrow v$ means that v has been “precolored”. A list assignment L is an f -assignment if $|L(v)| \geq f(v)$ for all $v \in V(G)$. We will let \deg_G be the function from $V(G)$ to \mathbb{Z} which maps each vertex to its degree. If $X \subset V(G)$, then we let \deg_X equal $\deg_{G[X]}$, where $G[X]$ is the induced subgraph of G consisting of the vertices in X .

Given a set of graphs \mathcal{F} and a graph H , a set $I \subseteq V(H)$ is \mathcal{F} -free if the graph H together with one additional vertex u adjacent to all of the vertices in I does not contain any subgraph isomorphic to a graph in \mathcal{F} .

Throughout the following definitions, let H be an induced subgraph of a graph G , let \mathcal{F} be a set of graphs, and let k be a positive integer.

Definition 1 (*(\mathcal{F}, k) -boundary-reducibility*) We say that H is an (\mathcal{F}, k) -boundary-reducible subgraph if there exists a set $R \subseteq V(H)$ such that $R \neq \emptyset$ and

- (FIX) for every $v \in R$, $H[R]$ is L -colorable for every $((k - \deg_G + \deg_R) \downarrow v)$ -assignment L , and
- (FORB) for every \mathcal{F} -free set $I \subseteq R$ of size at most $k - 2$, $H[R]$ is L -colorable for every $(k - \deg_G + \deg_R - 1_I)$ -assignment L .

Definition 2 (*Weak (\mathcal{F}, k) -boundary-reducibility*) We say that H is weakly (\mathcal{F}, k) -boundary-reducible if it satisfies (FORB) and there exists at least one vertex v satisfying (FIX) from Definition 1. In this case, we denote v by $\text{Fix}(H)$.

In both of the preceding definitions, we will occasionally refer to the set $V(H) \setminus R$ as the *boundary* of the configuration and the set R as the *reduced part* of the configuration. Note that (FORB) in particular implies that $\deg_G - \deg_R \leq k - 2$ for all $v \in R$.

Definition 3 (*(\mathcal{F}, k, b) -resolution*) Let G be an \mathcal{F} -free graph with lists of size k . An (\mathcal{F}, k, b) -resolution of G is a set $\{G_0, G_1, \dots, G_M\}$ of subgraphs of G such that for $i \geq 1$, H_i is an induced (\mathcal{F}, k) -boundary-reducible subgraph of G_{i-1} with reduced part R_i and

$$G_i := G - \bigcup_{j=1}^i R_j.$$

Additionally, for each $i \geq 1 |R_i| \leq b$ and G_M is itself a (\mathcal{F}, k) -boundary-reducible graph with empty boundary and order at most b . For technical reasons, let $G_{M+1} := \emptyset$.

A weak (\mathcal{F}, k, b) -resolution is defined analogously, except that it uses weak (\mathcal{F}, k) -boundary-reducibility in the place of (\mathcal{F}, k) -boundary-reducibility.

The following lemma is the main tool we use for proving weighted ϵ -flexibility.

Lemma 1 (Lemma 13 in [2]) *For integers $k \geq 3$ and $b \geq 1$ and for a set \mathcal{F} of forbidden subgraphs, let G be an \mathcal{F} -free graph with an (\mathcal{F}, k, b) -resolution. Then there exists an $\epsilon > 0$ such that G is weighted ϵ -flexible for lists of size k . Furthermore, if the request is widespread and G has a weak (\mathcal{F}, k, b) -resolution, then G is weakly $(\epsilon \cdot \frac{1}{b})$ -flexible for lists of size k .*

For the proof of Theorem 2 we prove a stronger version of Lemma 1 tailored to the setting of weak flexibility. For this, we define new “enhanced” versions of weak

(\mathcal{F}, k) -boundary-reducibility and of a weak (\mathcal{F}, k, b) -resolution. We will now require $\text{Fix}(H)$ to contain only vertices v satisfying $\deg_G(v) - \deg_R(v) \leq k - 3$. This change will allow us to consider smaller sets for the (FORB) condition.

Definition 4 (*Enhanced weak (\mathcal{F}, k) -boundary-reducibility*) A graph H is *enhanced weakly (\mathcal{F}, k) -boundary-reducible* if there exist non-empty sets $\text{Fix}(H) \subseteq R \subseteq V(H)$ such that

- (FIX) for every $v \in \text{Fix}(H)$, $\deg_G(v) - \deg_R(v) \leq k - 3$ and $H[R]$ is L -colorable for every $((k - \deg_G + \deg_R) \downarrow v)$ -assignment L , and
- (FORB) for every \mathcal{F} -free set $I \subseteq R$ of size at most $k - 3$, $H[R]$ is L -colorable for every $(k - \deg_G + \deg_R - 1_I)$ -assignment L .

Before proceeding further, observe that (FORB) in the enhanced version is easier to check because I is of size at most $k - 3$, instead of $k - 2$ in the non-enhanced version. However, (FIX) in the enhanced version has an additional restriction on the degree of vertices in $\text{Fix}(H)$, which makes it more difficult to satisfy. Note that in general, the (FORB) condition on a single vertex v implies $\deg_G - \deg_R \leq k - 2$. However for vertices in $\text{Fix}(H)$, the (FIX) condition implies $\deg_G - \deg_R \leq k - 3$. In particular, a vertex of degree $k - 2$ is no longer reducible under the enhanced definition. We overcome this obstacle by allowing $(k - 2)$ -vertices in a resolution under special conditions. Forbidding books B_ℓ helps with satisfying these special conditions. By doing this we can have both: a vertex of degree $k - 2$ is reducible in our setting, and in addition we obtain subgraphs H that are reducible under the enhanced weak (\mathcal{F}, k) -boundary-reducibility definition, given that certain special circumstances occur. This rather technical improvement helps substantially in reducing the complexity of the analysis of the discharging process for the graph classes studied in this paper. Note that further generalization of this idea may be possible, but for lack of use in this paper we will not aim to formulate this in the full generality.

For a subgraph H of a graph G , let $N_G(H)$ be the induced subgraph of G on all neighbors of the vertices in H . If G is a graph satisfying the conditions of Definition 4 and $I \subseteq R$ is an \mathcal{F} -free set of size $k - 2$ so that $G[R]$ is L -colorable for every $(k - \deg_G + \deg_R - 1_I)$ -assignment L , then we call I *loose*.

Let G be a graph, H its subgraph and $v \in V(G - H)$. We say that v is *H-tight* if $\deg_G(v) = k - 2$, $N_G(v) \subseteq V(H)$, and $N_G(v)$ is not loose in H .

Definition 5 (*Enhanced weak $(\mathcal{F}, k, b, \beta)$ -resolution*) Let G be an \mathcal{F} -free graph with lists of size k . An *enhanced weak $(\mathcal{F}, k, b, \beta)$ -resolution* of G is a set $\{G_0, G_1, \dots, G_M\}$ of subgraphs of G , such that all the following three conditions hold:

1. For $1 \leq i \leq M$, there exists a subgraph H_i of G_{i-1} satisfying that
 - H_i is an induced enhanced weak (\mathcal{F}, k) -boundary-reducible subgraph of G_{i-1} with reducible part R_i such that $|R_i| \leq b$, or
 - H_i is an induced weak (\mathcal{F}, k) -boundary-reducible subgraph of G_{i-1} with reducible part R_i , such that $|R_i| \leq b$ and for all $v \in \text{Fix}(H_i)$ either

$$|N_{G_{i-1}}(v) \cap H_j| \leq k - 3 \quad \text{or} \quad N_{G_{i-1}}(v) \cap H_j$$

is a loose set in H_j for all $j > i$, or

- H_i is a single vertex with $\deg_{G_{i-1}}(v) = k - 2$.

2. For every $1 \leq i \leq M - 1$,

$$G_i := G - \bigcup_{j=1}^i R_j,$$

G_M is a weak (\mathcal{F}, k) -boundary-reducible graph with empty boundary and order at most b , $G_{M+1} := \emptyset$, and $H_{M+1} := G_M$.

3. The following is satisfied:

(TIGHT) For every $1 \leq j \leq M$, there are at most β different H_j -tight vertices v_i , where $V(H_i) = \{v_i\}$ with $i < j$.

Note that in Definition 5, H_i can be H_j -tight only if H_i is a single vertex with $\deg_{G_{i-1}}(v) = k - 2$. A natural way to satisfy (TIGHT) condition is to show that whenever there is an H_j such that more than β subgraphs $H_{a_1}, \dots, H_{a_\beta}, H_{a_{\beta+1}}$ are H_j -tight, then

$$H_j \cup \bigcup_{i \in \{1, \dots, \beta, \beta+1\}} H_{a_i} \in \mathcal{F}.$$

If two adjacent vertices have many common neighbors, we get a book, which will be in \mathcal{F} .

We are now ready to state and prove our main lemma.

Lemma 2 (Reducible configurations for weak flexibility) *For integers $k \geq 4$, $b \geq 1$, $\beta \geq 0$, and for a set \mathcal{F} of forbidden subgraphs, let G be a \mathcal{F} -free graph with an enhanced weak $(\mathcal{F}, k, b, \beta)$ -resolution. Then, there exists an $\epsilon > 0$ such that G is weakly ϵ -flexible for lists of size k .*

The proof of the lemma is similar to the proof of Lemma 1 as it appeared in [2, Lemma 13]. In particular, we explicitly formulate a few arguments in their proof as a separate claim (Claim 4 below) that we use in our proof. We will also need the following Lemma 3, which is Lemma 12 in [2].

Let G be a graph with a weak (\mathcal{F}, k, b) -resolution \mathcal{R} . Let $\text{AllFix}(G)$ denote the union of all $\text{Fix}(H)$ over all reducible subgraphs H in the resolution \mathcal{R} .

Lemma 3 (Lemma 12 in [2]) *Let b be an integer. Let G be a graph with list assignment L of size k on $V(G)$. Suppose G has a weak (\mathcal{F}, k, b) -resolution, G is L -colorable, and there exists a probability distribution on the L -colorings φ of G such that*

for every $v \in \text{AllFix}(G)$ and $c \in L(v)$, $\text{Prob}[\varphi(v) = c] \geq \varepsilon$. Then G with L is weakly $(\varepsilon \cdot \frac{1}{b})$ -flexible.

Proof of Lemma 2 For $1 \leq j \leq M + 1$, let \mathcal{H}_j be the set of all H_j -tight subgraphs where the (TIGHT) property applies. Let $H_i \in \mathcal{H}_j$ for some i and j . This means that H_i is a single vertex with $k - 2 \geq 2$ neighbors in H_j . Hence $\mathcal{H}_i = \emptyset$.

Now, we refactor the enhanced weak $(\mathcal{F}, k, b, \beta)$ -resolution \mathcal{R} into an enhanced weak $(\mathcal{F}, k, b + \beta, 0)$ -resolution \mathcal{R}' . To do so, we attach all H_j -tight subgraphs to H_j and thus we create a larger configuration H'_j . The vertices in tight subgraphs are not part of any Fix set. Formally

$$H'_j := \begin{cases} \emptyset & \text{if exists } i \text{ such that } H_j \in \mathcal{H}_i \\ H_j \cup \bigcup_{H \in \mathcal{H}_j} H & \text{otherwise} \end{cases}$$

and $\text{Fix}(H'_i) = \text{Fix}(H_i)$ if $H'_i \neq \emptyset$ and $\text{Fix}(H'_i) = \emptyset$ otherwise. Observe that by the (TIGHT) property, the size of the resulting H'_j will be upper-bounded by $b + \beta$ and that H'_j is enhanced weakly (\mathcal{F}, k) -boundary-reducible or only weakly (\mathcal{F}, k) -boundary-reducible (provided its neighbourhood is always a loose or small set) if it is not empty. We simultaneously remember both \mathcal{R} and \mathcal{R}' , since each time we are using H'_j (or G'_j) we are referring to \mathcal{R}' and each time we are using H_j (or G_j) we are referring to \mathcal{R} .

The next step is to create a probability distribution on L -colorings φ of G_i for all i starting with G'_M . Let $p = k^{-(b+\beta)}$ and $\varepsilon' = p^{k-1}$. We are going to show that each i satisfies the following properties:

- (i) for every $v \in \text{AllFix}(G'_i)$ and a color $c \in L(v)$, the probability that $\varphi(v) = c$ is at least ε' , and
- (ii) for every color c and every \mathcal{F} -free set I in G'_i of size at most $k - 3$, the probability that $\varphi(v) \neq c$ for all $v \in I$ is at least $p^{|I|}$, and
- (iii) for every color c and every loose \mathcal{F} -free set I in G'_i of size exactly $k - 2$, the probability that $\varphi(v) \neq c$ for all $v \in I$ is at least $p^{|I|}$.

Note that for G'_{M+1} all of the properties trivially hold. Note that Property (i) on $G'_0 = G_0 = G$ immediately implies that G with L is weakly $(\varepsilon' \cdot \frac{1}{b})$ -flexible by Lemma 3 and therefore weakly ε -flexible for $\varepsilon = \frac{\varepsilon'}{b}$. □

We will make use of the following claim proven implicitly in [2].

Claim 4 (Implicit in the proof Lemma 13 in [2]) *Suppose that we have an enhanced weak $(\mathcal{F}, k, b + \beta, 0)$ -resolution and a probability distribution on L -colorings of G'_{i+1} satisfying Properties (i), (ii), and (iii) on G'_{i+1} .*

If for each vertex $v \in \text{Fix}(H'_i)$ and for each $I = N(v) \cap H'_j$ where $j > i$ one of the following holds:

- (a) $|I| = k - 2$ and I is loose in H'_j , or
- (b) $|I| < k - 2$

then there exists a probability distribution on L -colorings of G'_i such that Properties (i), (ii), and (iii) are satisfied on G'_i .

In order to use Claim 4, we need verify (a) and (b). If H'_i is not a single vertex v with $\deg_{G_i}(v) = k - 2$, then (a) or (b) hold by the definition of H'_i . Hence we need to check the case of H'_i being a single vertex v with $\deg_{G_i}(v) = k - 2$. We do it by showing v is not H'_j -tight for any $j \geq i$ in the following claim. It implies that for H'_i , either (a) or (b) is satisfied. In particular, we will show that we got rid of all tight subgraphs when we refactored \mathcal{R} into \mathcal{R}' .

Claim 5 *There are no $i < j$ such that H'_i is H'_j -tight.*

Proof Suppose for contradiction that H'_i is H'_j -tight for some $i < j$. By the definition, H'_i is one vertex v with degree $k - 2$ in G_{i-1} . By the definition of \mathcal{R}' , v is not H_ℓ -tight for any $\ell > i$. In particular, v is not H_j -tight. Since v is H'_j -tight, \mathcal{H}_j is not empty. Hence H'_j is a union of H_j and vertices W , where every $w \in W$ has $k - 2$ neighbors in H_j . Since v is not H_j -tight, it has at most $k - 3$ neighbors in H_j and at least one in W . Notice that every vertex w in W has $k - 2$ neighbors in \mathcal{H}_j hence a list of $k - 1$ colors suffices for extending any coloring of \mathcal{H}_j to w greedily. This and the (FORB) property for H_j imply that v is not H'_j -tight because $N(v)$ in H'_j is loose, which is a contradiction. ◇

We conclude that Claim 5 enables us to use Claim 4 directly on \mathcal{R}' . This finishes the proof of Lemma 2. □

For a positive integer d , a d -vertex, a d^+ -vertex, and a d^- -vertex are a vertex of degree d , at least d , and at most d , respectively. A d -face, a d^+ -face, and a d^- -face are defined analogously. A (d_1, d_2, d_3) -face is a 3-face where the degrees of the vertices on the face are d_1, d_2, d_3 . We will sometimes call 3-faces *triangles*. A *diamond* D is a graph isomorphic to K_4 minus an edge. The 2-vertices of D are called the *side vertices*, and the 3-vertices are called the *middle vertices* of D . Let G be a graph. By $T(a, b, c)$ we denote a triangle in G with vertices of degree a, b , and c in G , and by $Dia(a - b, c, d)$ a diamond in G with middle vertices of degrees a and b and side vertices of degrees c and d .

Lemma 4 *Let G be a plane $\{C_5, C_6, C_7\}$ -free graph. Suppose that v is the middle vertex of k distinct diamonds, and v is adjacent to m faces of size 3 or 4 that are not part of a diamond in which v is a middle vertex. Then $\deg(v) \geq 3k + 2m$ and $k \leq \lfloor \frac{\deg(v)}{3} \rfloor$.*

Proof If not, then v is adjacent to three faces f, g, h , each of them of size at most 4, such that f, g share an edge and g, h share an edge. But this induces a cycle C_i with $5 \leq i \leq 7$, a contradiction. □

In all the figures in the paper, black vertices have all their incident edges drawn, whereas a white vertex may have more edges incident than drawn (since white vertices are in the boundary).

4 Proof of Theorem 1

4.1 Reducible Configurations

Let $\mathcal{F} = \{K_4, C_5, C_6, C_7, B_5\}$. In this section we will provide a handful of $(\mathcal{F}, 4)$ -boundary-reducible configurations.

Lemma 5 *The following configurations are $(\mathcal{F}, 4)$ -boundary-reducible. See Figs. 2 and 3 for reference. If boundary is not mentioned, it is empty.*

- (C1) *A vertex of degree at most 2.*
- (C2) *Three 3-vertices appearing on a path of length 2.*
- (C3) *The triangle $T(3, 3, 3)$.*
- (C4) *Let u be a 3-vertex adjacent to the middle 4-vertex of the diamond $D = \text{Dia}(4 - 3, 4, 5^+)$. Let v denote the 5^+ -vertex that is a side vertex of D . Then $D \cup \{u\}$ is reducible with boundary v .*
- (C5) *$\text{Dia}(3 - 3, 5^+, 5^+)$ with 5^+ -vertices in the boundary.*
- (C6) *$\text{Dia}(3 - 5^+, 3, 5^+)$ with 5^+ -vertices in the boundary.*
- (C7) *The diamond $\text{Dia}(5 - 4, 3, 3)$.*
- (C8) *Let $D_1 = \text{Dia}(4 - 4, 5, 3)$ and $D_2 = (5 - 3, 4, 4^+)$ be two diamonds sharing the same 5 -vertex. Let v denote the 4^+ -vertex that is a side vertex of D_2 . Then the subgraph $D_1 \cup D_2$ is reducible with v in the boundary.*
- (C9) *Let $D_1 = D_2 = \text{Dia}(3 - 4, 4, 5^+)$ be two diamonds whose middle 4-vertices are connected by an edge. Let v_1 and v_2 denote the two 5^+ -vertices that are the side vertices of D_1 and D_2 . Then the subgraph $D_1 \cup D_2$ is reducible with v_1 and v_2 in the boundary.*
- (C10) *Let $D_1 = \text{Dia}(4 - 3, 5^+, 5)$ and $D_2 = \text{Dia}(5 - 3, 4, 4^+)$ be two diamonds sharing a middle 5-vertex. Let v_1 denote the 5^+ -vertex that is a side vertex of D_1 and let v_2 denote the 4^+ -vertex that is a side vertex of D_2 . Then the subgraph $D_1 \cup D_2$ is reducible with v_1 and v_2 in the boundary.*
- (C11) *A diamond $\text{Dia}(4 - 4, 3, 4)$ along with a 3-vertex adjacent to one of the middle 4-vertices.*
- (C12) *Let $D_1 = \text{Dia}(3 - 4, 4, 5^+)$ and $D_2 = \text{Dia}(4 - 4, 4, 3)$ be two diamonds whose middle 4-vertices are connected by an edge. Let v denote the 5^+ -vertex that is a side vertex of D_1 . Then the subgraph $D_1 \cup D_2$ is reducible with v in the boundary.*
- (C13) *Let $D_1 = \text{Dia}(3 - 5, 5, 5)$ and $D_2, D_3 = \text{Dia}(5 - 3, 4, 5^+)$ be two diamonds where the two side 5-vertices of D_1 are middle vertices of D_2 and D_3 . Let v_1 and v_2 denote the two side 5^+ -vertices of D_2 and D_3 , respectively. Then $D_1 \cup D_2 \cup D_3$ is reducible with v_1 and v_2 in the boundary.*





Fig. 2 Reducible configurations for Theorem 1 (Part 1). The labels give the list sizes remaining after accounting for the external neighbors and boundary vertices

Fig. 3 Reducible configurations for Theorem 1 (Part 2). The labels give the list sizes remaining after accounting for the external neighbors and boundary vertices



We wish to point out that the reducible configurations are meant to be induced subgraphs by definition, and we will use them as such in the discharging part of the proof. The only configuration, where two external edges can be identified is (C2) and it gives (C3), which we explicitly list. It can be straightforwardly checked that no identification of vertices in (C1)–(C15) is possible since otherwise, it creates a forbidden subgraph.


Proof of Lemma 5 It is straightforward to check that each configuration (C1)–(C15) satisfies the (FIX) and (FORB) conditions in Definition 1. However, checking all the cases is rather tedious. Hence we developed a simple computer program that does it, see <http://lidicky.name/pub/flexibility>.¹ In particular, a greedy coloring works in all cases. For an interested reader who wishes to check some cases by hand, we added list sizes to Figs. 2 and 3. We also provide here proofs showing that (C2) and (C5) are $(\mathcal{F}, 4)$ -boundary-reducible. Together, these two configurations demonstrate how to prove that the remaining configurations are reducible.

The two reducible configurations H_1 and H_2 corresponding to (C2) and (C5), respectively are depicted in Fig. 4. The reduced parts $R_1 = H_1$ and $R_2 \subset H_2$ are provided as well. Finally, we have labeled each vertex in the figure with the value of the function $4 - \deg_{H_i} + \deg_{R_i}$ for $i \in \{1, 2\}$.

By definition, checking the (FIX) condition for any subgraph H with reducible part R is equivalent to showing that for each $v \in V(R)$, R can be properly colored after assigning each vertex a list of size $((4 - \deg_H + \deg_R) \downarrow v)$. It is clear by inspection that this is the case for (C2) = $H_1 = R_1$, and hence we only need to check the (FORB) condition for (C2). Since (FIX) is already verified, it implies (FORB) for subsets of size one in R . It remains to verify (FORB) for subsets of size two in R .

¹ This program is also available as a part of the sources in our arXiv submission [9] (file `reducible_configurations.sage`).

Fig. 4 Reducible configurations (C2) and (C5). The reduced parts consist of the black vertices



If we apply (FORB) to a and c , then both a and c will be left with one available color in their lists. Vertex b still has three colors in its list. Therefore, we can greedily color a , c , and b in this order to obtain a proper coloring for (C2). If we apply (FORB) to a and b , then the color for a will be fixed, and each of b and c will be left with two possible colors. Therefore, we can greedily color a , b , and c in this order to obtain a proper coloring for (C2). By symmetry, the case of applying (FORB) to b and c is also verified, implying that (C2) is reducible.

Let H_2 be a subgraph of G isomorphic to (C5). Let $R_2 \subset H_2$ denote the reducible part of (C5), i.e. the subgraph of H_2 induced by vertices u_1, \dots, u_4 . For each $i = 1, \dots, 4$, we will check the (FIX) condition for u_i . Let L_i be an arbitrary list assignment where each vertex in R is assigned a list of size $((4 - \deg_H + \deg_R) \downarrow u_i)$. We will now show that R can be properly colored. In each case we list the order of vertices in greedy coloring.

- $L_1 : u_1, u_2, u_4, u_3.$
- $L_2 : u_2, u_1, u_4, u_3.$
- $L_3 : u_3, u_4, u_2, u_1.$
- $L_4 : u_4, u_3, u_2, u_1.$

Next we need to verify that H satisfies the (FORB) condition. However, only one subset of R of size two is \mathcal{F} -free: $\{u_1, u_2\}$. In that case R can be colored greedily in the following order u_1, u_2, u_4, u_3 . Thus, (C5) is a reducible configuration. \square

4.2 Discharging

In this section we prove the following lemma, which by Lemma 1 implies Theorem 1.

Lemma 6 *Let G be a connected $\{K_4, C_5, C_6, C_7, B_5\}$ -free plane graph. Then G contains at least one of the reducible configurations (C1)–(C13).*

Proof Suppose for contradiction that G is a connected $\{K_4, C_5, C_6, C_7, B_5\}$ -free plane graph that contains none of the configurations (C1)–(C13). We use discharging to obtain a contradiction with Euler’s formula.

We denote the initial charge by ch . For every vertex v , we let $ch(v) = \deg(v) - 4$, and for every face f we let $ch(f) = \ell(f) - 4$, where $\ell(F)$ is the length of the facial walk. For convenience we will also assign charge to the edges of G . The initial charge is 0 for each edge. By Euler’s formula, the total sum of initial charges is -8 .

We sequentially apply the following rules (see also Figs. 5 and 6) that move the charge around, while keeping the sum of charges unchanged. The charge at the end is called the *final charge*. The final charges will be all nonnegative, contradicting that their sum is -8 .

- (R1) Every 8^+ -face sends charge $1/2$ to every incident 3-face and 4-face for every edge they have in common.
- (R2) For every edge e that is not incident with any 3-face or 4-face the following applies. If e is a bridge, e receives charge 1 from the unique face incident with e . If e is not a bridge, e receives charge $1/2$ for each of the two faces incident to e .
- (R3) For every vertex u and an incident edge $e = uz$ with charge 1:
 - (R3a) If u and z are both 3-vertices, then e sends charge $1/2$ to u .
 - (R3b) If u is a 3-vertex and z is 4^+ -vertex, then e sends charge 1 to u .
 - (R3c) If z is a 4^+ -vertex, u is the middle 4-vertex of the diamond $Dia(4 - 3, 4, 4^+)$, and v is the 3-vertex on this diamond, then e sends 1 to v .
 - (R3d) If z is a 4^+ -vertex, u is one of the middle 4-vertices of the diamond $Dia(4 - 4, 3, 4)$, and v is the 3-vertex on this diamond, then e sends $1/2$ to v .
- (R4) Every 4-face sends charge 1 to every incident 3-vertex.
- (R5) Let f be a 3-face that is not part of a diamond. If exactly one vertex u of f has degree 3, then f sends $1/2$ to u .
- (R6) The following rules apply for a 5-vertex u . If u is a middle vertex in
 - (R6a) $Dia(5 - 3, 4, 4)$ or $Dia(5 - 3, 5, 4)$, then u sends 1 to the middle 3-vertex;
 - (R6b) $Dia(5 - 3, 6^+, 4^+)$ or $Dia(5 - 3, 5, 5)$, then u sends $1/2$ to the middle 3-vertex;
 - (R6c) $Dia(5 - 5^+, 3, 3)$, then u sends $1/4$ to every of the two side 3-vertices;
 - (R6d) $Dia(5 - 4^+, 4^+, 3)$, then u sends $1/2$ to the side 3-vertex.
- (R7) The following rules apply for every 5-vertex u and a diamond D , where v is a side vertex of D . If D is
 - (R7a) $Dia(4 - 4, 5, 3)$, then u sends $1/2$ to the side 3-vertex;
 - (R7b) $Dia(4 - 3, 5, 5^+)$, then u sends $1/2$ to the middle 3-vertex;
 - (R7c) $Dia(5 - 3, 5, 4^+)$ and u has not already sent 1 to another diamond under rule (R6a), then u sends $1/2$ to the middle 3-vertex.
- (R8) The following rules apply for every 6^+ -vertex u and a diamond D , where u is a side vertex of D . If D is

- (R8a) $Dia(5 - 3, 6^+, 4^+)$, then u sends $1/2$ to the middle 3-vertex;
 - (R8b) $Dia(4 - 4, 6^+, 3)$, then u sends $1/2$ to the side 3-vertex;
 - (R8c) $Dia(4 - 3, 6^+, 4^+)$, then u sends $1/2$ to the middle 3-vertex.
- (R9) The following rules apply for every 6^+ -vertex u and a diamond D , where u is a middle vertex of D . If D is
- (R9a) $Dia(6^+ - 4^+, 3, 3)$ then u sends $1/2$ to each of the side vertices;
 - (R9b) $Dia(6^+ - 4^+, 4^+, 3)$, then u sends 1 to the side 3-vertex;
 - (R9c) $Dia(6^+ - 3, 4^+, 4^+)$ then u sends 1 to the other middle 3-vertex.

Claim 6 *The final charge of every face of G is nonnegative.*

Proof Given that G does not contain any faces of length 5, 6 or 7, we consider 8^+ -faces, 4-faces, and 3-faces as three separate cases covering everything.

Suppose that f is an 8^+ -face. Then the initial charge of f is equal to $\ell(f) - 4$. By (R1) and (R2), f sends at most $\frac{1}{2}$ for each of these edge that is not a bridge and charge 1 to each bridge by (R2). This means that f sends at most $\left\lfloor \frac{\ell(f)}{2} \right\rfloor \leq \ell(f) - 4$ total charge. Since (R1) and (R2) are the only rules requiring an 8^+ -face to send out charge, every 8^+ -face has nonnegative final charge.

Suppose that f is a 4-face. Then f has its initial charge 0. Since $C_5, C_6,$ and C_7 are forbidden subgraphs, f must be incident with four 8^+ -faces. By (R1), each face sharing an edge with f sends charge $\frac{1}{2}$ to f for every edge they have in common, leaving f with a total charge of 2 before applying (R2)–(R9). Given that (C2) is a reducible configuration, f cannot contain more than two 3-vertices. Thus, (R4) applies to f at most twice, which decreases the charge at f by at most 2. Since no other rules apply to 4-faces, f has nonnegative final charge.

Next suppose that f is a 3-face that is not contained in a diamond. Every face incident to f must be an 8^+ -face since $C_5, C_6,$ and C_7 are forbidden subgraphs. This




Fig. 5 Discharging rules described in Lemma 6 used for Theorem 1 (Part 1)




Fig. 6 Discharging rules described in Lemma 6 used for Theorem 1 (Part 2)

means that after applying (R1), f has charge $\frac{1}{2}$. Among rules (R2)–(R9), only (R5) requires a 3-face to send out charge. If (R5) applies to f , then it only requires f to send a charge of $\frac{1}{2}$. This means that f has nonnegative final charge.

Lastly, assume that f is a 3-face contained in a diamond D . Then f shares one edge with another 3-face. Since C_5 , C_6 , and C_7 are forbidden subgraphs, f shares its other two edges with 8^+ -faces. By (R1), f receives charge at least $\frac{1}{2}$ for each edge it shares with an 8^+ -face, leaving f with charge at least 0 before applying (R2)–(R9). None of these rules, however, demand charge from a 3-face that is contained in a diamond, implying that f will end with nonnegative charge. As we have considered all possible faces in G , this completes the proof of Claim 6. \diamond

Claim 7 *The final charge of every edge of G is nonnegative.*

Proof Let $e = uz$ be an edge of G . If e is incident with a 3-face or a 4-face, then none of the rules apply to e and there is nothing to prove. Otherwise, e has charge 1 after applying (R2). As (R3) is the only rule that requires any edge to send out charge, it suffices to verify that e will never be asked to give more than 1 charge under (R3).

If u and z are both 3-vertices, then only (R3a) applies to e and the edge sends exactly $\frac{1}{2}$ to each of u and z . If u is a 3-vertex and z is a 4^+ -vertex, then only (R3b) applies, and e send exactly 1 to u .

If u is a 4^+ -vertex and z is a 5^+ -vertex, then (R3) does not apply with z and e sends charge at most 1 using either (R3c) or (R3d).

The remaining case is that both u and z are 4-vertices. The rules demand e to send charge more than 1 if by symmetry (R3c) applies with u and one of (R3c) and (R3d) applies with z . However, this would give reducible configurations (C9) and (C12), respectively. Therefore, no edge in G that begins with charge 1 will ever be asked to send out more than 1 total charge, completing the proof of Claim 7. \diamond

Claim 8 *The final charge of every 4^+ -vertex is nonnegative.*

Proof Suppose that v is a 4-vertex. The initial charge of v is 0, and there are no rules requiring v to send out charge, so v will end with nonnegative charge.

Next suppose that v is a 5-vertex. Then the initial charge of v is 1. Only (R6) and (R7) require a 5-vertex to distribute charge. Therefore, we may assume that v is incident with at least one diamond. Given that G does not contain any C_5 , C_6 , C_7 , or B_5 subgraphs, v is incident with at most two diamonds.

First suppose that v is incident with exactly one diamond D . If v is a middle vertex of D then only (R6) applies to v , and if v is a side vertex of D then only (R7) applies to v . As neither of these two rules will require v to send out charge more than 1, v will end with nonnegative charge.

Next suppose that v is incident with two diamonds D_1 and D_2 . Since G does not contain any C_5 , C_6 , C_7 , or B_5 subgraphs, D_1 and D_2 must be edge disjoint. Since $\deg(v) = 5$, v cannot be a middle vertex of both diamonds. If v is a side vertex of both diamonds, then only (R7) applies to v . As (R7) will not require v to send charge more than $\frac{1}{2}$ to either diamond, v will end with nonnegative charge.

Therefore, we may assume that v is a middle vertex of D_1 and a side vertex of D_2 . In this case, it is possible that both (R6) and (R7) apply to v . Among the subcases of (R6), only (R6a) requires v to send out charge for more than $\frac{1}{2}$, and (R7) will never ask v send out charge more than $\frac{1}{2}$. Given that configuration (C8) is reducible, (R6a) cannot apply with (R7a). Next, given that configuration (C10) is reducible, (R6a) cannot apply with (R7b). By assumption of (R7c), (R6a) cannot apply with (R7c). Therefore v is never asked to send more than 1, implying that v will end with non-negative charge.

Now suppose that v is a 6^+ -vertex. The only rules that apply to v are (R8) and (R9). Under these rules, v sends at most 1 to all diamonds that contain v as a middle vertex, and v sends at most $1/2$ to all diamonds that contain v as a side vertex. Assume that v is the middle vertex of k distinct diamonds, and incident to m other faces of size 3. By Lemma 4, the final charge of v is at least

$$\begin{aligned} \deg(v) - 4 - k - \frac{m}{2} &= \deg(v) - 4 - \frac{3k + 2m}{4} - \frac{k}{4} \geq \frac{3 \deg(v)}{4} - 4 - \frac{1}{4} \left\lfloor \frac{\deg(v)}{3} \right\rfloor \\ &\geq \frac{2 \deg(v)}{3} - 4, \end{aligned}$$

and $\frac{2 \deg(v)}{3} - 4$ is nonnegative whenever $\deg(v) \geq 6$. This completes the proof of Claim 8. ◇

Claim 9 *The final charge of every 3-vertex that is not contained in a diamond is nonnegative.*

Proof Let v be a 3-vertex that is not contained in a diamond. Then the initial charge of v is -1 . As there are no rules requiring v to send out charge, we only need to verify that v will receive charge at least 1. First suppose that v is not incident to any 3-faces or 4-faces. Then each of the three edges incident to v receive charge 1 under (R2). Next, each of these edges sends $\frac{1}{2}$ to v by (R3), leaving v with a charge of $\frac{1}{2}$.

Now suppose that v is incident to at least one 4-face f . By (R4), v receives 1 from f and we are done. Therefore, we may assume that v is not incident to any 4-face, and that v is incident to at least one 3-face T . By assumption, T is not contained in a diamond.

- Case 1: T contains another 3-vertex. In this case, v must be adjacent to a 4^+ -vertex u that is not contained T , since (C2) is reducible. Then by (R3b), v receives a charge of 1 from the edge uv .
- Case 2: T contains two 4^+ -vertices. Again, let u be the neighbor of v that is not contained in T . Here, v will receive at least $\frac{1}{2}$ from (R3). With that being said, T will send the remaining $\frac{1}{2}$ to v under (R5).

This completes the proof of Claim 9. ◇

Claim 10 *The final charge of every 3-vertex that is incident to a diamond is nonnegative.*

Proof Assume that v is a 3-vertex incident to a diamond D . Since (C2) and (C3) are reducible, there is at most one other 3-vertex incident to D . Since the initial charge of v is -1 , and there is no rule requiring a 3-vertex to send charge, it suffices to show that v will always receive charge at least 1 after applying rules (R1)–(R9). We consider the following cases.

Case 1 v is the only 3-vertex incident to D and v is a side vertex of D . Given the list of forbidden subgraphs in G , the other two faces incident to v must be 8^+ -faces. Hence by (R3), v receives charge at least $\frac{1}{2}$ from the only edge incident to v that is not a part of D . There are three subcases to Case 1 showing how v gets another $\frac{1}{2}$ of charge.

1. $D = \text{Dia}(4 - 4, 4, 3)$

Let x and y denote the two middle vertices of D . Since (C11) is reducible, each of the neighbors of x and y that are not contained in D must be 4^+ -vertices. Therefore by (R3d), v receives $\frac{1}{2}$ from the each of two edges incident to x and y that are not contained in D .

2. $D = \text{Dia}(4 - 4, 5^+, 3)$

Let u denote the other side vertex in D . If u is a 5-vertex, then v receives $\frac{1}{2}$ from u by (R7a). If u is a 6^+ -vertex, then v receives $\frac{1}{2}$ from u by (R8b).

3. $D = \text{Dia}(5^+ - 4^+, 4^+, 3)$

Let u denote the 5^+ -vertex that is the middle vertex of D . If $d(u) = 5$, then u sends charge $\frac{1}{2}$ to v by (R6d). If $d(u) \geq 6$, then u sends charge $\frac{1}{2}$ by (R9b).

In all three cases, the final charge of v is nonnegative.

Case 2 v is the only 3-vertex incident to D and v is a middle vertex of D . There are four subcases to Case 2. In each v receives charge 1 which leads to nonnegative final charge.

1. $D = \text{Dia}(4 - 3, 4, 4^+)$

Let u denote the middle 4-vertex of D . Since (C4) is reducible, the unique neighbor z of u not contained in D must be a 4^+ -vertex. Therefore, the edge uz will send charge 1 to v by (R3c), leaving v with nonnegative charge.

2. $D = \text{Dia}(4 - 3, 5^+, 5^+)$

Let x and y denote the two side vertices of D . If $d(x) = 5$, then x will send charge $\frac{1}{2}$ to v by (R7b). If $d(x) \geq 6$, then x will send charge $\frac{1}{2}$ to v by (R8c). As the rules apply to y identically as they do to x , it follows that v will end with nonnegative charge.

3. $D = \text{Dia}(5 - 3, 4^+, 4^+)$

Let u denote the middle 5-vertex of D and let x and y denote each of the side 4^+ -vertices of D with $d(x) \leq d(y)$. If $d(x) = d(y) = 4$, then u sends charge 1 to v by (R6a). If $d(x) = 4$ and $d(y) = 5$, then u sends charge 1 to v by (R6a). If $d(x) = 4$ and $d(y) \geq 6$, then u sends charge $\frac{1}{2}$ to v by (R6b) and y sends $\frac{1}{2}$ to v by (R8a). If $d(x) = d(y) = 5$, then u sends charge $\frac{1}{2}$ to v by (R6b). Since (C13) is reducible, x and y will not both send charge 1 to a vertex of a different diamond under rule

(R6a). Therefore, v receives charge $\frac{1}{2}$ from either x or y by (R7c). If $d(y) \geq 6$, y sends charge $\frac{1}{2}$ to v by (R8a) and u sends charge $\frac{1}{2}$ to v by (R6b). This leaves v with nonnegative charge.

4. $D = \text{Dia}(6^+ - 3, 4^+, 4^+)$

Let u denote the middle vertex of D . Then u sends charge 1 to v by (R9c). This leaves v with nonnegative charge.

Case 3 There are two 3-vertices incident to D , one of which is v . Let x denote the other 3-vertex incident to D . Since (C5) and (C6) are reducible configurations, both x and v are side vertices of D . Since (C7) is reducible, we may assume that if one of the middle vertices of D is a 4-vertex, then the other middle vertex is a 6^+ -vertex. There are two subcases to Case 3.

1. $D = \text{Dia}(6^+ - 4, 3, 3)$.

Let u denote the 6^+ -vertex incident to D . By (R9a), u sends charge $\frac{1}{2}$ to v . Since v is a side vertex of D , and $\text{deg}(v) = 3$, it follows that v is incident to exactly one edge that is not contained in D . By (R3), v will receive charge at least $\frac{1}{2}$ from this edge, leaving v with nonnegative charge. The case of x is symmetric.

2. $D = \text{Dia}(5^+ - 5^+, 3, 3)$

Since v is a side vertex of D , and $\text{deg}(v) = 3$, it follows that v is incident to exactly one edge that is not contained in D . By (R3), v will receive charge at least $\frac{1}{2}$ from this edge. Let a and b denote the middle vertices of D . If $d(a) = 5$, then v receives charge $\frac{1}{4}$ from a by (R6c). If $d(a) \geq 6$, then v receives charge $\frac{1}{2}$ from a by (R9a). As the rules apply to b identically as they do to a , it follows that v will end with nonnegative charge. Again, the case of x is symmetric.

Since we have covered all cases where v is contained in a diamond, this completes the proof of Claim 10. \diamond

Claims 6–10 show that the final charge of every vertex, face, and edge is nonnegative. Hence the sum of the charges is also nonnegative, which is a contradiction with the sum of the initial charges being -8 . This finishes the proof of Lemma 6. \square

5 Proof of Theorem 2

5.1 Reducible Configurations

We will use the following list of enhanced weakly reducible configurations. See Fig. 7 for illustration of these configurations.

- (D1) A vertex of degree at most 2.
- (D2) $T(3, 3, 3)$ and $T(3, 3, 4)$.

Fig. 7 Reducible configurations for Theorem 2. The labels give the list sizes remaining after accounting for the external neighbors and boundary vertices. The vertices whose colors cannot be fixed are drawn as squares. These vertices cannot be fixed because either their coloring does not extend or they have two external neighbors, with the one exception being (D4)

- (D3) Two diamonds $D_1 = Dia(6 - 3, 4, 3)$ and $D_2 = Dia(6 - 3, 4, 4)$ sharing a middle 6-vertex.
- (D4) $Dia(3 - 3, 4^+, 4^+)$ where the side vertices are in the boundary.
- (D5) $Dia(4 - 5, 3, 3)$.
- (D6) $Dia(4 - 3, 4, 4)$.
- (D7) $Dia(5 - 3, 4, 4)$ with another 3-vertex adjacent to the 5-vertex.
- (D8) $Dia(5 - 5, 3, 3)$ with another 3-vertex adjacent to one of the 5-vertices.
- (D9) Three 3-vertices u, v, w such that uv and vw are edges, and u and w are independent.
- (D10) $Dia(5 - 3, 4, 3)$.
- (D11) $T(5, 3, 3)$ with another 3-vertex adjacent to the 5-vertex.
- (D12) Two triangles $T_1 = T_2 = T(6, 3, 3)$ sharing the 6-vertex.

In the next section we will prove the following theorem, showing that (D1)–(D12) are unavoidable. We remark that no identification of vertices in (D1)–(D12) is possible since otherwise, it creates a forbidden subgraph. It is possible that some external edges can be identified in (D8), (D9), and (D11). We explicitly list those cases in Fig. 7 as (D8'), (D9'), (D11'), and (D11'').

Theorem 11 *Every $\{K_4, C_5, C_6, C_7\}$ -free planar graph contains one of (D1)–(D12).*

Let $\mathcal{F} = \{K_4, C_5, C_6, C_7, B_\ell\}$ for any fixed ℓ . We will show that (D2), (D3), and (D5)–(D12) are enhanced weakly $(\mathcal{F}, 4)$ -boundary-reducible configurations. We will also show that (D4) is only a weakly $(\mathcal{F}, 4)$ -boundary-reducible configuration. However, the only neighbors of the vertices in the reducible part of (D4) are the non-adjacent vertices in the boundary, and all non-adjacent pairs that do not forbid \mathcal{F} in each of (D1)–(D12) form loose sets, implying that the condition required by the enhanced weak resolution is satisfied.

In order to use Lemma 2, we want to have an enhanced weak $(\mathcal{F}, 4, b, \beta)$ -resolution for some β and b . First, we check the condition (TIGHT) in the following lemma.

Lemma 7 *Let G be an \mathcal{F} -free graph containing H , where H is one of (D2)–(D12). The number of H -tight vertices is at most $\beta \leq 10\ell$.*

Proof Let H be one of (D2)–(D12) and v be an H -tight vertex adjacent to u and w in H . First, suppose that u and w are not adjacent. There are no non-edges in (D2) and (D4). The non-edge in (D9) forms a loose set. By inspection of each pair of non-adjacent vertices in (D5)–(D8) and (D10)–(D12), we observed that if v was adjacent to any of these pairs, we would obtain a C_5 or a C_6 , contradicting that G is \mathcal{F} -free.



















Fig. 8 Configuration (D10)



Second, suppose that uw is an edge. As B_ℓ is in \mathcal{F} , the number of H -tight vertices for uv is at most $\ell - 3$. Since H is one of (D2)–(D12), it has at most 10 edges. This bounds that the total number of H -tight vertices as $\beta \leq 10(\ell - 3) \leq 10\ell$. \square

Lemma 8 *The configurations (D2), (D3), (D5)–(D12) are enhanced weakly $(\mathcal{F}, 4)$ -boundary-reducible. See Fig. 7 for illustration.*

Proof Given the rules for enhanced weak (\mathcal{F}, k) -boundary reducibility, it is straightforward to verify that configurations (D2)–(D12) are reducible. We also provide a computer program at <http://lidicky.name/pub/flexibility²> to do so. One notable difference is that the greedy algorithm is not always sufficient. We also added test for Gallai tree, which helped. Just greedy algorithm could end with a diamond, where middle vertices have lists L of size 3 and side vertices would have lists of size 2, which is not a Gallai tree and hence it is L -colorable, but not in a greedy way.

In order to highlight the difference between regular and weak reducibility, we will give a short proof that (D10) is weakly (\mathcal{F}, k) -boundary reducible, but not (\mathcal{F}, k) -boundary reducible. Let R be a subgraph of a graph G defined by configuration (D10). Let a, b, c and d be vertices of R . The initial list sizes of a list assignment L as defined by the function $(4 - (\deg_G + \deg_R))$ are given in Fig. 8.

First we will show that we cannot fix the color of a and still properly color R . Indeed, if the lists of b and c are identical and both contained the color assigned to a , there would be no proper L -coloring of R .

That being said, if we fix the color of any other vertex in R , then we will still be able to properly L -color R . Therefore, we can only apply (FIX) to a subset of the vertices of R . Given the graphs in \mathcal{F} , it immediately follows that the graph $H = R$ is weakly (\mathcal{F}, k) -boundary reducible, but as we have show, it is not (\mathcal{F}, k) -boundary reducible.

In the enhanced version, the main trick is that we never need to check (FORB) on two adjacent vertices. We do that by allowing (FIX) only on vertices, where their external neighbors are non-adjacent. The easiest way to do so is to use (FIX) only on vertices that have at most 1 external neighbor. In case of (D10), the only option for (FIX) is the vertex d . \square

² This program is also available as a part of the sources in our arXiv submission [9] (file `reducible_configurations.sage`).

The above works in all cases except (D1) and (D4). As the case (D1) was already discussed, we now justify the usage of the configuration (D4).

Lemma 9 *The configuration (D4) is weakly $(\mathcal{F}, 4)$ -boundary-reducible and for all vertices x in its reducible part holds that $|N(x) \cap R_j| \leq 1$ or $N(x) \cap R_j$ is a loose set, where R_j is a reducible part of some other configuration.*

Proof The check of the reducibility is straightforward.

The only vertices adjacent to the vertices in the reducible part are the two vertices in the boundary which are non-adjacent as $K_4 \in \mathcal{F}$. Therefore, it is sufficient to check non-adjacent non- \mathcal{F} -forbidding vertices in configurations (D1)–(D12). As was already discussed the only such a non-edge is the non-edge in (D9) which forms a loose set. □

5.2 Discharging Rules

In this section, we prove the following Lemma 10, that makes Theorem 2 a corollary of Lemma 2.

Lemma 10 *Let G be a connected $\{K_4, C_5, C_6, C_7\}$ -free plane graph. Then G contains at least one of the reducible configurations (D1)–(D12).*

Proof Assume for contradiction that G is a $\{K_4, C_5, C_6, C_7\}$ -free plane graph with no (D1)–(D12). We will use discharging to arrive to a contradiction.

For every vertex v assign the initial charge $ch(v) := 2 \deg(v) - 6$, and for every face f assign $ch(f) := \ell(f) - 6$, where $\ell(f)$ is the length of the facial walk around f . By Euler’s formula, the total initial charge of all vertices and faces is -12 . We sequentially apply the following rules (see also Fig. 9) that transfer charge. The charge after applying all the rules is called the *final charge*. We will show that the final charge is nonnegative for every vertex and every face, which is a contradiction with the total sum of all charges being -12 .

- (R1) Every 8^+ -face sends charge $\frac{1}{4}$ to every incident 3-face and 4-face for every edge they have in common.
- (R2) For every 3-vertex v that is incident to a triangle t and an edge uv that is not part of any triangle, the following applies. The two faces³ that are incident to uv , each send the following charge to t :

(R2a) $\frac{1}{8}$ if $\deg(u) = 3$,

(R2b) $\frac{1}{4}$ if $\deg(u) \geq 4$.

³ May be the same face twice if uv is a bridge.

- (R3) Every 4-vertex sends charge 1 to every 3-face and 4-face adjacent to it.
- (R4) Every 5-vertex sends charge 1 to every 4-face adjacent to it.
- (R5) Every 5-vertex that is a middle vertex in $Dia(5 - 3, 4, 4)$, $Dia(5 - 3, 3, 5^+)$, or $Dia(5 - 5, 3, 3)$ sends charge $\frac{3}{2}$ to every 3-face of such diamond.
- (R6) Every 5-vertex v , where rule (R5) does not apply, sends charge 1 to every 3-face of a diamond having v as a middle vertex.
- (R7) For every 3-face $f = \{v, u, w\}$ and 5-vertex v such that f is not part of a diamond having v as a middle vertex, the following applies.
- (R7a) If both $\deg(u) \geq 4$ and $\deg(w) \geq 4$, then v sends charge 1 to f .
- (R7b) Otherwise v sends charge 2 to f .
- (R8) Every 6^+ -vertex v sends charge 1 to every 4-face adjacent to it, and 2 to every 3-face f adjacent to it, unless f is part of a diamond having v as a middle vertex.
- (R9) Every 6-vertex v sends charge $\frac{7}{4}$ to every 3-face of every $Dia(6 - 3, 3, 4)$ that contains v .
- (R10) Every 6-vertex v sends charge $\frac{3}{2}$ to every 3-face of $Dia(6 - 3, 4, 4)$ that contains v .
- (R11) Every 6-vertex v sends charge $\frac{5}{4}$ to every 3-face of any diamond d having v as a middle vertex, where (R9) and (R10) did not apply.
- (R12) Every 7^+ -vertex v sends charge $\frac{7}{4}$ to every 3-face of any diamond having v as a middle vertex.
- (R13) For every two 3-faces f, g that form a diamond, if g has positive charge while f has negative charge, then g gives f all its positive charge.

Claim 12 *The final charge of every vertex is nonnegative.*

Proof There are no vertices of degree less than 3, by (D1). The initial charge of a 3-vertex is 0, and this does not change in the discharging process. A 4-vertex v has initial charge 2. It can be adjacent to at most two 4-faces, or otherwise a cycle C_k with $5 \leq k \leq 7$ is created. Therefore (R3) applies on v at most twice and no other rules apply. Hence v has a nonnegative final charge.

Let v be a 5-vertex that is not a middle vertex of a diamond. Note that v can be adjacent to at most two faces of size at most 4, or otherwise a cycle C_k with $5 \leq k \leq 7$ is created. Thus, the initial charge of v is 4, and (R4) and (R7) are applied together at most twice, implying that v has nonnegative final charge.

Let v be a 5-vertex that is a middle vertex of a diamond d . Then v is adjacent to at most one more face f of size at most 4, and f does not share any edge with d , or otherwise a cycle C_k with $5 \leq k \leq 7$ is created. If (R5) does not apply to v , then by (R4), (R6) and (R7), v sends 1 to each of the two 3-faces in d and at most 2 to f , leaving v with final nonnegative charge. Suppose (R5), where v sends charge 3 to the faces in d , applies to v . If v sends charge of at most 1 to f , then it has final nonnegative charge. So by (R4) and (R7a) we may assume that f is a triangle $\{v, u, w\}$ with $d(u) = 3$ (and $d(w) \leq 4$). See Fig. 10 for an illustration. But then G contains (D7),




Fig. 9 Discharging rules described in Lemma 10 used for Theorem 2

(D11), or (D8) as d is $Dia(5 - 3, 4, 4)$, $Dia(5 - 3, 3, 5^+)$, or $Dia(5 - 5, 3, 3)$, respectively. Hence (R7b) does not apply to v and the final charge is nonnegative.

Let v be a 6-vertex that is the middle vertex of k diamonds and it is adjacent to m faces of size 3 or 4 that are not part of a diamond in which v is a middle vertex. By Lemma 4, $6 \geq 3k + 2m$. Recall that $ch(v) = 6$. Suppose $k = 2$, then $m = 0$. By (D12) and (D3), (R9) cannot apply twice and (R9) cannot apply at the same time as (R10). Then by (R9)–(R11), the final charge of v is at least $6 - 3.5 - 2.5 = 0$ or




Fig. 10 Three options in a subcase of Claim 12 where v is a 5-vertex that is a middle vertex of a diamond

$6 - 3 - 3 = 0$. If $k = 1$ and $m \leq 1$, then by (R8)–(R11), the final charge of v is at least $6 - 3.5 - 2 > 0$. Finally, if $k = 0$ and $m \leq 3$ then by (R8), the final charge of v is at least $6 - 3 \cdot 2 = 0$.

Let v be a 7^+ -vertex that is the middle vertex of k distinct diamonds, and v is adjacent to m faces of sizes 3 and 4 that are not part of a diamond in which v is a middle vertex. Then by Lemma 4, $\deg(v) \geq 3k + 2m$. By (R12) v sends total weight of $3.5k$ to the k diamonds in which v is a middle vertex, and by (R8) it sends at most $2m$ to the other faces of size at most 4 it is adjacent to.

Altogether, the final charge of v is at least

$$\begin{aligned} 2 \deg(v) - 6 - 3.5k - 2m &= 2 \deg(v) - 6 - (3k + 2m) - k/2 \\ &\geq \deg(v) - 6 - \frac{1}{2} \cdot \left\lfloor \frac{\deg(v)}{3} \right\rfloor, \end{aligned}$$

where the last inequality follows from Lemma 4, and $\deg(v) - 6 - \frac{1}{2} \cdot \left\lfloor \frac{\deg(v)}{3} \right\rfloor \geq 0$ whenever $\deg(v) \geq 7$. \diamond

Claim 13 *The final charge of every face that is not contained in a diamond is nonnegative.*

Proof By (R1) and (R2), an 8^+ -face f sends out a total charge of at most $\frac{\ell(f)}{4}$. Thus the final charge of f is at least $\ell(f) - 6 - \frac{\ell(f)}{4} = \frac{3\ell(f)}{4} - 6$ which is nonnegative if $\ell(f) \geq 8$.

Let f be a 3-face that is not part of any diamond. Then the faces sharing an edge with f must be of size at least 8, since otherwise one of them is of size at most 4,




Fig. 11 Three possible triangles distinguished within Claim 13 that depicts Cases (1), (2), and (3), from left to right

which forces a diamond or a cycle C_i with $5 \leq i \leq 7$ together with f . Hence (R1) applies three times with f and f has charge $-3 + \frac{3}{4} = -2.25$ after (R1).

By (D1) and (D2), one of the following holds (see Fig. 11):

- (1) f is $T(3, 3, 5^+)$, or
- (2) f is $T(3, 4^+, 4^+)$, or
- (3) f is $T(4^+, 4^+, 4^+)$.

In case (1), (R2a) applies twice giving charge $\frac{4}{8}$ to f . In addition, (R7b) or (R8) applies and the final charge of f is at least $-3 + \frac{3}{4} + \frac{1}{2} + 2 \geq 0$.

In case (2), (R2) applies once, giving charge $\frac{2}{8}$ to f . Rules (R3), (R7) and (R8) apply together twice with f , each time f receives charge at least 1, and thus the final charge of f is at least $-3 + \frac{3}{4} + \frac{1}{4} + 2 \geq 0$.

In case (3), rules (R3), (R7), and (R8) together apply three times to f and thus the final charge of f is at least $-3 + \frac{3}{4} + 3 > 0$.

If f is a 4-face, the faces sharing an edge with f must be of size at least 8, since otherwise one of them is of size at most 4, which forces a cycle C_i of size $5 \leq i \leq 7$ with f . Hence (R1) applies four times with f contributing charge $\frac{4}{4}$. By (D9) f has at least two 4^+ -vertices. Thus at least one of (R3), (R4), and (R8) applies to f , giving charge 1 to f . Hence the final charge of f is at least $-2 + 1 + 1 \geq 0$. ◊

Claim 14 *The final charge of every 3-face that is contained in a diamond is nonnegative.*

Proof In the light of (R13), we will consider the faces that form a diamond together in pairs and show that as a pair, they receive sufficient charge. Let f and g be 3-faces sharing an edge, i.e., they form a diamond. Observe that in this case the other faces sharing edges with f and g must be of size at least 8, for otherwise one of them is of size at most 4, which forces a cycle C_i of size $5 \leq i \leq 7$ with f and g . Therefore, (R1) applies twice to each f and g resulting in charge ch' such that $ch'(f) = ch'(g) = -3 + \frac{2}{4} = -2.5$. Hence we aim to show that f and g together receive at least 5 more charge. We denote the vertices of f by u, v, x where u, v are shared with g , and by y the third vertex of g .

By symmetry, we assume that $\deg(u) \geq \deg(v)$ and $\deg(y) \geq \deg(x)$. Note that by (D4), $\deg(u) \geq 4$ and by (D1) the degree of each of the other vertices is at least 3.

We split the analysis into cases based on the type of diamond f and g form.

- $Dia(4 - 3, \star, \star)$
By (D2) and (D6), $\deg(x) \geq 4$ and $\deg(y) \geq 5$. Hence we are in case $Dia(4 - 3, 4^+, 5^+)$. For u , (R3) applies twice, for x one of (R3), (R7b), or (R8) applies, and for y one of (R7b), or (R8) applies. Thus the charge f and g receive using these rules is at least $3 \cdot 1 + 2 = 5$. Hence the final charges of f and g are nonnegative.
- $Dia(5 - 3, 3, 3)$, $Dia(5 - 3, 3, 4)$
Reducible by (D9) and (D10).
- $Dia(5 - 3, 3, 5^+)$
In this case, (R5) applies to u and (R7b) or (R8) applies to y . This gives charge $2 \cdot 1.5 + 2 = 5$. Hence the final charges of f and g are nonnegative.
- $Dia(5 - 3, 4, 4)$
In this case, (R5) applies to u . In addition (R3) applies to both x and y . This gives charge $2 \cdot 1.5 + 1 + 1 = 5$. Hence the final charges of f and g are nonnegative.
- $Dia(5 - 3, 4^+, 5^+)$
In this case, (R6) applies to u . In addition (R3), (R7b), or (R8) applies to x and (R7b), or (R8) to y . This gives charge at least $2 \cdot 1 + 1 + 2 = 5$. Hence the final charges of f and g are nonnegative.
- $Dia(6 - 3, 3, 3)$
Reducible by (D9).
- $Dia(6 - 3, 3, 4)$
By (R9), u contributes charge 3.5 and by (R4), y contributes charge 1. Let z be a neighbor of x that is not u or v . By (D9), $\deg(z) \geq 4$ Hence the application of (R2b) around x contributes charge 1/2. This gives charge $3.5 + 1 + 0.5 = 5$ in total. Hence the final charges of f and g are nonnegative.
- $Dia(6 - 3, 3, 5^+)$
By (R11), u contributes charge 2.5 and by (R7b) or (R8), y contributes charge 2. Let z be a neighbor of x that is not u or v . By (D9), $\deg(z) \geq 4$ Hence the application of (R2) around x contributes charge 1/2. This gives charge $2.5 + 2 + 0.5 = 5$ in total. Hence the final charges of f and g are nonnegative.
- $Dia(6 - 3, 4, 4)$
By (R10), u contributes charge 3 and by (R3), x and y each contribute charge 1. This gives charge $3 + 1 + 1 = 5$ in total. Hence the final charges of f and g are nonnegative.
- $Dia(6 - 3, 4^+, 5^+)$
By (R11), u contributes charge 2.5, by (R3), (R7b) or (R8), x contributes charge at least 1, and by (R7b) or (R8), y contributes charge 2. This gives charge at least $2.5 + 1 + 2 = 5$ in total. Hence the final charges of f and g are nonnegative.
- $Dia(7^+ - 3, 3, 3)$
Reducible by (D9).

- $Dia(7^+ - 3, 3, 4^+)$
 By (R12), u contributes charge 3.5, by (R3), (R7b) or (R8), y contributes charge at least 1. Let z be the neighbor of x that is not u or v . By (D9), $\deg(z) \geq 4$ Hence the application of (R2b) around x contributes charge 1/2. This gives charge at least $3.5 + 1 + 0.5 = 5$ in total. Hence the final charges of f and g are nonnegative.
- $Dia(7^+ - 3, 4^+, 4^+)$
 By (R12), u contributes charge 3.5, by (R3), (R7b) or (R8), x and y each contribute charge at least 1. This gives charge at least $3.5 + 1 + 1 = 5.5$ in total. Hence the final charges of f and g are nonnegative.
- $Dia(4 - 4, \star, \star)$ and $Dia(5 - 4, \star, \star)$
 By (D5), $\deg(y) \geq 4$. By (R3) or (R6), u and v together contribute charge 4, by (R3), y each contributes charge 1. This gives charge at least $4 + 1 = 5$ in total. Hence the final charges of f and g are nonnegative.
- $Dia(6^+ - 4^+, \star, \star)$
 By (R3), (R6), (R11), and (R12), u and v together contribute charge at least $1.25 + 1.25 + 1 + 1$. If (R3), (R7), or (R8) applies to y , then the total charge is at least 5.5. Hence we can assume the case $Dia(6^+ - 4^+, 3, 3)$. Then (R2a) or (R2b) applies at each x and y and the total contribution is at least 0.5. This gives charge at least $4.5 + 0.5 = 5$ in total. Hence the final charges of f and g are nonnegative.
- $Dia(5 - 5, 3, 3)$
 By (R5), u and v together contribute charge at least $4 \times 1.5 = 6$. Hence the final charges of f and g are nonnegative.
- $Dia(5 - 5, 3^+, 4^+)$
 By (R6), u and v together contribute charge at least $4 \times 1 = 4$. By (R3), (R7a), or (R8), y contributes charge at least 1. This gives charge at least $4 + 1 = 5$ in total. Hence the final charges of f and g are nonnegative.

This concludes the proof of Claim 14. ◇

Since all final charges are nonnegative, this concludes the proof of Lemma 10. □

Acknowledgements The work was initiated at Dagstuhl Seminar on “Graph Colouring: from Structure to Algorithms”(19271). Part of the work was carried out when T. Masařík was visiting Iowa State University. He thanks Steve Butler for kind hospitality. We would like to thank Ilkyoo Choi for fruitful discussions at the beginning of this project.

Funding B. Lidický was supported in part by NSF grant DMS-1855653. T. Masařík received funding from the European Research Council (ERC) under the European Union’s Horizon 2020 research and innovation programme Grant Agreement 714704. He completed a part of this work while he was a post-doc at Simon Fraser University in Canada. S. Zerbib was supported by NSF grant DMS-1953929.



Declarations

Conflict of interest The authors have no relevant financial or non-financial interests to disclose.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

1. Bradshaw, P., Masařík, T., Stacho, L.: Flexible list colorings in graphs with special degeneracy conditions. *J. Graph Theory* (2022). <https://doi.org/10.1002/jgt.22849>
2. Choi, I., Clemen, F.C., Ferrara, M., Horn, P., Ma, F., Masařík, T.: Flexibility of planar graphs—sharpening the tools to get lists of size four. *Discrete Appl. Math.* **306**, 120–132 (2022). <https://doi.org/10.1016/j.dam.2021.09.021>
3. Cranston, D.W., West, D.B.: An introduction to the discharging method via graph coloring. *Discrete Math.* **340**(4), 766–793 (2017). <https://doi.org/10.1016/j.disc.2016.11.022>
4. Dvořák, Z., Masařík, T., Musílek, J., Pangrác, O.: Flexibility of planar graphs of girth at least six. *J. Graph Theory* **95**(3), 457–466 (2020). <https://doi.org/10.1002/jgt.22567>
5. Dvořák, Z., Masařík, T., Musílek, J., Pangrác, O.: Flexibility of triangle-free planar graphs. *J. Graph Theory* **96**(4), 619–641 (2021). <https://doi.org/10.1002/jgt.22634>
6. Dvořák, Z., Norin, S., Postle, L.: List coloring with requests. *J. Graph Theory* **92**(3), 191–206 (2019). <https://doi.org/10.1002/jgt.22447>
7. Fijavž, G., Juvan, M., Mohar, B., Škrekovski, R.: Planar graphs without cycles of specific lengths. *Eur. J. Combin.* **23**(4), 377–388 (2002). <https://doi.org/10.1006/eujc.2002.0570>
8. Lam, P.C.B., Baogang, X., Liu, J.: The 4-choosability of plane graphs without 4-cycles. *J. Combin. Theory Ser. B* **76**(1), 117–126 (1999). <https://doi.org/10.1006/jctb.1998.1893>
9. Lidický, B., Masařík, T., Murphy, K., Zerbib, S.: On weak flexibility in planar graphs. *arXiv preprint* (2020). <https://doi.org/10.48550/ARXIV.2009.07932>
10. Masařík, T.: Flexibility of planar graphs without 4-cycles. *Acta Math. Univ. Comen.* **88**(3), 935–940 (2019). <http://www.iam.fmph.uniba.sk/amuc/ojs/index.php/amuc/article/view/1182>
11. Wang, W., Lih, K.-W.: Choosability and edge choosability of planar graphs without five cycles. *Appl. Math. Lett.* **15**(5), 561–565 (2002). [https://doi.org/10.1016/s0893-9659\(02\)80007-6](https://doi.org/10.1016/s0893-9659(02)80007-6)
12. Yang, D., Yang, F.: Flexibility of planar graphs without C_4 and C_5 . *arXiv preprint* (2020). <https://doi.org/10.48550/ARXIV.2006.05243>

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Authors and Affiliations

Bernard Lidický¹  · **Tomáš Masařík^{2,3}**  · **Kyle Murphy^{1,4}** · **Shira Zerbib¹** 

Bernard Lidický
lidicky@iastate.edu

Kyle Murphy
kyle.murphy@dsu.edu

Shira Zerbib
zerbib@iastate.edu

- ¹ Department of Mathematics, Iowa State University, Ames, IA, USA
- ² Faculty of Mathematics, Informatics and Mechanics, University of Warsaw, Warsaw, Poland
- ³ Department of Mathematics, Simon Fraser University, Burnaby, BC, Canada
- ⁴ Present Address: Dakota State University in Madison, Madison, SD, USA