Preface to Special Topic on Atmospheric Greenhouse Gas Measurement and Application in China

Pengfei HAN¹, Ning ZENG², Bo YAO³, Weijian ZHOU⁴, Liqi CHEN⁵, Shaoqiang WANG⁶, Honggang LV⁷, Wei XIAO⁸, Lingyun ZHU⁹, and Jiaping XU¹⁰

¹State Key Laboratory of Numerical Modeling for Atmospheric Sciences and Geophysical Fluid Dynamics, Institute of Atmospheric Physics, Chinese Academy of Sciences, Beijing 100029, China

²Department of Atmospheric and Oceanic Science, University of Maryland, College Park, MD 20742, USA

³Meteorological Observation Centre of China Meteorological Administration, Beijing 100081, China

⁴Institute of Earth Environment, Chinese Academy of Sciences, Xi'an 710061, China

⁵Third Institute of Oceanography, Ministry of Natural Resources, Xiamen 361005, China

⁶Institute of Geographic Sciences and Natural Resources Research, Chinese Academy of Sciences, Beijing 100101, China

⁷National Marine Environment Forecasting Center, Beijing 100081, China

⁸Nanjing University of Information Science & Technology, Nanjing 210044, China

⁹Shanxi Institute of Meteorological Sciences, Taiyuan 030002, China

¹⁰Jiangsu Institute of Meteorological Sciences, Nanjing 210009, China

Citation: Han, P. F., and Coauthors, 2020: Preface to Special Topic on Atmospheric Greenhouse Gas Measurement and Application in China. *Adv. Atmos. Sci.*, **37**(6), 555–556, <https://doi.org/10.1007/s00376-020-9300-x>.

China initiated a national carbon trading market in December 2017. Commitments and actions to reduce greenhouse gas (GHG) emissions require consistent, reliable and timely information on GHG emissions. GHG monitoring and modeling studies provide GHG emission estimates to evaluate and guide progress towards emission reductions. GHG monitoring has mainly focused on global-scale background networks over the last few decades, while recent efforts have been made on regional and urban scales, such as projects in the Beijing–Tianjin–Hebei city cluster, in Paris, Washington–Baltimore, Indianapolis, and Los Angeles. These “top-down” atmospheric observations, together with “bottom-up” GHG emissions inventories and biosphere fluxes, are integrated into GHG monitoring and application information systems. The Integrated Global Greenhouse Gas Information System (or IG³IS) was established by WMO/UNEP to facilitate such efforts after the 17th World Meteorological Congress. As one of the countries heavily impacted by climate change, China has implemented many actions to respond to climate change. Atmospheric GHG measurement techniques have been developed rapidly in China over the last decade, and this work has been applied in evaluating GHG emissions, identifying GHG sources, quantifying GHG fluxes, and supporting GHG-related policymaking.

The First China Greenhouse Gas Monitoring Conference was held in Beijing on 30th and 31st May 2019, which attracted more than 200 people from over 80 participating universities and institutes. This AAS–AOSL–ACCR joint special issue on “*Atmospheric Greenhouse Gas Measurement and Application in China*” provides a selection of research highlights from this conference. The selected papers cover all the sections of the conference, as follows:

(1) CO₂ background monitoring. An example here is the paper by [Mai et al. \(2020\)](#), which measured background atmospheric CO₂ and the potential source regions in the Pearl River Delta region.

(2) CO₂ city area (emission sources) monitoring. The paper by [Xiong et al. \(2020\)](#) reports on a two-year observation of fossil-fuel CO₂ spatial distribution in Xi'an City using ¹⁴C isotope data. Also, [Bao et al. \(2020\)](#) present a novel study in which a low-cost non-dispersive infrared sensor was used to observe the vertical CO₂ distribution in a high emissions area of Shijiazhuang, Hebei Province. Meanwhile, [Zhang et al. \(2020\)](#) conducted a three-year analysis of the variations and transmission characteristics of CO₂ in Taiyuan, Shanxi Province.

(3) Non-CO₂ GHG monitoring. [Pu et al. \(2020\)](#) estimated the hydrofluorocarbon emissions during 2012–16 in the Yangtze River Delta region based on weekly sampling at Lin'an Background Station, Zhejiang Province.

(4) GHG flux measurement. An example here is the work presented by [Zhu et al. \(2020\)](#), which compared the ozone

fluxes over a maize field measured with gradient methods and the eddy covariance technique.

(5) New technologies and applications for GHG measurement. For instance, [Ji et al. \(2020\)](#) show the temporal and vertical distribution of methane in Xianghe derived from ground-based Fourier transform infrared and Picarro measurements. Using the Aircore method, [Yi et al. \(2020\)](#) conducted direct observations of atmospheric transport and stratosphere–troposphere exchange from CO₂ profile measurements in Inner Mongolia.

(6) Applications of observational data. [Fu et al. \(2020\)](#) used a global chemical transport model (GEOS-Chem) and observations at 14 surface stations from the World Data Centre for Greenhouse Gases to study the sensitivity of the simulated CO₂ concentration to interannual variations of its sources and sinks over East Asia.

These efforts provide accurate scientific understanding, technical support and solutions for major issues such as energy conservation and emissions reduction, carbon market transactions and low-carbon development. Along with this special issue, the China Carbon Monitoring Alliance (<http://cma.sense1.cn/>) is under way, and a data resources platform has also been established to provide opportunities for collaborations.

REFERENCES

- Bao, Z. X., and Coauthors, 2020: Observation and modeling of vertical carbon dioxide distribution in a heavily polluted suburban environment. *Atmospheric and Oceanic Science Letters*, **13**, 4, <https://doi.org/10.1080/16742834.2020.1746627>.
- Fu, Y., H. Liao, X. J. Tian, H. Gao, Z. N. Cai, and R. Han, 2020: Sensitivity of the simulated CO₂ concentration to inter-annual variations of CO₂ sources and sinks over East Asia. *Advances in Climate Change Research*, <https://doi.org/10.1016/j.accres.2020.03.001>.
- Ji, D. H., and Coauthors, 2020: Deriving temporal and vertical distributions of methane in Xianghe using ground-based Fourier transform infrared and gas-analyzer measurements. *Adv. Atmos. Sci.*, **37**(6), <https://doi.org/10.1007/s00376-020-9233-4>.
- Mai, B. R., X. J. Deng, F. Zhang, H. He, T. Luan, F. Li, and X. Liu, 2020: Background characteristics of atmospheric CO₂ and the potential source regions in the Pearl River Delta region of China. *Adv. Atmos. Sci.*, **37**(6), <https://doi.org/10.1007/s00376-020-9238-z>.
- Pu, J. B., H. H., Xu, B. Yao, Y. Yu, Y. J. Jiang, Q. L. Ma, and L. Q. Chen, 2020: Estimate of hydrofluorocarbon emissions for 2012–2016 in the Yangtze River Delta, China. *Adv. Atmos. Sci.*, **37**(6), <https://doi.org/10.1007/s00376-020-9242-3>.
- Xiong, X. H., and Coauthors, 2020: Two-year observation of fossil fuel carbon dioxide spatial distribution in Xi'an City. *Adv. Atmos. Sci.*, <https://doi.org/10.1007/s00376-020-9241-4>.
- Yi, Y., and Coauthors, 2020: Direct observations of atmospheric transport and stratosphere–troposphere exchange from high-precision carbon dioxide and carbon monoxide profile measurements. *Adv. Atmos. Sci.*, **37**(6), <https://doi.org/10.1007/s00376-020-9227-2>.
- Zhang, F. S., L. Y. Zhu, S. M. Yan, X. A. Gao, and K. N. Pei, 2020: Analysis of the carbon dioxide mole fraction variation and its transmission characteristics in Taiyuan. *Atmospheric and Oceanic Science Letters*, <https://doi.org/10.1080/16742834.2020.1737501>.
- Zhu, Z. L., X. Z. Tang, and F. H. Zhao, 2020: Comparison of ozone fluxes over a maize field measured with gradient methods and the eddy covariance technique. *Adv. Atmos. Sci.*, **37**(6), <https://doi.org/10.1007/s00376-020-9217-4>.