CORRECTION



## Correction to: Association between discordance of LDL-C and non-HDL-C and clinical outcomes in patients with stent implantation: from the FU-Registry

Michiyo Shiiba<sup>1,2</sup> · Bo Zhang<sup>3</sup> · Shin-ichiro Miura<sup>2,4</sup> · Amane Ike<sup>2</sup> · Daisuke Nose<sup>2,5</sup> · Takashi Kuwano<sup>2</sup> · Satoshi Imaizumi<sup>2</sup> · Makoto Sugihara<sup>2</sup> · Atushi Iwata<sup>2</sup> · Hiroaki Nishikawa<sup>2</sup> · Akira Kawamura<sup>2</sup> · Kazuyuki Shirai<sup>5,6</sup> · Shin'ichiro Yasunaga<sup>3</sup> · Keijiro Saku<sup>2,4</sup>

Published online: 19 February 2018 © Springer Japan KK, part of Springer Nature 2018

## Correction to: Heart Vessels (2018) 33:102–112 https://doi.org/10.1007/s00380-017-1036-x

Unfortunately, the Fig. 1 was published incorrectly in the original publication of the article. The correct figure is as below.

In addition, under the Discussion section, the following sentence from the last paragraph was published incorrectly as, "This study has several ... this study was not a prospective cohort study...". The correct sentence is as follows, "This study has several ... this study was a prospective cohort study...".

The original article can be found online at https://doi.org/10.1007/s00380-017-1036-x.

Shin-ichiro Miura miuras@cis.fukuoka-u.ac.jp

- Keijiro Saku saku-k@fukuoka-u.ac.jp
- <sup>1</sup> Fukuoka Jo Gakuin University, Fukuoka 811-3113, Japan
- <sup>2</sup> Department of Cardiology, Fukuoka University School of Medicine, 7-45-1 Nanakuma Jonan-ku, Fukuoka 814-0180, Japan
- <sup>3</sup> Department of Biochemistry, Fukuoka University School of Medicine, Fukuoka 814-0180, Japan
- <sup>4</sup> Department of Molecular Cardiovascular Therapeutics, Fukuoka University School of Medicine, Fukuoka 814-0180, Japan
- <sup>5</sup> Division of Cardiology, Hakujyuji Hospital, Fukuoka 819-8511, Japan
- <sup>6</sup> Department of Cardiovascular Medicine, Fukuoka University Chikushi Hospital, Fukuoka 818-8502, Japan




Fig.1 Flow chart of the study (the FU-Registry). MACEs were defined to include these outcomes: all deaths, MI, TLR-PCI, TLR-CABG, and stent thrombosis