Correction to: Increased prevalence of cerebral microbleeds in patients with low left ventricular systolic function

Tomohiko Watanabe¹ · Yumiko Kanzaki¹ · Yohei Yamauchi¹ · Takahide Ito¹ · Yusuke Nishida¹ · Kenichiro Yamamura² · Tsuyoshi Komori² · Koichi Sohmiya¹ · Masaaki Hoshiga¹

Published online: 4 March 2020
© Springer Japan KK, part of Springer Nature 2020

Correction to: Heart and Vessels (2020) 35:384–390
<https://doi.org/10.1007/s00380-019-01503-0>

In the original publication of the article, under the results section, the following sentence “Of 118 patients with CMBs, an average number of CMBs were 2.6 ± 3.2 , 55 patients (47%) in lobar locations, 84 patients (71%) in deep or infratentorial locations, and 26 patients (22%) in both locations” was published incorrectly.

The correct sentence should read as “Of 118 patients with CMBs, an average number of CMBs were 2.6 ± 3.2 , 58 patients (49%) in lobar locations, 86 patients (73%) in deep or infratentorial locations, and 26 patients (22%) in both locations”.

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

The original article can be found online at <https://doi.org/10.1007/s00380-019-01503-0>.

✉ Yumiko Kanzaki
in3089@osaka-med.ac.jp

¹ Department of Cardiology, Osaka Medical College, 2-7 Daigaku-machi, Takatsuki 569-8686, Japan

² Department of Radiology, Osaka Medical College, Takatsuki, Japan