



Correction to: Clinical outcomes and predictors of restenosis in patients with femoropopliteal artery disease treated using polymer-coated paclitaxel-eluting stents or drug-coated balloons

Naoki Yoshioka^{1,2} · Takahiro Tokuda³ · Akio Koyama⁴ · Takehiro Yamada⁵ · Ryusuke Nishikawa⁶ · Kiyotaka Shimamura⁶ · Kensuke Takagi^{1,2} · Yasuhiro Morita¹ · Akihito Tanaka² · Hideki Ishii² · Itsuro Morishima¹  · Toyoaki Murohara² · on the ASIGARU PAD investigators

Published online: 19 October 2021
© Springer Japan KK, part of Springer Nature 2021

Correction to: Heart and Vessels
<https://doi.org/10.1007/s00380-021-01941-9>

The original article was updated.

In the original publication of the article, in the following sentence "... the multi-center registry data of 151 ...", the term "We assessed" was missed in the third line of abstract section. The correct sentence should read as follows "We assessed the multi-center registry data of 151 lesions from 151 patients who underwent EVT for symptomatic PAD in the superficial femoral and proximal popliteal arteries using PC-PES or DCB. ...".

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

The original article can be found online at <https://doi.org/10.1007/s00380-021-01941-9>.

✉ Itsuro Morishima
morishima-i@muc.biglobe.ne.jp

¹ Department of Cardiology, Ogaki Municipal Hospital, 4-86 Minaminokawa-cho, Ogaki, Gifu, Japan

² Department of Cardiology, Nagoya University Graduate School of Medicine, Nagoya, Japan

³ Department of Cardiology, Nagoya Heart Center, Nagoya, Japan

⁴ Department of Vascular Surgery, Ichinomiya Municipal Hospital, Ichinomiya, Japan

⁵ Department of Cardiology, Kizawa Memorial Hospital, Minokamo, Japan

⁶ Department of Cardiology, Shizuoka General Hospital, Shizuoka, Japan