



Correction: Combined assessment of frailty and nutritional status can be a prognostic indicator after percutaneous coronary intervention

Ryota Noike¹ · Hideo Amano¹ · Shojiro Hirano² · Masakazu Tsubono² · Yoshimasa Kojima² · Yosuke Oka² · Hiroto Aikawa² · Shingo Matsumoto² · Takayuki Yabe² · Takanori Ikeda¹

Published online: 17 December 2022
© Springer Japan KK, part of Springer Nature 2022

Correction: Heart and Vessels

<https://doi.org/10.1007/s00380-022-02176-y>

In the original publication of the article, in Table 1, under Comorbidity, the “Hypertension” values for “High-risk group ($n = 141$)” and “Low-risk group ($n = 267$)” were published incorrectly. The correct Hypertension values for “High-risk group ($n = 141$)” is “97 (68.8)” and for “Low-risk group ($n = 267$)” is “199 (74.5)”.

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

The original article can be found online at <https://doi.org/10.1007/s00380-022-02176-y>.

✉ Hideo Amano
amanohide@med.toho-u.ac.jp

¹ Division of Cardiovascular Medicine, Department of Internal Medicine, Toho University Graduate School of Medicine, 6-11-1 Omorinishi, Ota-ku, Tokyo 143-8541, Japan

² Division of Cardiovascular Medicine, Department of Internal Medicine, Faculty of Medicine, Toho University, Tokyo, Japan