

Hydrocephalus and cysticercosis

Somsri Wiwanitkit · Viroj Wiwanitkit

Received: 20 September 2011 / Accepted: 10 November 2011 / Published online: 24 November 2011
© Springer-Verlag 2011

Dear Editor:

The recent publication on hydrocephalus and cysticercosis is very interesting [1]. We would like to share some ideas on hydrocephalus and cysticercosis. Indeed, this condition is a deadly condition that needs proper management. Similar to Matushita et al.'s note, Sotelo proposed that "The use of one or various therapeutic measures largely depends on the peculiar combination of number, location, and biological stage of lesions as well as the degree of inflammatory response to the parasites [2]." However, to derive success in treatment, awareness and early diagnosis can lead to prompt effective treatment. There are some new alternative techniques for diagnosis to be mentioned. A good example is "Full moon" endoscopic sign which can

help diagnose the case and further minimally invasive therapeutic approach can be done [3].

References

1. Matushita H, Pinto FC, Cardeal DD, Teixeira MJ (2011) Hydrocephalus in neurocysticercosis. *Childs Nerv Syst* 27(10):1709–1721
2. Sotelo J (2011) Clinical manifestations, diagnosis, and treatment of neurocysticercosis. *Curr Neurol Neurosci Rep* 11(6):529–535
3. Ramos-Zúñiga R, de La Cruz-Ramírez J, Casillas-Espinosa PM, Sánchez-Prieto JA, López-Hernández MD (2011) "Full moon" endoscopic sign in intraventricular neurocysticercosis. *Minim Invasive Neurosurg* 54(2):90–94

S. Wiwanitkit (✉) · V. Wiwanitkit
Wiwanitkit House,
Bangkhuae,
Bangkok, Thailand 10160
e-mail: somsriwiwan@hotmail.com