



Correction to: European Society for Paediatric Oncology (SIOPE) MRI guidelines for imaging patients with central nervous system tumours

Shivaram Avula¹ · Andrew Peet^{2,3} · Giovanni Morana⁴ · Paul Morgan⁵ · Monika Warmuth-Metz⁶ · Tim Jaspan⁷ · European Society for Paediatric Oncology (SIOPE)-Brain Tumour Imaging Group

Published online: 19 July 2021

© Springer-Verlag GmbH Germany, part of Springer Nature 2021

Correction to: Child's Nervous System

<https://doi.org/10.1007/s00381-021-05199-4>

The original version of this article unfortunately contained an error. Table 2 went missing in the published version; hence, given here is the missing table.

The original article has been corrected.

The original article can be found online at <https://doi.org/10.1007/s00381-021-05199-4>.

✉ Shivaram Avula
shivaram.avula@alderhey.nhs.uk

¹ Department of Radiology, Alder Hey Children's NHS Foundation Trust, East Prescott Road, Liverpool L14 5AB, UK

² Institute of Cancer and Genomic Sciences, University of Birmingham, Birmingham, UK

³ Birmingham Women's and Children's Hospital NHS Foundation Trust, Birmingham, UK

⁴ Department of Neurosciences, University of Turin, Turin, Italy

⁵ Department of Medical Physics, Nottingham University Hospitals, Nottingham, UK

⁶ Institute of Diagnostic and Interventional Neuroradiology, University of Würzburg, Würzburg, Germany

⁷ Department of Radiology, Nottingham University Hospitals, Nottingham, UK

Table 2 Spine imaging

Essential Sequences*			
Sequence	Technique	Parameter	Plane
T ₁ W + Contrast	2D SE/TSE	Slice thickness ≤ 3 mm Slice gap < 0.5 mm	Sagittal whole spine (entire dural sac)
T ₁ W + Contrast	2D SE/TSE or 3D gradient	Slice thickness 4–5 mm No slice gap	Axial–suspicious areas*
Sequence	Technique		Plane
Matrix size—Minimum 256 (512 is desirable for better resolution)			
Optional Sequences			
T ₂ W		2D SE/TSE	Sagittal whole spine
T ₂ W		2D SE/TSE	Axial-suspicious areas
Heavily weighted T ₂ W		2D or 3D CISS/B FFE/FIESTA**	Sagittal ± axial

*In primary tumours of the spinal cord, T₂ W and pre contrast T₁ W sequences are essential

** The heavily weighted T₂ W sequence localised to a region of interest is useful in assessment of drop metastasis

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.