Correction to: 500-year tree-ring reconstruction of Salween River streamflow related to the history of water supply in Southeast Asia

Feng Chen^{1,2} · Huaming Shang² · Irina Panyushkina³ · David Meko³ · Jinbao Li⁴ · Yujiang Yuan² · Shulong Yu² · Fahu Chen⁵ · Daming He¹ · Xian Luo¹

Published online: 18 September 2019
© Springer-Verlag GmbH Germany, part of Springer Nature 2019

Correction to: *Climate Dynamics*
<https://doi.org/10.1007/s00382-019-04948-1>

In the original publication of the article Fig. 1 was incorrect. The correct Fig. 1 is given below:

Amendments of Fig. 1 have been made to the non-scientific content of the article.

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

The original article can be found online at <https://doi.org/10.1007/s00382-019-04948-1>.

✉ Feng Chen
feng653@163.com

✉ Fahu Chen
fhchen@itpcas.ac.cn

- ¹ Yunnan Key Laboratory of International Rivers and Transboundary Eco-Security, Institute of International Rivers and Eco-Security, Yunnan University, Kunming, China
- ² Key Laboratory of Tree-ring Physical and Chemical Research of China Meteorological Administration/Xinjiang Laboratory of Tree-ring Ecology, Institute of Desert Meteorology, China Meteorological Administration, Urumqi, China
- ³ Laboratory of Tree-Ring Research, University of Arizona, Tucson, Arizona, USA
- ⁴ Department of Geography, University of Hong Kong, Hong Kong, China
- ⁵ Institute of Tibetan Plateau Research, Chinese Academy of Sciences (CAS), Beijing, China

Fig. 1 Study region and data points. **a** Map of cross-border region of the Salween River Basin showing basin boundary and locations of the tree-ring sites, weather and hydrological stations. Table 1 lists the names of sites and stations. **b** Map of Southeast Asia showing major

ivers originating on the Tibetan Plateau. The numbers 1 and 2 denote the locations of MHS (Buckley et al. 2007) and BDNP (Buckley et al. 2010) sampling points