



Correction to: Cold-Eurasia contributes to arctic warm anomalies

Bingyi Wu^{1,2} · Shuoyi Ding^{1,2}

Published online: 1 November 2022
© Springer-Verlag GmbH Germany, part of Springer Nature 2022

Correction to: Climate Dynamics
<https://doi.org/10.1007/s00382-022-06445-4>

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Table 1 was missing from this article; the Table 1 should have appeared as shown below.

The original article has been corrected.

Table 1 Description for sensitivity experiments with different models

Experiment name	Prescribed forcing region	Prescribed ground albedo	Integration time
CAM4	50°–60°N, 50°–100°E	0.8	60 years
CAM5_1	50°–60°N, 60°–110°E	0.85	60 years
CAM5_2	40°–50°N, 60°–110°E	0.8	60 years
WACCM	50°–60°N, 50°–100°E	0.8	60 years

The original article can be found online at <https://doi.org/10.1007/s00382-022-06445-4>.

✉ Shuoyi Ding
dsy@fudan.edu.cn

¹ Department of Atmospheric and Oceanic Sciences,
and Institute of Atmospheric Sciences, Fudan University,
200438 Shanghai, China

² CMA-FDU Joint Laboratory of Marine Meteorology,
200438 Shanghai, China