

Roberto Zinicola  
Matthew D. Rutter  
Giuliano Falasco  
Jim C. Brooker  
Vincenzo Cennamo  
Sandro Contini  
Brian P. Saunders

## Haemorrhagic radiation proctitis: endoscopic severity may be useful to guide therapy

---

Published online: 12 March 2004  
© Springer-Verlag 2004

---

The online version of the original article can be found at <http://dx.doi.org/10.1007/s00384-003-0487-y>

---

R. Zinicola · M. D. Rutter · G. Falasco · J. C. Brooker ·  
B. P. Saunders (✉)  
Wolfson Unit for Endoscopy,  
St. Mark's Hospital,  
Watford Road, Harrow, HA1 3UJ, UK  
e-mail: b.saunders@ic.ac.uk  
Tel.: +44-208-2354225  
Fax: +44-208-4233588

V. Cennamo · S. Contini  
Endoscopia Digestiva,  
Ospedale Bellaria,  
Bologna, Italy

---

### Int J Colorectal Dis (2003) 18:439–444

In “Materials and methods,” an item (“— Fresh blood: 1 point”) was inadvertently omitted from the displayed list that presents the scoring of factors. The list should read:

- Distribution of telangiectasias
- Distal rectum (within 10 cm from anal verge): 1 point
- Entire rectum +/-sigmoid (more than 10 cm from anal verge): 2 points
  
- Surface area covered by telangiectasias
- Less than 50%: 1 point
- More than 50%: 2 points
  
- Presence of fresh blood
- No fresh blood: 0 points
- Fresh blood: 1 point