Correction to: Lateral lymph node dissection in advanced low rectal cancer treatment

Jie Li¹ · Akio Shiomi²

Published online: 22 September 2021

© Springer-Verlag GmbH Germany, part of Springer Nature 2021

Correction to: International Journal of Colorectal Disease
<https://doi.org/10.1007/s00384-021-03975-x>

In the original published version of this article, the figure legend of Fig. 1 was changed to clarify its originality.

Fig. 1 Lymphatic drainage system and LLNs in low rectal cancer **A** Lymphatic drainage in low rectal cancer includes two lymphatic flows; upward mesenteric lymphatic drainage, and lateral extra-mesenteric lymphatic drainage along internal iliac vessels (As shown in the blue dotted box). **B** The anatomical regions of LLNs in low rectal cancer

The new legend is as follows:

Figure 1B was adapted and modified from Yamaguchi et al. [65].

Reference

65. Yamaguchi T, Kinugasa Y, Shiomi A et al (2016) Robotic-assisted laparoscopic versus open lateral lymph node dissection for advanced lower rectal cancer. *Surg Endosc* 30:721–728. <https://doi.org/10.1007/s00464-015-4266-y>

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

The original article can be found online at <https://doi.org/10.1007/s00384-021-03975-x>.

✉ Akio Shiomi
a.shiomi@scchr.jp

¹ Department of General Surgery, the Second Affiliated Hospital of Xi'an Jiaotong University, West 5 Road, Xi'an 710004, China

² Division of Colon and Rectal Surgery, Shizuoka Cancer Center Hospital, 1007 Shimonagakubo, Nagaizumi-cho, Sunto-gun, Shizuoka 411-8777, Japan