



## Correction to: Evaluation of the efficacy of *Lactobacillus plantarum* HEAL9 and *Lactobacillus paracasei* 8700:2 on aspects of common cold infections in children attending day care: a randomised, double-blind, placebo-controlled clinical study

Irini Lazou Ahrén<sup>1</sup> · Anna Berggren<sup>1</sup> · Cristina Teixeira<sup>1</sup> · Titti Martinsson Niskanen<sup>1</sup> · Niklas Larsson<sup>1</sup>

Published online: 19 December 2019  
© The Author(s) 2019

**Correction to: European Journal of Nutrition**  
<https://doi.org/10.1007/s00394-019-02137-8>

The original version of this article unfortunately contained a mistake. A number in Table 6 has been corrected.

The percentage should be 10 (91) instead of 10 (0.9) in the second row of the column “Probi Defendum® (*n* = 44)”. The corrected Table 6 is given below.

**Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article’s Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article’s Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

**Table 6** Number of children with concomitant medication (PP population)

	Total ( <i>n</i> = 99)	Probi Defendum® ( <i>n</i> = 44)	Placebo ( <i>n</i> = 55)	<i>p</i> *
Subjects with any concomitant medication, <i>n</i> (%) <sup>a</sup>	35 (35.3)	11 (25.0)	24 (43.6)	0.06
RTI-related, <i>n</i> (%) <sup>b</sup>	28 (80.0)	10 (91)	18 (75.0)	NS
GI-related, <i>n</i> (%) <sup>b</sup>	2 (5.7)	0	2 (8.3)	NS
Other reasons, <i>n</i> (%) <sup>b</sup>	7 (20.0)	4 (36.4)	3 (12.5)	NS

<sup>a</sup>Percentage of all the subjects in the corresponding group

<sup>b</sup>Percentage of the subjects, in the corresponding group, with concomitant medication

\*Fischer’s exact test

The original article can be found online at <https://doi.org/10.1007/s00394-019-02137-8>.

✉ Irini Lazou Ahrén  
irini.lazouahren@probi.com

<sup>1</sup> Probi AB, Lund, Sweden